



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 04:17 pm GMT

PDB ID : 4UDT
Title : T cell receptor (TRAV22,TRBV7-9) structure
Authors : Rodstrom, K.E.J.; Regenthal, P.; Lindkvist-Petersson, K.
Deposited on : 2014-12-11
Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	206	
2	B	243	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4010 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T CELL RECEPTOR ALPHA CHAIN, T-CELL RECEPTOR ALPHA CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	1680	1050	289	334	7	0	13	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	CYS	THR	engineered mutation	UNP P01848

- Molecule 2 is a protein called PROTEIN TRBV7-9, T-CELL RECEPTOR BETA-2 CHAIN C REGION.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	241	2015	1263	361	384	7	0	13	0

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP A0A5A3
B	98	GLY	-	linker	UNP A0A5A3
B	99	GLY	-	linker	UNP A0A5A3
B	100	TYR	-	linker	UNP A0A5A3
B	101	GLU	-	linker	UNP A0A5A3
B	102	GLN	-	linker	UNP A0A5A3
B	103	TYR	-	linker	UNP A0A5A3
B	104	PHE	-	linker	UNP A0A5A3
B	105	GLY	-	linker	UNP A0A5A3
B	106	PRO	-	linker	UNP A0A5A3
B	107	GLY	-	linker	UNP A0A5A3
B	108	THR	-	linker	UNP A0A5A3
B	109	ARG	-	linker	UNP A0A5A3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	110	LEU	-	linker	UNP A0A5A3
B	111	THR	-	linker	UNP A0A5A3
B	112	VAL	-	linker	UNP A0A5A3
B	113	THR	-	linker	UNP A0A5A3
B	114	GLU	-	linker	UNP A0A5A3
B	128	VAL	GLU	engineered mutation	UNP A0A5B9
B	170	CYS	SER	engineered mutation	UNP A0A5B9
B	188	ALA	CYS	engineered mutation	UNP A0A5B9
B	202	ASP	ASN	engineered mutation	UNP A0A5B9

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	C O	0	0
			6	3 3		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	143	Total	O	0	0
			143	143		
4	B	166	Total	O	0	0
			166	166		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: T CELL RECEPTOR ALPHA CHAIN, T-CELL RECEPTOR ALPHA CHAIN C REGION

Chain A: 



- Molecule 2: PROTEIN TRBV7-9, T-CELL RECEPTOR BETA-2 CHAIN C REGION

Chain B: 



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	39.07Å 79.75Å 69.03Å 90.00° 105.86° 90.00°	Depositor
Resolution (Å)	39.91 – 1.35 39.87 – 1.35	Depositor EDS
% Data completeness (in resolution range)	96.1 (39.91-1.35) 96.1 (39.87-1.35)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.35Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.156 , 0.193 0.165 , 0.203	Depositor DCC
R_{free} test set	4341 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	18.2	Xtrriage
Anisotropy	0.276	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4010	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1717	0.57	0/2335
2	B	0.35	0/2068	0.59	0/2812
All	All	0.35	0/3785	0.58	0/5147

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1680	0	1579	10	0
2	B	2015	0	1881	14	0
3	A	6	0	8	0	0
4	A	143	0	0	1	0
4	B	166	0	0	0	0
All	All	4010	0	3468	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:31[B]:ASN:OD1	2:B:32[B]:ARG:NH1	1.93	0.99
1:A:45[B]:ILE:O	1:A:45[B]:ILE:HD13	1.70	0.91
2:B:31[B]:ASN:CG	2:B:32[B]:ARG:HH11	1.81	0.83
2:B:14[A]:THR:HG21	2:B:20:VAL:CG1	2.09	0.82
2:B:31[B]:ASN:CG	2:B:32[B]:ARG:NH1	2.34	0.81
1:A:34:GLN:HE22	2:B:102:GLN:H	1.34	0.76
2:B:14[A]:THR:HG21	2:B:20:VAL:HG11	1.75	0.68
2:B:14[A]:THR:HG21	2:B:20:VAL:HG12	1.76	0.68
1:A:46[B]:ASN:HD22	1:A:46[B]:ASN:N	1.98	0.61
1:A:109[B]:LYS:HE3	1:A:109[B]:LYS:HA	1.83	0.59
1:A:54:THR:HG22	1:A:63[B]:THR:HG22	1.85	0.56
2:B:12:LYS:HE3	2:B:14[A]:THR:CG2	2.35	0.56
1:A:45[B]:ILE:HD13	1:A:45[B]:ILE:C	2.28	0.52
2:B:32[A]:ARG:NE	2:B:49:TYR:OH	2.43	0.52
2:B:47:LEU:HD22	2:B:62:LEU:HD22	1.91	0.51
1:A:148:GLN:NE2	4:A:2110:HOH:O	2.46	0.46
1:A:38:GLN:HA	1:A:43:GLN:O	2.15	0.45
1:A:163:ASP:HB3	1:A:165[B]:ARG:HE	1.81	0.45
2:B:145[A]:LEU:HG	2:B:147:THR:HG23	1.99	0.44
1:A:3:ILE:HD13	1:A:28[A]:ASP:OD1	2.18	0.44
2:B:31[B]:ASN:OD1	2:B:32[B]:ARG:HG2	2.20	0.42
2:B:190[A]:SER:OG	2:B:192:ARG:NH1	2.54	0.40
2:B:31[B]:ASN:ND2	2:B:32[B]:ARG:NH1	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/206 (103%)	207 (97%)	6 (3%)	0	100	100
2	B	252/243 (104%)	245 (97%)	7 (3%)	0	100	100
All	All	465/449 (104%)	452 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/185 (102%)	188 (100%)	1 (0%)	88	74
2	B	215/213 (101%)	213 (99%)	2 (1%)	78	53
All	All	404/398 (102%)	401 (99%)	3 (1%)	81	64

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	187	ASN
2	B	241	ARG
2	B	243	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	34	GLN
1	A	38	GLN
1	A	148	GLN
1	A	172	ASN
1	A	187	ASN
2	B	136	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	1204	-	5,5,5	0.30	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1204	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1204	GOL	C1-C2-C3-O3
3	A	1204	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/206 (98%)	0.49	21 (10%) 6 7	13, 23, 49, 77	0
2	B	241/243 (99%)	0.13	14 (5%) 23 25	12, 20, 45, 58	0
All	All	443/449 (98%)	0.29	35 (7%) 12 13	12, 22, 48, 77	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	203	PRO	21.9
1	A	41[A]	TRP	7.3
2	B	62	LEU	6.8
1	A	130	SER	6.1
1	A	93	ALA	6.0
2	B	41	GLY	5.9
2	B	61	LEU	5.2
2	B	40	LEU	4.7
1	A	98	TYR	4.5
1	A	182	ASP	4.3
2	B	183	ASN	4.0
1	A	42	GLY	4.0
1	A	96	GLY	3.9
1	A	181	SER	3.4
1	A	179	ASN	3.3
1	A	95	SER	3.3
2	B	181	ALA	3.0
2	B	219	ASN	3.0
2	B	3	THR	2.9
1	A	2	GLY	2.9
2	B	182	LEU	2.7
1	A	92	SER	2.7
1	A	94	THR	2.6
1	A	57	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	45[A]	ILE	2.5
1	A	202[A]	SER	2.4
1	A	43	GLN	2.4
2	B	39	THR	2.3
1	A	40	PRO	2.3
1	A	165[A]	ARG	2.2
2	B	221	GLU	2.2
2	B	209	CYS	2.1
2	B	223	THR	2.1
2	B	63	SER	2.1
1	A	97	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1204	6/6	0.78	0.11	41,43,44,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.