



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 16, 2023 – 12:32 pm GMT

PDB ID : 4UEW
Title : Structure of H₂-treated anaerobically purified *D. fructosovorans* NiFe- hydrogenase
Authors : Volbeda, A.; Martin, L.; Liebgott, P.-P.; Fontecilla-Camps, J.C.
Deposited on : 2014-12-20
Resolution : 2.08 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	 89% 9%
1	B	264	 3% 91% 8%
1	C	264	 91% 8%
2	Q	549	 91% 8%
2	R	549	 % 93% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	S	549	 A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a small red segment on the left labeled '5%', a large green segment in the middle labeled '91%', and a small yellow segment on the right labeled '8%'. The total length of the bar represents 100%.

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	Q	1564	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 20352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PERIPLASMIC [NIFE] HYDROGENASE SMALL SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	262	Total 1973	C 1256	N 330	O 372	S 15	0	0	0
1	B	262	Total 1980	C 1260	N 330	O 375	S 15	0	2	0
1	C	260	Total 1964	C 1251	N 328	O 370	S 15	0	1	0

- Molecule 2 is a protein called PERIPLASMIC [NIFE] HYDROGENASE LARGE SUBUNIT.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	Q	544	Total 4204	C 2680	N 728	O 774	S 22	0	9	0
2	R	545	Total 4181	C 2664	N 724	O 771	S 22	0	3	0
2	S	544	Total 4169	C 2654	N 724	O 769	S 22	0	1	0

- Molecule 3 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Fe S 8 4 4	0	0
3	A	1	Total Fe S 8 4 4	0	0
3	B	1	Total Fe S 8 4 4	0	0
3	B	1	Total Fe S 8 4 4	0	0
3	C	1	Total Fe S 8 4 4	0	0
3	C	1	Total Fe S 8 4 4	0	0

- Molecule 4 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe₃S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Fe	S	0	0
			7	3	4		
4	B	1	Total	Fe	S	0	0
			7	3	4		
4	C	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 6	C 3	O 3	0	0
5	A	1	Total 6	C 3	O 3	0	0
5	Q	1	Total 6	C 3	O 3	0	0
5	Q	1	Total 6	C 3	O 3	0	0
5	Q	1	Total 6	C 3	O 3	0	0
5	Q	1	Total 6	C 3	O 3	0	0
5	R	1	Total 6	C 3	O 3	0	0
5	R	1	Total 6	C 3	O 3	0	0
5	R	1	Total 6	C 3	O 3	0	0
5	R	1	Total 6	C 3	O 3	0	0
5	R	1	Total 6	C 3	O 3	0	0
5	R	1	Total 6	C 3	O 3	0	0
5	S	1	Total 6	C 3	O 3	0	0

- Molecule 6 is CARBONMONOXIDE-(DICYANO) IRON (three-letter code: FCO) (formula: C₃FeN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	Q	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
6	R	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		
6	S	1	Total	C	Fe	N	O	0	0
			7	3	1	2	1		

- Molecule 7 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	Q	1	Total	Ni	0	0
			1	1		
7	R	1	Total	Ni	0	0
			1	1		
7	S	1	Total	Ni	0	0
			1	1		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	Q	1	Total	Mg	0	0
			1	1		
8	R	1	Total	Mg	0	0
			1	1		
8	S	1	Total	Mg	0	0
			1	1		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	310	Total 310	O 310	0	0
9	B	191	Total 191	O 191	0	0
9	C	216	Total 216	O 216	0	0
9	Q	405	Total 405	O 405	0	0
9	R	342	Total 342	O 342	0	0
9	S	237	Total 237	O 237	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PERIPLASMIC [NIFE] HYDROGENASE SMALL SUBUNIT

Chain A: 



- Molecule 1: PERIPLASMIC [NIFE] HYDROGENASE SMALL SUBUNIT

Chain B: 



- Molecule 1: PERIPLASMIC [NIFE] HYDROGENASE SMALL SUBUNIT

Chain C: 





- Molecule 2: PERIPLASMIC [NIFE] HYDROGENASE LARGE SUBUNIT

Chain Q: 





• Molecule 2: PERIPLASMIC [NIFE] HYDROGENASE LARGE SUBUNIT



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.77Å 100.40Å 183.43Å 90.00° 91.55° 90.00°	Depositor
Resolution (Å)	21.98 – 2.08 21.98 – 2.08	Depositor EDS
% Data completeness (in resolution range)	87.6 (21.98-2.08) 87.7 (21.98-2.08)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.18 (at 2.08Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.195 , 0.235 0.195 , 0.235	Depositor DCC
R_{free} test set	6255 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.537	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	0.076 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	20352	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NI, SF4, MG, FCO, F3S, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.74	4/2027 (0.2%)	0.75	4/2759 (0.1%)
1	B	0.53	1/2043 (0.0%)	0.61	0/2781
1	C	0.68	8/2023 (0.4%)	0.66	2/2754 (0.1%)
2	Q	0.58	2/4334 (0.0%)	0.68	3/5881 (0.1%)
2	R	0.50	2/4292 (0.0%)	0.64	3/5825 (0.1%)
2	S	0.43	1/4271 (0.0%)	0.60	3/5796 (0.1%)
All	All	0.56	18/18990 (0.1%)	0.65	15/25796 (0.1%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	517	GLU	CD-OE2	-10.55	1.14	1.25
1	C	261	TYR	CE1-CZ	-10.53	1.24	1.38
2	Q	464	GLU	CD-OE1	-9.32	1.15	1.25
1	C	261	TYR	CG-CD2	-9.06	1.27	1.39
1	C	261	TYR	CG-CD1	-8.65	1.27	1.39

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	517	GLU	OE1-CD-OE2	-9.88	111.45	123.30
2	Q	464	GLU	OE1-CD-OE2	-7.94	113.77	123.30
2	S	161	LYS	CD-CE-NZ	7.87	129.79	111.70
1	A	203	ASP	CB-CG-OD2	7.55	125.09	118.30
1	C	215	GLU	OE1-CD-OE2	-7.03	114.86	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1973	0	1911	13	0
1	B	1980	0	1918	21	0
1	C	1964	0	1906	19	0
2	Q	4204	0	4192	31	0
2	R	4181	0	4154	24	0
2	S	4169	0	4140	28	0
3	A	16	0	0	0	0
3	B	16	0	0	0	0
3	C	16	0	0	0	0
4	A	7	0	0	0	0
4	B	7	0	0	0	0
4	C	7	0	0	0	0
5	A	18	0	24	3	0
5	Q	24	0	32	5	0
5	R	36	0	48	3	0
5	S	6	0	8	0	0
6	Q	7	0	0	1	0
6	R	7	0	0	0	0
6	S	7	0	0	1	0
7	Q	1	0	0	0	0
7	R	1	0	0	0	0
7	S	1	0	0	0	0
8	Q	1	0	0	0	0
8	R	1	0	0	0	0
8	S	1	0	0	0	0
9	A	310	0	0	4	0
9	B	191	0	0	2	0
9	C	216	0	0	5	0
9	Q	405	0	0	2	0
9	R	342	0	0	4	0
9	S	237	0	0	5	0
All	All	20352	0	18333	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:202:PHE:CE2	2:R:233:VAL:HG11	1.83	1.13
1:B:202:PHE:CE1	1:B:216:LEU:HD13	1.87	1.09
1:B:202:PHE:CD2	2:R:233:VAL:HG11	1.95	1.00
9:C:2216:HOH:O	2:Q:189:LYS:HE3	1.73	0.88
2:Q:337:LYS:HE2	5:Q:1564:GOL:H2	1.56	0.85

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/264 (98%)	254 (98%)	6 (2%)	0	100	100
1	B	262/264 (99%)	254 (97%)	8 (3%)	0	100	100
1	C	259/264 (98%)	251 (97%)	8 (3%)	0	100	100
2	Q	550/549 (100%)	533 (97%)	17 (3%)	0	100	100
2	R	545/549 (99%)	534 (98%)	11 (2%)	0	100	100
2	S	542/549 (99%)	524 (97%)	18 (3%)	0	100	100
All	All	2418/2439 (99%)	2350 (97%)	68 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/210 (99%)	206 (99%)	2 (1%)	76	81
1	B	210/210 (100%)	209 (100%)	1 (0%)	88	92
1	C	208/210 (99%)	208 (100%)	0	100	100
2	Q	443/438 (101%)	438 (99%)	5 (1%)	73	78
2	R	437/438 (100%)	432 (99%)	5 (1%)	73	78
2	S	435/438 (99%)	429 (99%)	6 (1%)	67	72
All	All	1941/1944 (100%)	1922 (99%)	19 (1%)	76	81

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	S	397	GLN
2	S	416	THR
2	S	473	ASP
2	S	408	LEU
2	R	92	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	B	14	ASN
1	B	61	HIS
1	C	14	ASN
1	C	172	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CSO	Q	543	7,2	3,6,7	0.54	0	0,6,8	-	-
2	CSO	S	543	7,2	3,6,7	0.83	0	0,6,8	-	-
2	CSO	R	543	7,2	3,6,7	0.66	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CSO	Q	543	7,2	-	1/1/5/7	-
2	CSO	S	543	7,2	-	1/1/5/7	-
2	CSO	R	543	7,2	-	1/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	Q	543	CSO	N-CA-CB-SG
2	R	543	CSO	N-CA-CB-SG
2	S	543	CSO	N-CA-CB-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 6 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	F3S	A	1266	1	0,9,9	-	-	-		
4	F3S	B	1266	1	0,9,9	-	-	-		
5	GOL	R	1565	-	5,5,5	0.32	0	5,5,5	0.20	0
6	FCO	Q	1550	7,2	0,6,6	-	-	-		
5	GOL	R	1564	-	5,5,5	0.33	0	5,5,5	0.18	0
5	GOL	R	1563	-	5,5,5	0.26	0	5,5,5	0.43	0
3	SF4	C	1265	1	0,12,12	-	-	-		
4	F3S	C	1266	1	0,9,9	-	-	-		
5	GOL	S	1561	-	5,5,5	0.38	0	5,5,5	0.24	0
5	GOL	Q	1563	-	5,5,5	0.27	0	5,5,5	0.26	0
3	SF4	C	1267	1	0,12,12	-	-	-		
5	GOL	R	1562	-	5,5,5	0.26	0	5,5,5	0.18	0
5	GOL	R	1566	-	5,5,5	0.36	0	5,5,5	0.27	0
3	SF4	A	1265	1	0,12,12	-	-	-		
5	GOL	Q	1562	-	5,5,5	0.20	0	5,5,5	0.30	0
5	GOL	Q	1561	-	5,5,5	0.26	0	5,5,5	0.18	0
5	GOL	A	1271	-	5,5,5	0.30	0	5,5,5	0.55	0
3	SF4	B	1265	1	0,12,12	-	-	-		
5	GOL	A	1273	-	5,5,5	0.39	0	5,5,5	0.54	0
6	FCO	S	1550	7,2	0,6,6	-	-	-		
3	SF4	A	1267	1	0,12,12	-	-	-		
3	SF4	B	1267	1	0,12,12	-	-	-		
5	GOL	Q	1564	-	5,5,5	0.34	0	5,5,5	0.46	0
6	FCO	R	1550	7,2	0,6,6	-	-	-		
5	GOL	R	1561	-	5,5,5	0.31	0	5,5,5	0.61	0
5	GOL	A	1272	-	5,5,5	0.33	0	5,5,5	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	R	1565	-	-	4/4/4/4	-
4	F3S	A	1266	1	-	-	0/3/3/3
4	F3S	B	1266	1	-	-	0/3/3/3
5	GOL	R	1564	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	R	1563	-	-	0/4/4/4	-
3	SF4	C	1265	1	-	-	0/6/5/5
5	GOL	S	1561	-	-	0/4/4/4	-
4	F3S	C	1266	1	-	-	0/3/3/3
5	GOL	Q	1563	-	-	0/4/4/4	-
3	SF4	C	1267	1	-	-	0/6/5/5
5	GOL	R	1562	-	-	4/4/4/4	-
5	GOL	R	1566	-	-	4/4/4/4	-
5	GOL	Q	1562	-	-	0/4/4/4	-
3	SF4	A	1265	1	-	-	0/6/5/5
5	GOL	Q	1561	-	-	0/4/4/4	-
5	GOL	A	1273	-	-	2/4/4/4	-
5	GOL	A	1271	-	-	3/4/4/4	-
3	SF4	B	1265	1	-	-	0/6/5/5
3	SF4	A	1267	1	-	-	0/6/5/5
3	SF4	B	1267	1	-	-	0/6/5/5
5	GOL	Q	1564	-	-	2/4/4/4	-
5	GOL	R	1561	-	-	1/4/4/4	-
5	GOL	A	1272	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1271	GOL	C1-C2-C3-O3
5	R	1562	GOL	O1-C1-C2-C3
5	R	1566	GOL	O1-C1-C2-O2
5	R	1566	GOL	O1-C1-C2-C3
5	R	1566	GOL	C1-C2-C3-O3

There are no ring outliers.

7 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	Q	1550	FCO	1	0
5	R	1564	GOL	1	0
5	Q	1563	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	R	1562	GOL	2	0
5	A	1273	GOL	3	0
6	S	1550	FCO	1	0
5	Q	1564	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	262/264 (99%)	-0.56	0 100 100	9, 17, 27, 46	4 (1%)
1	B	262/264 (99%)	0.17	9 (3%) 45 50	21, 36, 57, 71	6 (2%)
1	C	260/264 (98%)	-0.05	0 100 100	19, 33, 56, 70	4 (1%)
2	Q	543/549 (98%)	-0.52	0 100 100	9, 19, 33, 42	7 (1%)
2	R	544/549 (99%)	-0.19	4 (0%) 87 89	16, 29, 47, 65	6 (1%)
2	S	543/549 (98%)	0.41	25 (4%) 32 36	20, 48, 77, 90	7 (1%)
All	All	2414/2439 (98%)	-0.12	38 (1%) 72 75	9, 29, 63, 90	34 (1%)

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	264	GLY	5.5
2	S	400	PHE	4.2
2	S	280	TRP	3.9
2	S	160	LEU	3.9
2	S	134	LEU	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CSO	S	543	7/8	0.97	0.09	29,31,32,33	1
2	CSO	R	543	7/8	0.99	0.06	18,18,19,19	1
2	CSO	Q	543	7/8	0.99	0.06	9,9,10,10	1

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	GOL	R	1566	6/6	0.67	0.25	55,60,61,62	0
5	GOL	R	1562	6/6	0.83	0.19	49,50,52,53	0
5	GOL	Q	1563	6/6	0.83	0.20	43,49,53,53	0
5	GOL	Q	1564	6/6	0.85	0.20	36,38,41,46	0
5	GOL	S	1561	6/6	0.85	0.15	34,40,41,42	0
5	GOL	R	1563	6/6	0.89	0.15	34,36,38,39	0
5	GOL	A	1273	6/6	0.91	0.17	43,43,46,47	0
5	GOL	R	1564	6/6	0.91	0.20	40,43,47,53	0
5	GOL	R	1565	6/6	0.93	0.13	40,41,42,42	0
5	GOL	Q	1561	6/6	0.94	0.16	35,36,38,39	0
5	GOL	A	1272	6/6	0.95	0.16	39,39,42,46	0
8	MG	S	1553	1/1	0.95	0.05	34,34,34,34	0
3	SF4	B	1265	8/8	0.96	0.07	45,47,48,50	0
5	GOL	R	1561	6/6	0.96	0.08	28,29,30,30	0
5	GOL	A	1271	6/6	0.96	0.11	25,30,32,36	0
8	MG	R	1553	1/1	0.97	0.03	24,24,24,24	0
5	GOL	Q	1562	6/6	0.98	0.11	20,22,23,25	0
3	SF4	B	1267	8/8	0.98	0.06	24,25,26,26	0
4	F3S	B	1266	7/7	0.99	0.04	32,32,34,34	0
4	F3S	C	1266	7/7	0.99	0.05	20,21,21,22	0
3	SF4	A	1267	8/8	0.99	0.07	8,8,9,9	0
3	SF4	C	1265	8/8	0.99	0.07	19,21,21,22	0
6	FCO	R	1550	7/7	0.99	0.08	19,20,20,21	0
6	FCO	S	1550	7/7	0.99	0.12	29,30,31,32	0
7	NI	S	1551	1/1	0.99	0.08	27,27,27,27	0
8	MG	Q	1553	1/1	0.99	0.06	9,9,9,9	0
3	SF4	C	1267	8/8	0.99	0.07	23,23,24,25	0
4	F3S	A	1266	7/7	0.99	0.06	9,10,10,10	0
3	SF4	A	1265	8/8	1.00	0.06	13,14,15,15	0
6	FCO	Q	1550	7/7	1.00	0.07	9,9,9,9	0
7	NI	Q	1551	1/1	1.00	0.07	10,10,10,10	0
7	NI	R	1551	1/1	1.00	0.06	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.