

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 15, 2023 – 03:43 PM EDT

PDB ID	:	7UFK
Title	:	Crystal structure of chimeric omicron RBD (strain BA.2) complexed with hu-
		man ACE2
Authors	:	Zhang, W.; Shi, K.; Geng, Q.; Ye, G.; Aihara, H.; Li, F.
Deposited on	:	2022-03-22
Resolution	:	2.38  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	4.02b-467
:	1.8.5 (274361), CSD as541be (2020)
:	1.13
:	2.36
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	5.8.0158
:	7.0.044 (Gargrove)
:	Engh & Huber $(2001)$
:	Parkinson et al. (1996)
:	2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motrie	Whole archive	Similar resolution
	$(\# { m Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	5509(2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973(2.40-2.36)
Sidechain outliers	138945	5975(2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	597	5% 89%	10%
1	В	597	5% 88%	11%
2	Е	217	9% 77% 9%	13%
2	F	217	75% 11%	13%
3	С	2	100%	



Mol	Chain	Length	Quality of chain	
3	Ι	2	100%	
3	K	2	100%	
4	D	3	100%	
4	G	3	33% 67%	
5	J	4	75%	25%



#### 7UFK

## 2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 13092 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Angiotensin-converting enzyme 2.

Mol	Chain	Residues		Atoms					AltConf	Trace
1	А	596	Total 4862	C 3111	N 805	O 917	S 29	0	0	0
1	В	596	Total 4862	C 3111	N 805	O 917	S 29	0	0	0

• Molecule 2 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	188	Total	С	Ν	0	$\mathbf{S}$	0	1	0
			1504	968	249	277	10	0		
0	Б	100	Total	С	Ν	0	S	0	0	0
	Г	100	1501	966	249	277	9	0	0	0

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	С	2	Total         C         N         O           28         16         2         10	0	0	0
3	Ι	2	Total         C         N         O           28         16         2         10	0	0	0
3	K	2	Total         C         N         O           28         16         2         10	0	0	0

• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total 39	C 22	N 2	O 15	0	0	0
4	G	3	Total 39	C 22	N 2	O 15	0	0	0

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluco pyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	J	4	Total 50	C 28	N 2	O 20	0	0	0

• Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	Total Zn 1 1	0	0
6	В	1	Total Zn 1 1	0	0

• Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	Total Cl 1 1	0	0
7	В	1	Total Cl 1 1	0	0

• Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	А	1	Total         C         N         O           14         8         1         5	0	0
8	А	1	Total         C         N         O           14         8         1         5	0	0
8	А	1	Total         C         N         O           14         8         1         5	0	0
8	А	1	Total C N O 14 8 1 5	0	0
8	В	1	Total C N O 14 8 1 5	0	0
8	В	1	Total C N O 14 8 1 5	0	0
8	E	1	Total         C         N         O           14         8         1         5	0	0
8	F	1	Total C N O 14 8 1 5	0	0

• Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	А	1	Total Na 1 1	0	0

• Molecule 10 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
10	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
10	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	А	8	Total O 8 8	0	0
11	В	11	Total O 11 11	0	0
11	Ε	2	Total O 2 2	0	0
11	$\mathbf{F}$	1	Total O 1 1	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Angiotensin-converting enzyme 2



V483 A484 A484 A484 C485 C485 C485 F497 F615 E515 E515 E515 E515 E515 E515 C505 C505 C505 C505 C505 C505 C505 C	
• Molecule 2: Spike protein S1	
Chain F: 75% 11% 13%	
AKC VALL VALL PRO CLY VALL VALL VALL VALL VAL VALL VAL VALL VAL VA	
1468 1468 1472 1472 1472 1472 1472 1472 1472 1473 1500 1500 1500 1500 1500 1500 1500 150	
• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-opyranose	D-gluc
Chain C: 100%	
• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-opyranose	D-gluc
Chain I: 100%	
• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-opyranose	D-gluc
Chain K: 100%	
MAG3 MAG2	
• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose	4)-2-ac

Chain D:

100%



• Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 33%

67%



#### NAG1 NAG2 BMA3

 $\bullet \ Molecule \ 5: \ alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose$ 

Chain J: 75% 25%

NAG1 NAG2 BMA3 MAN<del>4</del>



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	80.40Å 117.86Å 113.02Å	Deneriten
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $92.13^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	29.01 - 2.38	Depositor
Resolution (A)	81.54 - 2.38	EDS
% Data completeness	51.4 (29.01-2.38)	Depositor
(in resolution range)	48.7(81.54-2.38)	EDS
R <sub>merge</sub>	0.04	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.82 (at 2.40 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
P.P.	0.192 , $0.235$	Depositor
$n, n_{free}$	0.192 , $0.233$	DCC
$R_{free}$ test set	2116 reflections $(4.90%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	60.7	Xtriage
Anisotropy	0.035	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.30 , 35.8	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
	0.007 for -h,-l,-k	
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
	0.027 for h,-k,-l	
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	13092	wwPDB-VP
Average B, all atoms $(Å^2)$	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BMA, NA, CL, ZN, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.23	0/4999	0.42	0/6792
1	В	0.23	0/4999	0.43	0/6792
2	Е	0.24	0/1546	0.49	0/2099
2	F	0.25	0/1543	0.49	0/2094
All	All	0.24	0/13087	0.44	0/17777

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4862	0	4632	32	0
1	В	4862	0	4633	34	0
2	Е	1504	0	1422	10	0
2	F	1501	0	1427	15	0
3	С	28	0	25	0	0
3	Ι	28	0	25	0	0
3	K	28	0	25	0	0
4	D	39	0	34	0	0
4	G	39	0	34	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	J	50	0	43	0	0
6	А	1	0	0	0	0
6	В	1	0	0	0	0
7	А	1	0	0	0	0
7	В	1	0	0	0	0
8	А	56	0	52	0	0
8	В	28	0	26	0	0
8	Ε	14	0	13	0	0
8	F	14	0	13	0	0
9	А	1	0	0	0	0
10	В	12	0	18	0	0
11	А	8	0	0	0	0
11	В	11	0	0	0	0
11	Е	2	0	0	0	0
11	F	1	0	0	0	0
All	All	13092	0	12422	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:245:ARG:NH2	1:A:603:PHE:O	2.29	0.66
1:B:55:THR:OG1	1:B:58:ASN:ND2	2.28	0.64
1:A:90:ASN:HB3	1:A:93:VAL:HG22	1.80	0.63
1:B:520:LEU:HD22	1:B:579:MET:HE2	1.81	0.62
1:B:177:ARG:NH1	1:B:181:GLU:OE2	2.33	0.62
1:A:177:ARG:HD3	1:A:498:CYS:HB2	1.83	0.60
1:A:520:LEU:HD22	1:A:579:MET:HE2	1.83	0.60
1:A:142:LEU:HB3	1:A:147:GLY:HA3	1.82	0.60
2:F:367:VAL:HG13	2:F:368:LEU:HD13	1.85	0.59
1:A:268:GLY:O	1:A:277:ASN:ND2	2.31	0.59
2:E:485:GLY:H	2:E:488:CYS:HB2	1.68	0.58
1:B:425:SER:OG	1:B:427:ASP:OD1	2.22	0.56
2:E:393:SER:O	2:E:523:THR:OG1	2.23	0.55
1:A:177:ARG:NH1	1:A:181:GLU:OE2	2.40	0.55
1:B:457:GLU:HG2	1:B:513:ILE:HB	1.90	0.54
1:B:233:ILE:HD13	1:B:450:LEU:HD13	1.90	0.54
1:A:524:GLN:HG2	1:A:583:PRO:HG2	1.90	0.53
2:F:401:VAL:HG22	2:F:509:ARG:HG2	1.91	0.53



	louo pugom	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:245:ARG:NH1	1:B:258:PRO:O	2.41	0.53
1:B:126:ILE:HA	1:B:129:THR:HG22	1.92	0.52
1:A:157:ASP:HB3	1:A:160:GLU:HB3	1.92	0.52
1:A:233:ILE:HD13	1:A:450:LEU:HD13	1.91	0.52
1:B:19:SER:HB2	1:B:24:GLN:HE21	1.74	0.51
1:A:294:THR:HG23	1:A:365:THR:HA	1.93	0.50
1:A:293:VAL:HB	1:A:423:LEU:HB3	1.93	0.50
2:E:401:VAL:HG22	2:E:509:ARG:HG2	1.94	0.50
2:F:454:ARG:NH2	2:F:469:SER:O	2.43	0.50
1:B:132:VAL:HG12	1:B:171:GLU:HG3	1.95	0.48
2:E:454:ARG:HD3	2:E:457:ARG:HB2	1.95	0.48
1:A:457:GLU:HG2	1:A:513:ILE:HB	1.95	0.48
1:B:157:ASP:HB3	1:B:160:GLU:HB3	1.95	0.48
1:B:241:HIS:CE1	1:B:245:ARG:HH21	2.31	0.48
1:A:394:ASN:OD1	1:A:395:GLY:N	2.47	0.48
1:A:108:LEU:HD23	1:A:112:LYS:HB3	1.95	0.47
1:B:100:LEU:HG	1:B:391:LEU:HD21	1.96	0.47
1:B:143:LEU:HD23	1:B:143:LEU:H	1.80	0.47
1:B:400:PHE:HZ	1:B:570:LEU:HB2	1.80	0.47
2:F:472:ILE:HD12	2:F:484:ALA:HB2	1.96	0.47
1:B:118:THR:O	1:B:122:THR:OG1	2.33	0.46
1:A:143:LEU:H	1:A:143:LEU:HD23	1.80	0.46
1:A:291:ILE:HG23	1:A:424:LEU:HD21	1.96	0.46
1:A:293:VAL:HG23	1:A:297:MET:HE2	1.97	0.46
1:B:245:ARG:NH2	1:B:605:GLY:O	2.31	0.46
1:A:446:ILE:HD13	1:A:523:PHE:HZ	1.81	0.46
2:F:395:VAL:HG23	2:F:524:VAL:HG21	1.98	0.46
1:B:291:ILE:H	1:B:291:ILE:HG13	1.64	0.45
1:B:394:ASN:OD1	1:B:395:GLY:N	2.49	0.45
2:E:472:ILE:HD12	2:E:484:ALA:HB2	1.99	0.45
1:B:305:GLN:O	1:B:309:LYS:HB2	2.16	0.45
1:B:455:MET:HE3	1:B:455:MET:HB3	1.89	0.45
2:E:350:VAL:HA	2:E:400:PHE:HB2	1.99	0.44
1:B:52:THR:HG22	1:B:359:LEU:HD13	1.99	0.44
1:B:538:PRO:HD2	1:B:541:LYS:HD3	2.00	0.44
1:B:578:ASN:OD1	1:B:579:MET:N	2.49	0.44
2:F:367:VAL:C	2:F:369:TYR:H	2.21	0.44
2:F:389:ASP:OD1	2:F:389:ASP:N	2.47	0.44
2:E:379:CYS:HA	2:E:432:CYS:HA	1.99	0.43
1:A:288:LYS:HE3	1:A:433:GLU:HB2	2.01	0.43
1:B:20:THR:HG22	1:B:22:GLU:H	1.83	0.43



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:B:247:LYS:HB2	1:B:282:THR:HG22	2.00	0.43
1:A:538:PRO:HD2	1:A:541:LYS:HD3	2.01	0.43
2:F:481:ASN:OD1	2:F:481:ASN:N	2.52	0.43
1:B:47:SER:O	1:B:51:ASN:ND2	2.46	0.43
1:A:20:THR:HG22	1:A:22:GLU:H	1.83	0.43
1:B:456:LEU:HD23	1:B:512:PHE:CD2	2.54	0.43
2:F:379:CYS:HA	2:F:432:CYS:HA	2.01	0.42
1:B:184:VAL:HG22	1:B:464:PHE:HE1	1.84	0.42
1:A:553:LYS:HE2	1:A:573:VAL:O	2.20	0.42
1:A:309:LYS:HD2	1:A:328:TRP:CH2	2.54	0.42
1:B:478:TRP:HA	1:B:481:LYS:HB2	2.02	0.42
2:E:350:VAL:HG22	2:E:422:ASN:HB3	2.02	0.42
2:F:497:PHE:CD2	2:F:507:PRO:HB3	2.55	0.42
1:B:180:TYR:HA	1:B:183:TYR:HB3	2.01	0.42
1:A:336:PRO:HG2	1:A:340:GLN:O	2.20	0.42
1:B:134:ASN:HD21	1:B:136:ASP:HB2	1.85	0.42
2:F:367:VAL:O	2:F:368:LEU:HB2	2.19	0.42
1:A:126:ILE:HA	1:A:129:THR:HG22	2.01	0.42
2:E:497:PHE:CD2	2:E:507:PRO:HB3	2.55	0.41
1:A:134:ASN:HD21	1:A:136:ASP:HB2	1.83	0.41
1:A:336:PRO:HB2	1:A:340:GLN:HB3	2.03	0.41
1:B:38:ASP:OD1	2:F:498:ARG:NH2	2.53	0.41
2:E:387:LEU:HA	2:E:390:LEU:HD12	2.03	0.41
1:A:468:ILE:HG22	1:A:473:TRP:HD1	1.86	0.41
2:F:350:VAL:HG22	2:F:422:ASN:HB3	2.02	0.41
1:A:456:LEU:HD23	1:A:512:PHE:CD2	2.55	0.41
1:A:288:LYS:HD3	1:A:434:THR:HG23	2.03	0.40
1:A:578:ASN:OD1	1:A:579:MET:N	2.51	0.40
1:B:116:LEU:HD11	1:B:187:LYS:HE2	2.01	0.40
2:F:502:GLY:O	2:F:506:GLN:HG3	2.22	0.40
2:F:390:LEU:HD23	2:F:390:LEU:HA	1.96	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	594/597~(100%)	574 (97%)	20 (3%)	0	100 100
1	В	594/597~(100%)	574 (97%)	20 (3%)	0	100 100
2	Ε	185/217~(85%)	173~(94%)	12~(6%)	0	100 100
2	F	184/217~(85%)	171 (93%)	13 (7%)	0	100 100
All	All	1557/1628~(96%)	1492 (96%)	65~(4%)	0	100 100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	526/527~(100%)	518~(98%)	8 (2%)	65 79
1	В	526/527~(100%)	517~(98%)	9(2%)	60 76
2	Ε	163/189~(86%)	162~(99%)	1 (1%)	86 93
2	F	163/189~(86%)	160 (98%)	3(2%)	59 75
All	All	1378/1432~(96%)	1357~(98%)	21 (2%)	65 79

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	79	LEU
1	А	143	LEU
1	А	172	VAL
1	А	297	MET
1	А	381	TYR
1	А	401	HIS
1	А	518	ARG
1	А	557	MET
1	В	31	LYS
1	В	122	THR



Mol	Chain	Res	Type
1	В	143	LEU
1	В	162	LEU
1	В	287	GLN
1	В	341	LYS
1	В	381	TYR
1	В	401	HIS
1	В	518	ARG
2	Е	377	PHE
2	F	377	PHE
2	F	468	ILE
2	F	481	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	417	HIS
1	В	24	GLN
1	В	58	ASN
2	F	477	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Type Chain		Chain Res J	Link	Bo	ond leng	$_{\rm ths}$	Bond angles		
IVIOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	С	1	1,3	14,14,15	0.37	0	$17,\!19,\!21$	0.61	0
3	NAG	С	2	3	14,14,15	0.27	0	$17,\!19,\!21$	0.38	0
4	NAG	D	1	4,1	14,14,15	0.29	0	$17,\!19,\!21$	0.45	0
4	NAG	D	2	4	14,14,15	0.22	0	17,19,21	0.40	0
4	BMA	D	3	4	11,11,12	0.61	0	$15,\!15,\!17$	0.76	0
4	NAG	G	1	4,1	14,14,15	0.94	1 (7%)	17,19,21	0.95	2 (11%)
4	NAG	G	2	4	14,14,15	0.16	0	$17,\!19,\!21$	0.73	1 (5%)
4	BMA	G	3	4	11,11,12	0.57	0	$15,\!15,\!17$	0.85	0
3	NAG	Ι	1	1,3	14,14,15	0.26	0	$17,\!19,\!21$	0.46	0
3	NAG	Ι	2	3	14,14,15	0.24	0	17,19,21	0.40	0
5	NAG	J	1	5,1	14,14,15	0.30	0	$17,\!19,\!21$	0.48	0
5	NAG	J	2	5	14,14,15	0.21	0	17,19,21	0.37	0
5	BMA	J	3	5	11,11,12	0.60	0	$15,\!15,\!17$	0.91	0
5	MAN	J	4	5	11,11,12	0.68	0	$15,\!15,\!17$	1.01	2(13%)
3	NAG	K	1	1,3	14,14,15	0.27	0	$17,\!19,\!21$	0.47	0
3	NAG	K	2	3	14,14,15	0.24	0	17,19,21	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	С	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	С	2	3	-	1/6/23/26	0/1/1/1
4	NAG	D	1	4,1	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	1/2/19/22	0/1/1/1
4	NAG	G	1	4,1	-	4/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
3	NAG	Ι	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	Ι	2	3	-	2/6/23/26	0/1/1/1
5	NAG	J	1	5,1	-	0/6/23/26	0/1/1/1
5	NAG	J	2	5	-	2/6/23/26	0/1/1/1
5	BMA	J	3	5	-	0/2/19/22	0/1/1/1
5	MAN	J	4	5	-	1/2/19/22	0/1/1/1
3	NAG	К	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	К	2	3	-	2/6/23/26	0/1/1/1



All	(1)	bond	length	outliers	are	listed	below:	
-----	-----	------	--------	----------	-----	--------	--------	--

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	G	1	NAG	O5-C1	-3.39	1.38	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	J	4	MAN	C1-O5-C5	2.39	115.43	112.19
4	G	1	NAG	C3-C4-C5	2.25	114.25	110.24
4	G	2	NAG	C1-O5-C5	2.22	115.19	112.19
5	J	4	MAN	O2-C2-C3	-2.14	105.85	110.14
4	G	1	NAG	O4-C4-C3	-2.02	105.67	110.35

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	Κ	1	NAG	O5-C5-C6-O6
4	G	2	NAG	O5-C5-C6-O6
3	Κ	2	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6
3	Ι	1	NAG	O5-C5-C6-O6
3	Κ	1	NAG	C4-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6
3	Κ	2	NAG	O5-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
4	G	1	NAG	C8-C7-N2-C2
4	G	1	NAG	O7-C7-N2-C2
3	Ι	2	NAG	O5-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
3	Ι	1	NAG	C4-C5-C6-O6
3	С	2	NAG	O5-C5-C6-O6
4	D	3	BMA	O5-C5-C6-O6
5	J	4	MAN	O5-C5-C6-O6
4	G	1	NAG	O5-C5-C6-O6
3	Ι	2	NAG	C4-C5-C6-O6
5	J	2	NAG	C4-C5-C6-O6
5	J	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,





bond angles, torsion angles, and ring geometry for oligosaccharide.

















## 5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 5 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	T in le	Bo	ond leng	ths	B	ond ang	les
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	NAG	А	703	1	14,14,15	0.24	0	17,19,21	0.43	0
8	NAG	А	705	1	14,14,15	0.19	0	17,19,21	0.45	0
10	EDO	В	704	-	3,3,3	0.46	0	2,2,2	0.34	0
10	EDO	В	703	-	3,3,3	0.46	0	2,2,2	0.34	0
8	NAG	А	704	1	14,14,15	0.22	0	17,19,21	0.41	0
8	NAG	А	706	1	14,14,15	0.23	0	17,19,21	0.45	0
8	NAG	F	601	2	14,14,15	0.22	0	17,19,21	0.44	0
8	NAG	В	705	1	14,14,15	0.24	0	17,19,21	0.47	0
10	EDO	В	707	-	3,3,3	0.46	0	2,2,2	0.34	0
8	NAG	Е	601	2	14,14,15	0.26	0	17,19,21	0.41	0
8	NAG	В	706	1	14,14,15	0.24	0	17,19,21	0.44	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	А	703	1	-	1/6/23/26	0/1/1/1
8	NAG	А	705	1	-	2/6/23/26	0/1/1/1
10	EDO	В	704	-	-	0/1/1/1	-
10	EDO	В	703	-	-	0/1/1/1	-
8	NAG	А	704	1	-	2/6/23/26	0/1/1/1
8	NAG	А	706	1	-	0/6/23/26	0/1/1/1
8	NAG	F	601	2	-	2/6/23/26	0/1/1/1
8	NAG	В	705	1	-	0/6/23/26	0/1/1/1
10	EDO	В	707	-	-	0/1/1/1	-
8	NAG	Е	601	2	-	1/6/23/26	0/1/1/1
8	NAG	B	706	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	А	704	NAG	O5-C5-C6-O6
8	А	705	NAG	O5-C5-C6-O6
8	А	704	NAG	C4-C5-C6-O6
8	F	601	NAG	O5-C5-C6-O6
8	А	705	NAG	C4-C5-C6-O6
8	F	601	NAG	C4-C5-C6-O6
8	Ε	601	NAG	O5-C5-C6-O6
8	А	703	NAG	O5-C5-C6-O6
8	В	706	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	596/597~(99%)	0.55	29 (4%) 29 32	40, 72, 122, 164	0
1	В	596/597~(99%)	0.58	32 (5%) 25 28	33, 70, 122, 192	0
2	Ε	188/217~(86%)	0.74	20 (10%) 6 7	42, 69, 132, 177	0
2	F	188/217~(86%)	0.70	21 (11%) 5 6	54, 88, 145, 167	0
All	All	1568/1628~(96%)	0.60	102 (6%) 18 20	33, 72, 127, 192	0

All (102) RSRZ outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	RSRZ
1	В	136	ASP	9.2
1	В	339	VAL	8.7
2	Е	369	TYR	7.9
1	В	108	LEU	6.8
2	F	363	ALA	6.1
1	В	110	GLU	5.9
2	Е	387	LEU	5.9
1	А	299	ASP	5.7
1	В	140	GLU	5.7
1	В	137	ASN	5.7
1	В	192	ARG	5.7
2	F	367	VAL	5.7
1	В	139	GLN	5.6
2	Е	368	LEU	5.6
1	А	424	LEU	5.5
2	F	368	LEU	5.3
2	F	456	PHE	5.3
1	А	423	LEU	5.3
2	Е	365	TYR	5.3
2	F	391	CYS	5.2
1	А	338	ASN	5.2



Mol	Chain	Res	Type	RSRZ
2	Е	386	LYS	5.2
2	F	390	LEU	5.1
1	А	360	MET	4.9
1	А	333	LEU	4.9
1	А	301	ALA	4.9
1	А	363	LYS	4.9
1	В	193	ALA	4.7
2	Е	384	ALA	4.7
2	Е	373	PHE	4.7
2	F	373	PHE	4.4
2	F	369	TYR	4.4
1	В	135	PRO	4.4
2	Е	515	PHE	4.2
2	F	364	ASP	4.1
1	В	195	HIS	4.0
2	F	387	LEU	3.9
2	Е	370	ASN	3.8
1	В	202	TYR	3.7
2	F	338	PHE	3.6
2	F	370	ASN	3.6
1	А	362	THR	3.5
1	В	85	LEU	3.4
2	F	335	LEU	3.4
1	А	302	TRP	3.4
2	F	392	PHE	3.4
2	Е	367	VAL	3.4
1	В	105	SER	3.4
1	В	97	LEU	3.3
1	В	141	CYS	3.3
1	А	289	PRO	3.2
1	В	191	ALA	3.1
2	Е	385	THR	3.1
1	В	190	MET	3.0
2	Е	517	LEU	2.9
1	А	293	VAL	2.9
1	А	85	LEU	2.9
2	Е	366	SER	2.9
1	В	107	VAL	2.9
2	Е	524	VAL	2.9
1	В	73	LEU	2.9
1	В	179	LEU	2.8
2	Е	392	PHE	2.7



	J	1	I J	
Mol	Chain	Res	Type	RSRZ
1	А	418	LEU	2.7
1	А	419	LYS	2.7
2	Е	394	ASN	2.7
1	А	84	PRO	2.7
1	А	369	PHE	2.7
1	А	297	MET	2.6
1	В	138	PRO	2.6
1	В	132	VAL	2.6
1	В	163	TRP	2.5
2	F	525	CYS	2.5
1	В	189	GLU	2.5
1	В	162	LEU	2.5
1	А	525	PHE	2.5
2	Е	483	VAL	2.5
1	В	88	ILE	2.5
1	В	172	VAL	2.4
1	А	300	GLN	2.4
2	F	366	SER	2.4
1	А	366	MET	2.4
2	Е	363	ALA	2.4
2	F	415	THR	2.3
2	F	341	VAL	2.3
2	Е	513	LEU	2.2
1	А	303	ASP	2.2
2	F	389	ASP	2.2
1	А	544	ILE	2.2
2	F	386	LYS	2.2
1	В	407	ILE	2.2
1	В	490	PRO	2.2
2	F	388	ASN	2.1
1	А	305	GLN	2.1
1	В	57	GLU	2.1
1	А	494	ASP	2.1
1	А	295	ASP	2.1
1	А	97	LEU	2.0
2	Е	341	VAL	2.0
1	А	91	LEU	2.0
1	А	140	GLU	2.0
1	В	129	THR	2.0

Continued from previous page...



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B$ -factors( $Å^2$ )	Q<0.9
4	BMA	D	3	11/12	0.45	0.18	102,128,136,137	0
4	BMA	G	3	11/12	0.61	0.22	125,142,161,161	0
3	NAG	K	2	14/15	0.74	0.23	95,131,141,143	0
3	NAG	Ι	2	14/15	0.79	0.30	84,129,143,145	0
5	MAN	J	4	11/12	0.84	0.18	91,104,116,130	0
5	BMA	J	3	11/12	0.87	0.12	83,101,104,111	0
3	NAG	С	2	14/15	0.88	0.39	100,116,124,134	0
4	NAG	G	1	14/15	0.88	0.25	111,121,151,155	0
4	NAG	G	2	14/15	0.88	0.19	106,122,136,138	0
3	NAG	Ι	1	14/15	0.91	0.16	92,103,127,133	0
3	NAG	K	1	14/15	0.92	0.12	59,100,121,123	0
4	NAG	D	2	14/15	0.92	0.15	95,122,146,149	0
3	NAG	С	1	14/15	0.93	0.20	80,92,102,117	0
5	NAG	J	2	14/15	0.93	0.16	53,66,86,94	0
4	NAG	D	1	14/15	0.95	0.12	72,82,96,98	0
5	NAG	J	1	14/15	0.96	0.15	42,54,77,80	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.















## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
8	NAG	А	704	14/15	0.70	0.23	88,132,150,151	0
8	NAG	В	705	14/15	0.81	0.23	53,102,123,123	0
8	NAG	А	703	14/15	0.82	0.13	72,113,120,125	0
9	NA	А	707	1/1	0.82	0.11	70,70,70,70	0
8	NAG	А	706	14/15	0.85	0.12	78,105,122,132	0
8	NAG	В	706	14/15	0.88	0.21	72,110,119,119	0
8	NAG	F	601	14/15	0.89	0.17	92,107,131,136	0
8	NAG	А	705	14/15	0.89	0.16	83,95,117,122	0
8	NAG	Е	601	14/15	0.90	0.22	82,106,126,132	0
10	EDO	В	704	4/4	0.93	0.15	52,66,74,74	0
10	EDO	В	703	4/4	0.94	0.18	$56,\!57,\!59,\!73$	0
7	CL	В	702	1/1	0.94	0.07	72,72,72,72	0
10	EDO	В	707	4/4	0.95	0.13	54,57,60,64	0
6	ZN	В	701	1/1	0.97	0.16	$65,\!65,\!65,\!65$	0
6	ZN	A	701	1/1	0.98	0.18	80,80,80,80	0
7	CL	А	702	1/1	0.99	0.09	70,70,70,70	0

### 6.5 Other polymers (i)

There are no such residues in this entry.

