



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2021 – 03:08 AM EDT

PDB ID : 1UI8
Title : Site-directed mutagenesis of His592 involved in binding of copper ion in *Arthrobacter globiformis* amine oxidase
Authors : Matsunami, H.; Okajima, T.; Hirota, S.; Yamaguchi, H.; Hori, H.; Kuroda, S.; Tanizawa, K.
Deposited on : 2003-07-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	638	 2% 87% 10% . .
1	B	638	 2% 88% 9% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylethylamine oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	620	4863	3071	853	930	9	0	0	0
1	B	620	4863	3071	853	930	9	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	382	TPQ	TYR	modified residue	UNP P46881
A	592	ALA	HIS	engineered mutation	UNP P46881
B	382	TPQ	TYR	modified residue	UNP P46881
B	592	ALA	HIS	engineered mutation	UNP P46881

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	B	1	Total	Cu	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	507	Total	O	0	0
			507	507		
3	B	600	Total	O	0	0
			600	600		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylethylamine oxidase



- Molecule 1: Phenylethylamine oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	157.76Å 63.21Å 184.24Å 90.00° 111.64° 90.00°	Depositor
Resolution (Å)	10.00 – 1.80 23.84 – 1.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.80) 99.4 (23.84-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.38 (at 1.80Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.217 , 0.258 0.185 , 0.214	Depositor DCC
R_{free} test set	7785 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtrriage
Anisotropy	0.865	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 63.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10835	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9116e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CU, TPQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/4969	0.72	0/6766
1	B	0.47	0/4969	0.73	0/6766
All	All	0.47	0/9938	0.72	0/13532

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4863	0	4683	41	0
1	B	4863	0	4682	34	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	507	0	0	3	0
3	B	600	0	0	5	0
All	All	10835	0	9365	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:SER:HB2	1:A:85:ILE:HD11	1.56	0.87
1:A:324:LEU:HB2	1:A:342:ILE:HB	1.70	0.72
1:B:166:SER:H	1:B:203:ASN:HD21	1.40	0.69
1:A:251:VAL:HG22	1:A:306:GLN:HB3	1.82	0.60
1:B:45:LEU:HD21	1:B:61:ARG:HB2	1.83	0.60
1:B:237:HIS:HE1	1:B:239:GLU:OE1	1.87	0.58
1:B:251:VAL:CG2	1:B:306:GLN:HB3	2.34	0.56
1:B:302:TYR:O	1:B:303:LEU:HB2	2.05	0.56
1:B:146:GLU:O	1:B:150:ARG:HD3	2.06	0.55
1:B:324:LEU:HB2	1:B:342:ILE:HB	1.88	0.54
1:A:72:PRO:HG2	1:A:90:LEU:HB2	1.90	0.54
1:B:599:TRP:CD2	1:B:600:PRO:HA	2.43	0.54
1:A:600:PRO:HD2	1:B:623:LEU:HD21	1.90	0.54
1:B:237:HIS:HD2	1:B:246:ASP:OD1	1.91	0.54
1:B:484:VAL:HG12	1:B:539:LYS:HG3	1.90	0.54
1:A:100:VAL:HG11	1:A:105:PHE:CE1	2.44	0.53
1:A:483:ALA:HB1	1:A:543:VAL:HB	1.91	0.53
1:A:302:TYR:O	1:A:303:LEU:HB2	2.11	0.51
1:B:154:ARG:HD2	1:B:293:TRP:CD2	2.47	0.50
1:A:251:VAL:CG2	1:A:306:GLN:HB3	2.41	0.50
1:A:267:ARG:NH2	3:A:1292:HOH:O	2.44	0.49
1:A:280:MET:HE3	1:A:384:TYR:CE1	2.48	0.49
1:A:162:PHE:HB2	1:A:163:PRO:HD2	1.95	0.49
1:B:479:ARG:O	1:B:482:GLU:HG2	2.13	0.48
1:A:24:GLN:HE22	1:A:334:ASN:ND2	2.11	0.48
1:B:515:HIS:HE1	3:B:1074:HOH:O	1.95	0.48
1:B:125:ARG:NH2	1:B:191:ASP:OD2	2.46	0.48
1:B:56:GLU:HB2	3:B:1334:HOH:O	2.13	0.47
1:A:189:VAL:HG12	1:A:189:VAL:O	2.14	0.47
1:A:187:THR:O	1:A:188:ARG:HG2	2.15	0.47
1:A:237:HIS:HD2	1:A:246:ASP:OD1	1.98	0.46
1:A:599:TRP:CD2	1:A:600:PRO:HA	2.50	0.46
1:B:251:VAL:HG22	1:B:306:GLN:HB3	1.95	0.46
1:A:484:VAL:HG12	1:A:539:LYS:HG3	1.97	0.46
1:B:384:TYR:OH	1:B:433:HIS:CE1	2.68	0.46
1:A:117:ARG:HB3	1:A:189:VAL:HG11	1.98	0.46
1:A:506:LEU:HD13	1:B:421:GLN:HG2	1.98	0.46
1:B:134:VAL:O	1:B:136:PRO:HD3	2.16	0.46
1:B:267:ARG:HG2	1:B:267:ARG:HH11	1.81	0.45
1:B:490:ARG:NE	3:B:1118:HOH:O	2.48	0.45
1:B:468:ASN:HD22	1:B:468:ASN:H	1.65	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:ARG:NE	1:A:85:ILE:HD13	2.31	0.45
1:A:26:ILE:HD11	1:A:82:GLY:HA2	2.00	0.44
1:B:125:ARG:HD3	3:B:1123:HOH:O	2.16	0.44
1:A:490:ARG:NE	3:A:1305:HOH:O	2.51	0.44
1:A:246:ASP:HB2	1:A:258:HIS:HB2	2.00	0.43
1:A:111:LEU:HD12	1:A:179:VAL:CG1	2.49	0.43
1:B:178:TYR:CD2	1:B:188:ARG:HD2	2.54	0.43
1:A:347:GLU:OE2	1:A:370:ARG:HD3	2.19	0.43
1:A:422:LEU:HD11	1:A:428:ALA:HB2	2.01	0.43
1:A:303:LEU:O	1:A:306:GLN:HG2	2.18	0.43
1:B:422:LEU:HD11	1:B:428:ALA:HB2	2.00	0.43
1:A:401:LYS:HG2	1:A:606:THR:HG22	2.00	0.42
1:A:513:LYS:O	1:A:611:LEU:HA	2.19	0.42
1:A:20:ILE:HD12	1:A:327:VAL:HG12	2.01	0.42
1:A:111:LEU:HD12	1:A:179:VAL:HG11	2.02	0.42
1:B:137:LEU:HD12	1:B:293:TRP:HH2	1.85	0.41
1:B:313:LEU:HD12	1:B:313:LEU:HA	1.93	0.41
1:B:298:ASP:HA	1:B:302:TYR:HD1	1.85	0.41
1:A:280:MET:HE3	1:A:384:TYR:CZ	2.56	0.41
1:A:537:ALA:HA	1:A:587:THR:HG21	2.02	0.41
1:A:505:ARG:HD3	1:A:618:ASP:HB3	2.02	0.41
1:A:10:SER:HA	1:A:11:PRO:HD3	1.91	0.41
1:B:10:SER:HB3	3:B:1467:HOH:O	2.20	0.41
1:B:125:ARG:HH22	1:B:191:ASP:CG	2.24	0.40
1:A:102:GLU:HA	1:A:105:PHE:CD2	2.56	0.40
1:A:242:LYS:NZ	3:A:1332:HOH:O	2.51	0.40
1:A:61:ARG:HH22	1:A:556:ASP:CG	2.24	0.40
1:A:143:GLU:CD	1:A:143:GLU:H	2.25	0.40
1:B:20:ILE:HD12	1:B:327:VAL:HG12	2.03	0.40
1:B:401:LYS:HG2	1:B:606:THR:HG22	2.02	0.40
1:B:510:VAL:HB	1:B:616:PHE:HA	2.04	0.40
1:A:45:LEU:CD1	1:A:61:ARG:HB2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	617/638 (97%)	590 (96%)	26 (4%)	1 (0%)	47	33
1	B	617/638 (97%)	594 (96%)	22 (4%)	1 (0%)	47	33
All	All	1234/1276 (97%)	1184 (96%)	48 (4%)	2 (0%)	47	33

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	54	GLU
1	B	55	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	512/528 (97%)	504 (98%)	8 (2%)	62	54
1	B	512/528 (97%)	505 (99%)	7 (1%)	67	59
All	All	1024/1056 (97%)	1009 (98%)	15 (2%)	65	56

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	LEU
1	A	143	GLU
1	A	208	GLU
1	A	303	LEU
1	A	311	LEU
1	A	370	ARG
1	A	376	PHE
1	A	541	LEU
1	B	208	GLU
1	B	303	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	336	ARG
1	B	339	ARG
1	B	376	PHE
1	B	444	ASP
1	B	468	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	HIS
1	A	273	ASN
1	A	334	ASN
1	A	507	ASN
1	A	515	HIS
1	B	203	ASN
1	B	237	HIS
1	B	309	ASN
1	B	334	ASN
1	B	433	HIS
1	B	468	ASN
1	B	515	HIS
1	B	519	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPQ	B	382	2,1	13,14,15	2.43	6 (46%)	15,19,21	1.41	3 (20%)
1	TPQ	A	382	2,1	13,14,15	2.56	8 (61%)	15,19,21	1.40	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPQ	B	382	2,1	-	2/5/22/24	0/1/1/1
1	TPQ	A	382	2,1	-	2/5/22/24	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	382	TPQ	C1-C2	-4.48	1.42	1.49
1	B	382	TPQ	C1-C2	-4.30	1.42	1.49
1	B	382	TPQ	O2-C2	4.16	1.35	1.24
1	A	382	TPQ	O2-C2	3.88	1.34	1.24
1	A	382	TPQ	O5-C5	3.67	1.34	1.24
1	B	382	TPQ	O5-C5	3.57	1.34	1.24
1	A	382	TPQ	C3-C4	3.08	1.40	1.35
1	B	382	TPQ	C3-C4	2.67	1.39	1.35
1	A	382	TPQ	C6-C5	-2.59	1.37	1.44
1	B	382	TPQ	C6-C1	2.57	1.41	1.34
1	B	382	TPQ	C4-C5	-2.51	1.39	1.47
1	A	382	TPQ	C4-C5	-2.49	1.39	1.47
1	A	382	TPQ	C3-C2	-2.27	1.38	1.44
1	A	382	TPQ	C6-C1	2.04	1.40	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	382	TPQ	O2-C2-C3	-2.62	115.78	121.78
1	B	382	TPQ	O2-C2-C3	-2.58	115.87	121.78
1	A	382	TPQ	CB-CA-C	-2.38	107.01	111.47
1	B	382	TPQ	CB-CA-C	-2.22	107.30	111.47
1	B	382	TPQ	O5-C5-C4	-2.06	116.01	119.38

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	382	TPQ	C-CA-CB-C1
1	B	382	TPQ	C-CA-CB-C1
1	A	382	TPQ	N-CA-CB-C1
1	B	382	TPQ	N-CA-CB-C1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	619/638 (97%)	-0.33	15 (2%) 59 54	6, 16, 34, 96	0
1	B	619/638 (97%)	-0.46	10 (1%) 72 68	3, 12, 28, 91	0
All	All	1238/1276 (97%)	-0.40	25 (2%) 65 61	3, 14, 32, 96	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	55	ALA	12.8
1	B	53	SER	10.1
1	B	52	GLY	9.5
1	B	54	GLU	8.3
1	A	52	GLY	7.4
1	A	53	SER	7.0
1	A	54	GLU	6.4
1	B	51	ALA	6.0
1	A	51	ALA	5.7
1	A	9	ALA	4.9
1	B	9	ALA	4.1
1	A	266	ASP	3.3
1	A	50	GLY	3.2
1	B	266	ASP	3.0
1	B	628	ASN	2.9
1	B	50	GLY	2.7
1	B	55	ALA	2.4
1	A	265	GLY	2.2
1	A	507	ASN	2.2
1	A	189	VAL	2.2
1	A	324	LEU	2.2
1	A	56	GLU	2.1
1	A	314	GLY	2.1
1	A	313	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	225	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPQ	A	382	14/15	0.84	0.14	12,33,38,40	0
1	TPQ	B	382	14/15	0.89	0.12	10,30,36,37	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU	B	1002	1/1	0.96	0.41	2,2,2,2	0
2	CU	A	1001	1/1	0.97	0.37	2,2,2,2	0

6.5 Other polymers [i](#)

There are no such residues in this entry.