



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 17, 2024 – 07:50 PM EST

PDB ID : 3UJF
Title : Asymmetric complex of human neuron specific enolase-4-PGA/PEP
Authors : Qin, J.; Chai, G.; Brewer, J.; Lovelace, L.; Lebioda, L.
Deposited on : 2011-11-07
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	 2% 77% 19% ..
1	B	443	 2% 80% 15% ..

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 6835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-enolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	433	3314	2084	568	649	13	0	0	0
1	B	432	3306	2078	567	648	13	0	0	0


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	434	HIS	-	expression tag	UNP P09104
A	435	HIS	-	expression tag	UNP P09104
A	436	HIS	-	expression tag	UNP P09104
A	437	HIS	-	expression tag	UNP P09104
A	438	HIS	-	expression tag	UNP P09104
A	439	HIS	-	expression tag	UNP P09104
A	440	HIS	-	expression tag	UNP P09104
A	441	HIS	-	expression tag	UNP P09104
A	442	HIS	-	expression tag	UNP P09104
A	443	HIS	-	expression tag	UNP P09104
B	434	HIS	-	expression tag	UNP P09104
B	435	HIS	-	expression tag	UNP P09104
B	436	HIS	-	expression tag	UNP P09104
B	437	HIS	-	expression tag	UNP P09104
B	438	HIS	-	expression tag	UNP P09104
B	439	HIS	-	expression tag	UNP P09104
B	440	HIS	-	expression tag	UNP P09104
B	441	HIS	-	expression tag	UNP P09104
B	442	HIS	-	expression tag	UNP P09104
B	443	HIS	-	expression tag	UNP P09104

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mg 2 2	0	0
2	B	2	Total Mg 2 2	0	0

- Molecule 3 is 2-PHOSPHOGLYCERIC ACID (three-letter code: 2PG) (formula: $C_3H_7O_7P$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O P 11 3 7 1	0	0

- Molecule 4 is PHOSPHOENOLPYRUVATE (three-letter code: PEP) (formula: $C_3H_5O_6P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	P	0	0
			10	3	6	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	88	Total	O	0	0
			88	88		
5	B	102	Total	O	0	0
			102	102		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gamma-enolase



- Molecule 1: Gamma-enolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	109.61Å 119.34Å 68.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.39 – 2.10 37.39 – 2.10	Depositor EDS
% Data completeness (in resolution range)	89.2 (37.39-2.10) 89.2 (37.39-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.75 (at 2.10Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.194 , 0.259 0.194 , 0.257	Depositor DCC
R_{free} test set	2380 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtrriage
Anisotropy	0.342	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6835	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2PG, MG, PEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.03	2/3369 (0.1%)	0.87	4/4558 (0.1%)
1	B	1.00	2/3361 (0.1%)	0.89	0/4547
All	All	1.02	4/6730 (0.1%)	0.88	4/9105 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	406	TYR	CD2-CE2	6.30	1.48	1.39
1	B	34	VAL	CB-CG2	6.16	1.65	1.52
1	A	61	VAL	CB-CG2	5.40	1.64	1.52
1	B	303	TRP	CE3-CZ3	5.09	1.47	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	LEU	CA-CB-CG	6.26	129.71	115.30
1	A	402	ARG	NE-CZ-NH1	5.51	123.06	120.30
1	A	29	LEU	CA-CB-CG	5.20	127.25	115.30
1	A	402	ARG	NE-CZ-NH2	-5.07	117.77	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3314	0	3289	59	1
1	B	3306	0	3278	49	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	11	0	4	2	0
4	B	10	0	2	0	0
5	A	88	0	0	2	0
5	B	102	0	0	1	1
All	All	6835	0	6573	101	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:334:LYS:HA	1:A:334:LYS:HE3	1.49	0.93
1:A:271:THR:OG1	1:A:274:GLN:HG3	1.70	0.91
1:A:215:ASN:H	1:A:215:ASN:HD22	1.21	0.86
1:A:101:ASN:H	1:A:101:ASN:HD22	1.24	0.84
1:B:141:ASP:O	1:B:388:CYS:SG	2.37	0.83
1:A:102:LYS:HE2	1:A:346:ILE:O	1.85	0.76
1:B:101:ASN:HD22	1:B:101:ASN:H	1.36	0.73
1:A:400:SER:HB2	1:B:401:GLU:HB3	1.71	0.72
1:B:238:LYS:HE3	5:B:616:HOH:O	1.92	0.70
1:A:205:ASN:OD1	5:A:621:HOH:O	2.10	0.68
1:A:102:LYS:HB2	5:A:620:HOH:O	1.94	0.67
1:A:215:ASN:H	1:A:215:ASN:ND2	1.92	0.66
1:A:401:GLU:HB3	1:B:400:SER:HB2	1.80	0.64
1:A:101:ASN:H	1:A:101:ASN:ND2	1.96	0.62
1:B:294:PRO:HD2	1:B:303:TRP:CH2	2.34	0.62
1:A:73:ALA:O	1:A:77:ILE:HG13	1.98	0.62
1:A:182:ARG:NH1	1:B:54:GLN:O	2.33	0.61
1:A:342:LYS:HZ3	3:A:503:2PG:H2	1.66	0.61
1:B:76:LEU:HD23	1:B:92:LEU:HD23	1.83	0.61
1:B:73:ALA:HB3	1:B:74:PRO:HD3	1.83	0.60
1:B:340:LEU:HD23	1:B:342:LYS:HE3	1.84	0.60
1:A:164:MET:HG2	1:A:245:VAL:HG13	1.85	0.59
1:B:322:THR:HG23	1:B:341:LEU:HD12	1.85	0.58
1:B:357:LYS:O	1:B:361:GLU:HB2	2.04	0.58
1:A:400:SER:CB	1:B:401:GLU:HB3	2.34	0.58
1:A:143:ILE:HD13	1:A:423:ALA:HB2	1.87	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:LEU:CD2	1:B:92:LEU:HD23	2.36	0.56
1:A:41:GLY:O	1:A:44:GLU:HG3	2.06	0.55
1:A:143:ILE:C	1:A:143:ILE:HD12	2.27	0.55
1:B:155:GLY:O	1:B:261:LYS:CE	2.55	0.55
1:A:313:GLN:HA	1:A:337:ASN:HD21	1.73	0.54
1:B:328:GLU:HG2	1:B:358:LEU:HD21	1.90	0.54
1:A:68:ILE:HD11	1:A:109:ALA:HA	1.90	0.54
1:B:251:TYR:OH	1:B:254:GLY:HA2	2.09	0.53
1:A:293:ASP:OD2	1:A:317:ASP:HB3	2.10	0.52
1:A:244:ASP:HA	1:A:292:GLU:HB3	1.92	0.51
1:A:245:VAL:HG11	1:A:279:TYR:OH	2.10	0.51
1:A:227:LYS:HE3	1:A:286:TYR:CD1	2.46	0.51
1:B:149:PHE:O	1:B:168:MET:HA	2.11	0.51
1:B:94:LEU:HD22	1:B:102:LYS:HE3	1.91	0.51
1:B:101:ASN:HD22	1:B:101:ASN:N	2.08	0.51
1:B:264:THR:HG22	1:B:265:ASP:N	2.27	0.50
1:A:337:ASN:HD22	1:A:337:ASN:C	2.15	0.50
1:A:359:ALA:O	1:A:364:TRP:HB2	2.12	0.49
1:B:293:ASP:HA	1:B:303:TRP:CH2	2.47	0.49
1:A:101:ASN:HD22	1:A:101:ASN:N	2.02	0.49
1:A:340:LEU:HD23	1:A:342:LYS:HE3	1.95	0.49
1:B:97:ASP:OD1	1:B:104:LYS:HB3	2.12	0.49
1:B:153:ASN:N	1:B:153:ASN:OD1	2.45	0.49
1:B:155:GLY:O	1:B:261:LYS:HD3	2.13	0.49
1:A:319:LEU:O	1:A:326:ARG:HD3	2.13	0.49
1:B:257:ASP:OD2	1:B:260:PHE:HA	2.12	0.49
1:A:293:ASP:HA	1:A:303:TRP:CH2	2.48	0.48
1:B:155:GLY:O	1:B:261:LYS:NZ	2.46	0.48
1:A:183:LEU:HB3	1:A:239:ILE:HD11	1.96	0.48
1:B:156:SER:OG	1:B:249:GLU:HB3	2.13	0.48
1:B:416:LEU:HB2	1:B:420:ALA:HB2	1.95	0.48
1:B:340:LEU:CD2	1:B:342:LYS:HE3	2.44	0.47
1:A:313:GLN:O	1:A:314:ILE:HD13	2.13	0.47
1:A:370:HIS:ND1	1:A:402:ARG:NH1	2.58	0.46
1:B:395:GLY:HA3	1:B:402:ARG:HD2	1.97	0.46
1:A:61:VAL:O	1:A:65:VAL:HG23	2.15	0.46
1:B:429:ASN:HD22	1:B:429:ASN:N	2.13	0.46
1:A:314:ILE:H	1:A:337:ASN:HD21	1.63	0.46
1:B:141:ASP:HB3	1:B:421:ARG:NH1	2.31	0.45
1:B:429:ASN:HD22	1:B:429:ASN:H	1.63	0.45
1:A:183:LEU:O	1:A:187:VAL:HG23	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:HIS:HE1	1:B:15:GLY:O	1.99	0.45
1:A:314:ILE:H	1:A:337:ASN:ND2	2.14	0.45
1:A:4:LYS:HG3	1:A:24:TYR:HB2	1.98	0.45
1:B:323:ASN:HA	1:B:324:PRO:HD2	1.67	0.45
1:A:334:LYS:HE3	1:A:334:LYS:CA	2.31	0.44
1:A:119:LYS:NZ	1:A:377:ASP:OD2	2.40	0.44
1:B:12:ASP:OD1	1:B:12:ASP:C	2.55	0.44
1:B:6:TRP:HA	1:B:6:TRP:CE3	2.53	0.44
1:B:29:LEU:HD12	1:B:30:PHE:N	2.33	0.43
1:A:257:ASP:HB2	1:A:269:TYR:CD1	2.53	0.43
1:A:44:GLU:CD	1:A:345:GLN:HG2	2.38	0.43
1:A:371:ARG:O	1:A:374:GLU:HG2	2.19	0.43
1:B:178:ARG:NE	1:B:413:GLU:OE1	2.50	0.42
1:B:279:TYR:O	1:B:283:VAL:HG23	2.19	0.42
1:A:401:GLU:HB3	1:B:400:SER:CB	2.48	0.42
1:A:342:LYS:NZ	3:A:503:2PG:H2	2.32	0.42
1:B:57:LEU:HD12	1:B:57:LEU:N	2.34	0.42
1:A:102:LYS:HD3	1:A:110:ILE:HD12	2.01	0.42
1:B:133:ILE:HD13	1:B:383:LEU:HA	2.01	0.42
1:A:141:ASP:O	1:A:388:CYS:SG	2.56	0.42
1:A:164:MET:H	1:A:219:ASN:HD21	1.68	0.42
1:B:108:ASN:OD1	1:B:108:ASN:N	2.52	0.41
1:B:394:THR:HG23	1:B:406:TYR:CZ	2.54	0.41
1:A:426:ASN:HB2	1:A:433:LEU:HD11	2.02	0.41
1:B:6:TRP:HA	1:B:6:TRP:HE3	1.85	0.41
1:A:143:ILE:HD11	1:A:390:GLY:HA2	2.01	0.41
1:A:265:ASP:HA	1:A:266:PRO:HD3	1.89	0.41
1:A:165:GLN:HB3	1:A:166:GLU:HG3	2.03	0.41
1:B:336:CYS:SG	1:B:364:TRP:CH2	3.13	0.41
1:A:292:GLU:OE1	1:A:340:LEU:HD22	2.21	0.41
1:A:284:ARG:HG3	1:A:285:ASP:N	2.37	0.40
1:A:92:LEU:HD12	1:A:92:LEU:HA	1.96	0.40
1:A:201:LYS:HE3	1:B:261:LYS:HB3	2.04	0.40
1:A:370:HIS:CG	1:A:394:THR:HA	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:SER:OG	5:B:606:HOH:O[3_645]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	431/443 (97%)	403 (94%)	28 (6%)	0	100	100
1	B	430/443 (97%)	408 (95%)	20 (5%)	2 (0%)	29	26
All	All	861/886 (97%)	811 (94%)	48 (6%)	2 (0%)	47	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	269	TYR
1	B	399	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	350/360 (97%)	329 (94%)	21 (6%)	19	16
1	B	349/360 (97%)	330 (95%)	19 (5%)	22	20
All	All	699/720 (97%)	659 (94%)	40 (6%)	20	18

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	54	GLN
1	A	101	ASN
1	A	150	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	215	ASN
1	A	237	GLU
1	A	264	THR
1	A	312	ILE
1	A	320	THR
1	A	325	LYS
1	A	337	ASN
1	A	344	ASN
1	A	357	LYS
1	A	370	HIS
1	A	372	SER
1	A	388	CYS
1	A	389	THR
1	A	391	GLN
1	A	403	LEU
1	A	418	ASP
1	A	419	GLU
1	B	6	TRP
1	B	29	LEU
1	B	97	ASP
1	B	101	ASN
1	B	141	ASP
1	B	152	ILE
1	B	153	ASN
1	B	178	ARG
1	B	215	ASN
1	B	273	ASP
1	B	309	ASN
1	B	370	HIS
1	B	388	CYS
1	B	389	THR
1	B	403	LEU
1	B	415	GLU
1	B	418	ASP
1	B	419	GLU
1	B	429	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	A	135	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	150	ASN
1	A	189	HIS
1	A	215	ASN
1	A	219	ASN
1	A	337	ASN
1	A	354	GLN
1	A	426	ASN
1	B	101	ASN
1	B	139	ASN
1	B	150	ASN
1	B	189	HIS
1	B	309	ASN
1	B	345	GLN
1	B	429	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2PG	A	503	2	9,10,10	1.18	1 (11%)	11,14,14	1.86	2 (18%)
4	PEP	B	503	2	9,9,9	1.70	2 (22%)	11,13,13	2.18	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2PG	A	503	2	-	3/11/11/11	-
4	PEP	B	503	2	-	0/9/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	503	PEP	O2'-C1	-3.43	1.20	1.30
4	B	503	PEP	O2-C2	2.54	1.46	1.39
3	A	503	2PG	O2-C1	-2.26	1.23	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	PEP	O2'-C1-C2	4.62	121.79	113.91
3	A	503	2PG	P-O1P-C2	-4.58	112.53	123.04
4	B	503	PEP	O2'-C1-O1	-3.28	116.10	123.61
4	B	503	PEP	O2P-P-O1P	2.31	119.72	110.68
4	B	503	PEP	O2P-P-O2	-2.22	98.49	105.25
4	B	503	PEP	O2-C2-C3	-2.20	120.55	124.79
3	A	503	2PG	O2-C1-C2	2.18	119.27	113.03

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	503	2PG	O2-C1-C2-C3
3	A	503	2PG	O1-C1-C2-C3
3	A	503	2PG	O2-C1-C2-O1P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	503	2PG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	433/443 (97%)	-0.17	10 (2%) 60 65	19, 33, 50, 60	0
1	B	432/443 (97%)	-0.14	7 (1%) 72 75	18, 33, 49, 64	5 (1%)
All	All	865/886 (97%)	-0.15	17 (1%) 65 69	18, 33, 50, 64	5 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	418	ASP	3.8
1	A	253	ASP	3.7
1	B	263	PRO	3.1
1	A	418	ASP	2.9
1	B	141	ASP	2.6
1	B	159	GLY	2.5
1	A	380	ILE	2.4
1	A	54	GLN	2.4
1	A	6	TRP	2.4
1	A	302	ALA	2.4
1	B	54	GLN	2.3
1	A	33	ALA	2.2
1	B	403	LEU	2.2
1	A	57	LEU	2.1
1	A	265	ASP	2.1
1	B	395	GLY	2.1
1	A	140	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	A	502	1/1	0.95	0.14	13,13,13,13	0
2	MG	A	501	1/1	0.97	0.22	10,10,10,10	0
2	MG	B	501	1/1	0.98	0.11	5,5,5,5	0
2	MG	B	502	1/1	0.98	0.12	9,9,9,9	0
3	2PG	A	503	11/11	0.98	0.11	26,29,36,40	0
4	PEP	B	503	10/10	0.99	0.12	22,24,27,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.