



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 2, 2023 – 10:24 AM EDT

PDB ID : 3UMJ
Title : Crystal Structure of D311E Lipase
Authors : Ruslan, R.; Rahman, R.N.Z.R.A.; Leow, T.C.; Ali, M.S.M.; Basri, M.; Salleh, A.B.
Deposited on : 2011-11-13
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	 2% 85% 13%
1	B	387	 88% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1001	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thermostable lipase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	386	Total	C	N	O	S	0	0	0
			3048	1930	539	571	8			
1	B	387	Total	C	N	O	S	0	0	0
			3054	1933	540	573	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	311	GLU	ASP	engineered mutation	UNP Q842J9
B	311	GLU	ASP	engineered mutation	UNP Q842J9

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0
4	B	1	Total Cl 1 1	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	B	1	Total Na 1 1	0	0

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Ca 1 1	0	0
6	B	1	Total Ca 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	322	Total O 322 322	0	0
7	B	300	Total O 300 300	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thermostable lipase



- Molecule 1: Thermostable lipase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.32Å 81.16Å 100.14Å 90.00° 96.49° 90.00°	Depositor
Resolution (Å)	33.17 – 2.10 33.17 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.8 (33.17-2.10) 96.8 (33.17-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.54 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.155 , 0.212 0.157 , 0.214	Depositor DCC
R_{free} test set	2677 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	11.0	Xtrriage
Anisotropy	0.144	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6750	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GOL, CA, NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.08	4/3132 (0.1%)	0.92	9/4258 (0.2%)
1	B	1.04	3/3138 (0.1%)	0.92	5/4266 (0.1%)
All	All	1.06	7/6270 (0.1%)	0.92	14/8524 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	30	TRP	CB-CG	5.58	1.60	1.50
1	B	90	PHE	CE2-CZ	5.46	1.47	1.37
1	A	258	ALA	CA-CB	5.40	1.63	1.52
1	A	123	VAL	CB-CG1	5.18	1.63	1.52
1	B	251	LYS	CD-CE	5.17	1.64	1.51
1	B	226	GLU	CG-CD	5.13	1.59	1.51
1	A	370	PHE	CD1-CE1	5.13	1.49	1.39

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	307	LEU	CA-CB-CG	8.31	134.41	115.30
1	B	285	LEU	CB-CG-CD2	-7.12	98.90	111.00
1	A	103	ARG	NE-CZ-NH2	-6.56	117.02	120.30
1	B	317	ASP	CB-CG-OD2	-6.33	112.60	118.30
1	A	24	MET	CG-SD-CE	-5.91	90.75	100.20
1	A	357	ASP	CB-CG-OD1	5.82	123.54	118.30
1	A	103	ARG	NE-CZ-NH1	5.80	123.20	120.30
1	B	317	ASP	CB-CG-OD1	5.52	123.27	118.30
1	A	34	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	A	89	ARG	NE-CZ-NH2	-5.47	117.57	120.30
1	B	230	ARG	NE-CZ-NH1	5.39	122.99	120.30
1	B	76	ASP	CB-CG-OD1	5.30	123.07	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	ARG	NE-CZ-NH2	5.20	122.90	120.30
1	A	330	ARG	NE-CZ-NH2	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3048	0	2912	44	0
1	B	3054	0	2916	30	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	12	0	16	12	3
3	B	6	0	8	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	322	0	0	5	9
7	B	300	0	0	9	1
All	All	6750	0	5852	72	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:GLY:N	3:A:1001:GOL:H11	1.36	1.38
1:A:32:GLY:O	3:A:1001:GOL:C1	1.96	1.14
1:A:32:GLY:O	3:A:1001:GOL:H12	1.46	1.13
1:A:35:GLY:HA2	3:A:1001:GOL:O2	1.58	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:GLY:N	3:A:1001:GOL:C1	2.23	1.02
1:A:35:GLY:H	3:A:1001:GOL:C1	1.73	1.00
1:B:47:ARG:NH1	7:B:658:HOH:O	1.96	0.98
1:B:214:ARG:HD2	7:B:662:HOH:O	1.66	0.94
1:A:32:GLY:O	3:A:1001:GOL:O1	1.86	0.92
1:B:250:GLU:HG3	7:B:609:HOH:O	1.73	0.89
1:A:304:ASN:OD1	1:A:306:THR:HG22	1.74	0.88
1:A:106:ARG:HH21	1:A:153:HIS:HE1	1.22	0.88
1:A:35:GLY:H	3:A:1001:GOL:H11	1.04	0.85
1:A:308:GLY:HA3	1:A:330:ARG:HH22	1.43	0.84
1:A:387:GLN:OE1	7:A:683:HOH:O	1.95	0.84
1:B:216:GLN:HA	1:B:216:GLN:OE1	1.79	0.81
1:B:24:MET:HE1	1:B:204:TYR:HB2	1.65	0.78
1:A:114:GLN:HE22	1:A:240:ALA:HB1	1.50	0.76
1:B:312:ARG:HD3	7:B:589:HOH:O	1.83	0.76
1:B:250:GLU:CG	7:B:609:HOH:O	2.33	0.73
1:A:35:GLY:O	3:A:1001:GOL:O1	2.08	0.70
1:A:272:THR:O	7:A:542:HOH:O	2.09	0.70
1:B:24:MET:CE	1:B:204:TYR:HB2	2.21	0.70
1:B:182:ASP:OD1	7:B:526:HOH:O	2.12	0.68
1:B:250:GLU:CD	7:B:609:HOH:O	2.33	0.65
1:A:35:GLY:CA	3:A:1001:GOL:H11	2.24	0.65
1:A:304:ASN:OD1	1:A:306:THR:CG2	2.45	0.64
1:A:307:LEU:HD23	1:A:307:LEU:O	1.98	0.64
1:A:35:GLY:CA	3:A:1001:GOL:C1	2.77	0.62
1:A:153:HIS:O	7:A:708:HOH:O	2.17	0.59
1:A:17:THR:HG22	1:A:184:GLN:NE2	2.19	0.58
1:A:273:TYR:CE2	1:B:236:SER:HB3	2.42	0.55
1:B:84:LYS:HE3	7:B:467:HOH:O	2.09	0.51
1:A:208:LEU:HA	1:A:210:GLN:HE22	1.76	0.51
1:B:242:TYR:OH	1:B:251:LYS:CD	2.59	0.51
1:A:106:ARG:HH21	1:A:153:HIS:CE1	2.14	0.50
1:B:210:GLN:HE21	1:B:210:GLN:N	2.10	0.50
1:B:250:GLU:OE1	1:B:254:GLN:NE2	2.44	0.50
1:A:273:TYR:CE2	1:B:236:SER:CB	2.95	0.49
1:A:16:PHE:HA	1:A:55:PRO:O	2.12	0.48
1:A:120:ARG:HD2	1:A:348:TRP:CZ2	2.50	0.47
1:A:317:ASP:O	1:A:357:ASP:HB2	2.14	0.47
1:A:210:GLN:HE21	1:A:210:GLN:N	2.13	0.46
1:B:19:TRP:CE2	1:B:24:MET:HG3	2.51	0.46
1:B:303:ARG:HH11	1:B:303:ARG:HG3	1.81	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:THR:HG22	1:A:184:GLN:HE22	1.81	0.45
1:B:39:GLN:NE2	1:B:43:ASP:OD1	2.48	0.45
1:B:242:TYR:OH	1:B:251:LYS:HD2	2.17	0.45
1:A:343:LEU:HD13	1:A:386:LEU:HD21	2.00	0.44
1:B:17:THR:HG22	7:B:550:HOH:O	2.17	0.44
1:B:175:ASP:OD1	1:B:175:ASP:N	2.39	0.44
1:B:273:TYR:CZ	1:B:282:TYR:HB2	2.53	0.44
1:A:33:VAL:HG22	1:A:363:GLY:O	2.17	0.44
1:A:35:GLY:HA2	3:A:1001:GOL:C2	2.43	0.43
1:B:82:ALA:HB1	1:B:87:HIS:O	2.19	0.43
1:A:343:LEU:CD1	1:A:386:LEU:HD21	2.49	0.43
1:B:208:LEU:HA	1:B:210:GLN:HE22	1.84	0.43
1:A:19:TRP:CE2	1:A:24:MET:HG3	2.54	0.42
1:B:274:ARG:NH2	1:B:311:GLU:OE2	2.52	0.42
1:B:216:GLN:OE1	1:B:216:GLN:CA	2.60	0.42
1:B:357:ASP:OD1	1:B:357:ASP:C	2.57	0.42
1:A:11:VAL:CG1	1:A:51:LEU:HG	2.50	0.42
1:A:179:ARG:HD3	1:A:294:VAL:HG13	2.02	0.42
1:A:309:ILE:H	1:A:309:ILE:HG12	1.65	0.42
1:B:120:ARG:HD2	1:B:348:TRP:CZ2	2.55	0.41
1:A:174:VAL:HG13	7:A:453:HOH:O	2.20	0.41
1:A:297:PRO:HD2	7:A:610:HOH:O	2.21	0.41
1:A:210:GLN:HE21	1:A:211:TRP:H	1.70	0.40
1:B:16:PHE:HB3	1:B:114:GLN:HE21	1.86	0.40
1:A:296:ALA:N	1:A:297:PRO:CD	2.84	0.40
1:A:114:GLN:NE2	1:A:240:ALA:HB1	2.29	0.40
1:A:308:GLY:CA	1:A:330:ARG:HH22	2.25	0.40

All (10) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:391:HOH:O	7:A:706:HOH:O[4_446]	1.17	1.03
7:A:392:HOH:O	7:A:609:HOH:O[4_446]	1.28	0.92
3:A:1001:GOL:O3	7:A:686:HOH:O[4_446]	1.39	0.81
7:A:389:HOH:O	7:A:559:HOH:O[4_456]	1.43	0.77
7:A:390:HOH:O	7:A:707:HOH:O[4_446]	1.66	0.54
3:A:1001:GOL:C3	7:A:686:HOH:O[4_446]	1.97	0.23
3:A:1001:GOL:O1	3:A:1002:GOL:C1[4_446]	2.02	0.18
7:A:665:HOH:O	7:B:664:HOH:O[4_456]	2.06	0.14

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:396:HOH:O	7:A:605:HOH:O[4_446]	2.12	0.08
7:A:393:HOH:O	7:A:710:HOH:O[4_446]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/387 (99%)	372 (97%)	12 (3%)	0	100	100
1	B	385/387 (100%)	373 (97%)	11 (3%)	1 (0%)	41	41
All	All	769/774 (99%)	745 (97%)	23 (3%)	1 (0%)	51	54

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	104	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	317/318 (100%)	305 (96%)	12 (4%)	33	34
1	B	318/318 (100%)	305 (96%)	13 (4%)	30	31
All	All	635/636 (100%)	610 (96%)	25 (4%)	32	33

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	154	PHE
1	A	175	ASP
1	A	201	SER
1	A	209	ASP
1	A	210	GLN
1	A	251	LYS
1	A	274	ARG
1	A	277	LEU
1	A	335	ARG
1	A	357	ASP
1	A	373	ARG
1	B	17	THR
1	B	22	GLU
1	B	47	ARG
1	B	103	ARG
1	B	154	PHE
1	B	204	TYR
1	B	210	GLN
1	B	216	GLN
1	B	285	LEU
1	B	342	THR
1	B	344	LYS
1	B	373	ARG
1	B	381	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN
1	A	141	ASN
1	A	153	HIS
1	A	172	ASN
1	A	210	GLN
1	A	257	GLN
1	A	387	GLN
1	B	114	GLN
1	B	153	HIS
1	B	172	ASN
1	B	210	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	1002	-	5,5,5	0.77	0	5,5,5	2.31	2 (40%)
3	GOL	B	1001	-	5,5,5	0.34	0	5,5,5	0.77	0
3	GOL	A	1001	-	5,5,5	0.40	0	5,5,5	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1002	-	-	1/4/4/4	-
3	GOL	B	1001	-	-	0/4/4/4	-
3	GOL	A	1001	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	1002	GOL	O2-C2-C3	4.25	127.85	109.12
3	A	1002	GOL	C3-C2-C1	-2.11	103.48	111.70

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1001	GOL	C1-C2-C3-O3
3	A	1002	GOL	O1-C1-C2-C3
3	A	1001	GOL	O2-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1002	GOL	0	1
3	A	1001	GOL	12	3

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/387 (99%)	-0.44	6 (1%) 72 75	3, 9, 25, 38	0
1	B	387/387 (100%)	-0.44	1 (0%) 94 94	3, 11, 23, 31	0
All	All	773/774 (99%)	-0.44	7 (0%) 84 86	3, 9, 24, 38	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	307	LEU	4.5
1	A	3	LEU	4.0
1	A	174	VAL	3.9
1	B	217	PRO	3.3
1	A	306	THR	2.6
1	A	175	ASP	2.1
1	A	141	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NA	B	904	1/1	0.88	0.23	44,44,44,44	0
3	GOL	A	1002	6/6	0.92	0.11	16,20,23,23	0
3	GOL	A	1001	6/6	0.93	0.29	454,454,454,454	0
3	GOL	B	1001	6/6	0.95	0.10	18,21,21,25	0
5	NA	A	904	1/1	0.98	0.26	26,26,26,26	0
6	CA	B	902	1/1	0.98	0.05	21,21,21,21	0
6	CA	A	902	1/1	0.99	0.03	18,18,18,18	0
4	CL	A	903	1/1	0.99	0.04	15,15,15,15	0
2	ZN	B	901	1/1	1.00	0.03	13,13,13,13	0
4	CL	B	903	1/1	1.00	0.03	20,20,20,20	0
2	ZN	A	901	1/1	1.00	0.05	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.