



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 11:34 am BST

PDB ID : 6UNC  
Title : The crystal structure of Phosphopantetheinyl Hydrolase (PptH) from Mycobacterium tuberculosis  
Authors : Mosior, J.W.; Sacchettini, J.C.; TB Structural Genomics Consortium (TB-SGC)  
Deposited on : 2019-10-11  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	 2% 84% 11%
1	B	313	 2% 84% 5% 10%
1	C	313	 4% 84% 10%
1	D	313	 7% 79% 8% 12%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 9287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metallophos domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	278	2252	1448	394	400	5	5	0	0	0
1	B	282	2280	1462	401	407	5	5	0	0	0
1	C	281	2264	1454	394	406	5	5	0	0	0
1	D	277	2234	1434	395	395	5	5	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP I6YEE1
A	2	ASN	-	expression tag	UNP I6YEE1
A	3	ALA	-	expression tag	UNP I6YEE1
A	4	VAL	-	expression tag	UNP I6YEE1
B	1	SER	-	expression tag	UNP I6YEE1
B	2	ASN	-	expression tag	UNP I6YEE1
B	3	ALA	-	expression tag	UNP I6YEE1
B	4	VAL	-	expression tag	UNP I6YEE1
C	1	SER	-	expression tag	UNP I6YEE1
C	2	ASN	-	expression tag	UNP I6YEE1
C	3	ALA	-	expression tag	UNP I6YEE1
C	4	VAL	-	expression tag	UNP I6YEE1
D	1	SER	-	expression tag	UNP I6YEE1
D	2	ASN	-	expression tag	UNP I6YEE1
D	3	ALA	-	expression tag	UNP I6YEE1
D	4	VAL	-	expression tag	UNP I6YEE1

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mn 1 1	0	0
2	A	1	Total Mn 1 1	0	0
2	D	1	Total Mn 1 1	0	0
2	C	1	Total Mn 1 1	0	0


- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Fe 1 1	0	0
3	A	1	Total Fe 1 1	0	0
3	D	1	Total Fe 1 1	0	0
3	C	1	Total Fe 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	75	Total O 75 75	0	0
4	B	96	Total O 96 96	0	0
4	C	45	Total O 45 45	0	0
4	D	33	Total O 33 33	0	0





MSE  
ARG  
THR  
GLN

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.19Å 92.53Å 183.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.86 – 2.50 47.21 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (44.86-2.50) 93.5 (47.21-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, $R_{free}$	0.186 , 0.222 0.189 , 0.224	Depositor DCC
$R_{free}$ test set	2287 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.2	Xtrriage
Anisotropy	0.347	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9287	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2326	0.43	0/3182
1	B	0.24	0/2355	0.43	0/3221
1	C	0.24	0/2339	0.42	0/3202
1	D	0.24	0/2306	0.44	0/3152
All	All	0.24	0/9326	0.43	0/12757

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2252	0	2139	6	0
1	B	2280	0	2163	10	0
1	C	2264	0	2138	8	0
1	D	2234	0	2131	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	75	0	0	0	0
4	B	96	0	0	1	0
4	C	45	0	0	0	2
4	D	33	0	0	0	2
All	All	9287	0	8571	35	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:151:LYS:O	1:B:155:VAL:HG23	1.97	0.65
1:D:58:ARG:NH1	1:D:90:ARG:O	2.33	0.60
1:B:135:MSE:HE3	1:B:207:ASN:HB2	1.84	0.58
1:A:135:MSE:HE3	1:A:207:ASN:HB2	1.85	0.57
1:B:34:LYS:NZ	1:B:61:GLU:OE2	2.41	0.53
1:D:161:ARG:NH1	1:D:223:GLU:OE1	2.39	0.52
1:C:180:ASP:OD1	1:C:180:ASP:N	2.36	0.50
1:A:52:ALA:HA	1:A:79:VAL:HG22	1.94	0.49
1:B:180:ASP:N	1:B:180:ASP:OD1	2.37	0.49
1:D:135:MSE:HE3	1:D:207:ASN:HB2	1.95	0.48
1:B:133:VAL:HG11	1:B:194:LEU:HD21	1.95	0.47
1:C:110:GLU:HG2	1:C:111:MSE:HG3	1.96	0.47
1:D:60:ASP:OD1	1:D:60:ASP:N	2.47	0.47
1:D:137:LEU:HD13	1:D:233:THR:HG21	1.96	0.47
1:B:60:ASP:OD1	1:B:60:ASP:N	2.46	0.46
1:A:126:ARG:H	1:A:126:ARG:HG3	1.52	0.46
1:D:20:LEU:HB3	1:D:287:LEU:HB2	1.98	0.46
1:B:202:PRO:HB3	1:B:243:VAL:HG13	1.98	0.45
1:B:213:ARG:NH1	4:B:504:HOH:O	2.42	0.45
1:D:180:ASP:OD1	1:D:180:ASP:N	2.36	0.44
1:D:68:LEU:HD11	1:D:72:ARG:NH2	2.33	0.44
1:A:17:GLU:CB	1:A:126:ARG:HD2	2.47	0.44
1:B:126:ARG:HG3	1:B:126:ARG:H	1.55	0.43
1:C:271:PRO:HA	1:C:274:TRP:NE1	2.32	0.43
1:C:60:ASP:N	1:C:60:ASP:OD1	2.48	0.43
1:A:123:TRP:NE1	1:A:126:ARG:O	2.41	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:ILE:O	1:A:134:PRO:HD3	2.20	0.42
1:B:79:VAL:HG22	1:B:116:PRO:HG3	2.01	0.42
1:C:68:LEU:HD11	1:C:72:ARG:NH2	2.34	0.42
1:D:123:TRP:HE1	1:D:127:GLY:H	1.68	0.42
1:D:67:ASP:HB2	1:D:111:MSE:HE1	2.01	0.41
1:C:67:ASP:OD1	1:C:70:ARG:NH2	2.52	0.41
1:C:199:TRP:CD2	1:D:31:LEU:HD12	2.55	0.41
1:C:212:LEU:HD12	1:C:215:PRO:HD3	2.02	0.41
1:D:131:THR:OG1	1:D:201:GLN:NE2	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:530:HOH:O	4:D:517:HOH:O[4_545]	2.07	0.13
4:C:531:HOH:O	4:D:523:HOH:O[4_545]	2.10	0.10

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	276/313 (88%)	263 (95%)	12 (4%)	1 (0%)	34 54
1	B	280/313 (90%)	268 (96%)	12 (4%)	0	100 100
1	C	279/313 (89%)	266 (95%)	12 (4%)	1 (0%)	34 54
1	D	275/313 (88%)	259 (94%)	14 (5%)	2 (1%)	22 39
All	All	1110/1252 (89%)	1056 (95%)	50 (4%)	4 (0%)	34 54

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	ALA
1	D	278	LYS
1	D	75	LYS
1	C	280	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/265 (89%)	232 (98%)	4 (2%)	60	82
1	B	239/265 (90%)	233 (98%)	6 (2%)	47	73
1	C	237/265 (89%)	231 (98%)	6 (2%)	47	73
1	D	234/265 (88%)	225 (96%)	9 (4%)	33	58
All	All	946/1060 (89%)	921 (97%)	25 (3%)	46	72

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LYS
1	A	180	ASP
1	A	212	LEU
1	A	265	GLU
1	B	28	THR
1	B	79	VAL
1	B	155	VAL
1	B	180	ASP
1	B	212	LEU
1	B	265	GLU
1	C	28	THR
1	C	70	ARG
1	C	79	VAL
1	C	180	ASP
1	C	212	LEU
1	C	265	GLU
1	D	34	LYS
1	D	70	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	79	VAL
1	D	88	THR
1	D	90	ARG
1	D	180	ASP
1	D	212	LEU
1	D	265	GLU
1	D	278	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	201	GLN
1	D	201	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	273/313 (87%)	0.06	7 (2%) 56 59	34, 48, 72, 106	0
1	B	277/313 (88%)	0.21	5 (1%) 68 71	33, 45, 77, 101	0
1	C	276/313 (88%)	0.25	13 (4%) 31 33	38, 59, 88, 109	0
1	D	272/313 (86%)	0.49	21 (7%) 13 13	48, 71, 96, 105	0
All	All	1098/1252 (87%)	0.25	46 (4%) 36 39	33, 55, 90, 109	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	16	ALA	6.0
1	D	280	TYR	5.5
1	C	199	TRP	4.4
1	C	292	TYR	4.3
1	B	292	TYR	4.0
1	D	197	LEU	3.8
1	B	293	ALA	3.7
1	D	148	ALA	3.6
1	D	192	ALA	3.3
1	C	16	ALA	3.3
1	C	294	PRO	3.2
1	D	15	GLY	3.0
1	A	280	TYR	3.0
1	B	14	VAL	2.9
1	C	198	ASP	2.9
1	A	292	TYR	2.9
1	D	258	TYR	2.9
1	C	293	ALA	2.9
1	D	190	THR	2.9
1	C	280	TYR	2.9
1	C	112	GLY	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	260	GLY	2.7
1	D	243	VAL	2.6
1	C	110	GLU	2.6
1	C	36	VAL	2.5
1	A	31	LEU	2.5
1	C	15	GLY	2.5
1	B	31	LEU	2.4
1	D	196	GLN	2.4
1	D	242	ALA	2.4
1	D	205	LEU	2.4
1	A	128	GLY	2.3
1	A	36	VAL	2.3
1	D	188	ALA	2.3
1	D	278	LYS	2.3
1	C	32	GLY	2.3
1	A	16	ALA	2.2
1	D	235	ASP	2.2
1	A	278	LYS	2.2
1	D	233	THR	2.2
1	D	129	PRO	2.1
1	D	189	ALA	2.1
1	D	240	TYR	2.1
1	C	67	ASP	2.0
1	D	68	LEU	2.0
1	D	241	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FE	D	402	1/1	0.70	0.10	43,43,43,43	1
3	FE	B	402	1/1	0.90	0.08	35,35,35,35	1
3	FE	A	402	1/1	0.92	0.06	28,28,28,28	1
3	FE	C	402	1/1	0.93	0.06	38,38,38,38	1
2	MN	C	401	1/1	0.96	0.08	52,52,52,52	0
2	MN	D	401	1/1	0.98	0.07	48,48,48,48	0
2	MN	B	401	1/1	0.98	0.09	40,40,40,40	0
2	MN	A	401	1/1	0.99	0.09	38,38,38,38	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





**Electron density around FE D 402:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




















## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.