



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2024 – 09:42 AM EST

PDB ID : 3UO8  
Title : Crystal structure of the MALT1 paracaspase (P1 form)  
Authors : Jeffrey, P.D.; Yu, J.W.; Shi, Y.  
Deposited on : 2011-11-16  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	390	
1	C	390	
2	L	6	
2	M	6	

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 11804 atoms, of which 5800 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mucosa-associated lymphoid tissue lymphoma translocation protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	B	364	5728	1835	2869	463	540	21	0	0	0
1	C	363	5698	1831	2847	464	535	21	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	338	MET	-	expression tag	UNP Q9UDY8
B	720	LEU	-	expression tag	UNP Q9UDY8
B	721	GLU	-	expression tag	UNP Q9UDY8
B	722	HIS	-	expression tag	UNP Q9UDY8
B	723	HIS	-	expression tag	UNP Q9UDY8
B	724	HIS	-	expression tag	UNP Q9UDY8
B	725	HIS	-	expression tag	UNP Q9UDY8
B	726	HIS	-	expression tag	UNP Q9UDY8
B	727	HIS	-	expression tag	UNP Q9UDY8
C	338	MET	-	expression tag	UNP Q9UDY8
C	720	LEU	-	expression tag	UNP Q9UDY8
C	721	GLU	-	expression tag	UNP Q9UDY8
C	722	HIS	-	expression tag	UNP Q9UDY8
C	723	HIS	-	expression tag	UNP Q9UDY8
C	724	HIS	-	expression tag	UNP Q9UDY8
C	725	HIS	-	expression tag	UNP Q9UDY8
C	726	HIS	-	expression tag	UNP Q9UDY8
C	727	HIS	-	expression tag	UNP Q9UDY8

- Molecule 2 is a protein called Z-Val-Arg-Pro-DL-Arg-fluoromethylketone.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	L	4	Total	C	H	N	O	0	0	0
			78	22	42	10	4			
2	M	4	Total	C	H	N	O	0	0	0
			78	22	42	10	4			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	105	Total	O	0	0
			105	105		
3	C	112	Total	O	0	0
			112	112		
3	L	2	Total	O	0	0
			2	2		
3	M	3	Total	O	0	0
			3	3		





- Molecule 2: Z-Val-Arg-Pro-DL-Arg-fluoromethylketone

Chain M: 33% 17% 17% 33%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.23Å 61.16Å 71.13Å 108.59° 97.23° 117.23°	Depositor
Resolution (Å)	34.26 – 1.90 34.26 – 1.90	Depositor EDS
% Data completeness (in resolution range)	92.3 (34.26-1.90) 92.4 (34.26-1.90)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.24 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, $R_{free}$	0.207 , 0.250 0.201 , 0.246	Depositor DCC
$R_{free}$ test set	3197 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.3	Xtrriage
Anisotropy	0.337	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.023 for k,h,-h-k-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11804	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.32	0/2910	0.49	0/3936
1	C	0.33	0/2901	0.49	0/3920
2	L	0.43	0/36	0.66	0/47
2	M	0.44	0/36	0.72	0/47
All	All	0.33	0/5883	0.49	0/7950

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2859	2869	2876	55	0
1	C	2851	2847	2854	46	0
2	L	36	42	44	4	0
2	M	36	42	44	5	0
3	B	105	0	0	8	0
3	C	112	0	0	4	0
3	L	2	0	0	1	0
3	M	3	0	0	2	0
All	All	6004	5800	5818	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.



The worst 5 of 101 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:662:ASP:HB2	3:B:903:HOH:O	1.87	0.73
1:C:594:PHE:CE1	1:C:600:ILE:HD12	2.24	0.73
2:L:1:VAL:HA	3:L:101:HOH:O	1.92	0.69
1:B:502:GLN:H	2:L:1:VAL:HG23	1.57	0.68
1:B:499:PHE:HB2	1:B:510:ILE:HD13	1.77	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	360/390 (92%)	338 (94%)	22 (6%)	0	100	100
1	C	355/390 (91%)	342 (96%)	13 (4%)	0	100	100
2	L	2/6 (33%)	2 (100%)	0	0	100	100
2	M	2/6 (33%)	2 (100%)	0	0	100	100
All	All	719/792 (91%)	684 (95%)	35 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	316/347 (91%)	312 (99%)	4 (1%)	69	68
1	C	312/347 (90%)	312 (100%)	0	100	100
2	L	4/4 (100%)	4 (100%)	0	100	100
2	M	4/4 (100%)	3 (75%)	1 (25%)	0	0
All	All	636/702 (91%)	631 (99%)	5 (1%)	81	82

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	401	LEU
1	B	584	HIS
1	B	637	LEU
1	B	652	GLU
2	M	1	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	502	GLN
1	C	508	ASN
1	B	666	HIS
1	C	371	ASN
1	C	422	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	364/390 (93%)	0.63	50 (13%) 3 3	20, 41, 80, 158	0
1	C	363/390 (93%)	0.68	49 (13%) 3 3	19, 38, 76, 111	0
2	L	4/6 (66%)	0.54	1 (25%) 0 0	29, 36, 50, 67	0
2	M	4/6 (66%)	0.41	0 100 100	31, 36, 47, 57	0
All	All	735/792 (92%)	0.65	100 (13%) 3 3	19, 39, 79, 158	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	474	ILE	7.3
1	C	659	VAL	6.6
1	C	476	ILE	5.9
1	C	641	ILE	5.5
1	B	476	ILE	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.