



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 12:06 pm BST

PDB ID : 1UQ5  
Title : RICIN A-CHAIN (RECOMBINANT) N122A MUTANT  
Authors : Marsden, C.J.; Fulop, V.  
Deposited on : 2003-10-15  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2596 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RICIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	2076	1315	366	390	5	0	0	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	118	ALA	ASN	engineered mutation	UNP P02879

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	506	Total	O	0	0
			506	506		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RICIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.50Å 67.50Å 140.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.00 – 1.40 45.20 – 1.40	Depositor EDS
% Data completeness (in resolution range)	93.9 (46.00-1.40) 93.9 (45.20-1.40)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.20 (at 1.40Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.177 , 0.214 0.195 , 0.229	Depositor DCC
$R_{free}$ test set	2448 reflections (4.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.3	Xtrriage
Anisotropy	0.361	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 53.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2596	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.75	31/2122 (1.5%)	1.75	42/2887 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	ARG	CG-CD	-11.09	1.24	1.51
1	A	145	GLU	CD-OE2	10.90	1.37	1.25
1	A	228	SER	CB-OG	-10.54	1.28	1.42
1	A	39	ARG	CB-CG	10.19	1.80	1.52
1	A	243	TYR	CD1-CE1	8.43	1.51	1.39
1	A	145	GLU	CD-OE1	8.17	1.34	1.25
1	A	33	THR	CB-CG2	-7.93	1.26	1.52
1	A	185	GLU	CD-OE2	7.59	1.33	1.25
1	A	220	GLU	CD-OE1	-7.34	1.17	1.25
1	A	187	GLU	CD-OE2	7.00	1.33	1.25
1	A	215	SER	CB-OG	6.71	1.50	1.42
1	A	246	SER	CB-OG	-6.63	1.33	1.42
1	A	257	TYR	CZ-OH	6.54	1.49	1.37
1	A	135	GLU	CD-OE1	6.40	1.32	1.25
1	A	194	TYR	CB-CG	-6.30	1.42	1.51
1	A	166	ARG	CG-CD	6.17	1.67	1.51
1	A	154	TYR	CZ-OH	6.16	1.48	1.37
1	A	177	GLU	CD-OE1	-5.81	1.19	1.25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	10	ASN	CG-OD1	-5.70	1.11	1.24
1	A	162	PRO	N-CD	5.69	1.55	1.47
1	A	194	TYR	CE1-CZ	5.67	1.46	1.38
1	A	33	THR	CA-CB	5.65	1.68	1.53
1	A	135	GLU	CG-CD	5.64	1.60	1.51
1	A	115	TYR	CG-CD1	5.61	1.46	1.39
1	A	257	TYR	CD2-CE2	5.52	1.47	1.39
1	A	85	ARG	NE-CZ	5.40	1.40	1.33
1	A	196	ARG	CD-NE	5.33	1.55	1.46
1	A	196	ARG	CG-CD	5.32	1.65	1.51
1	A	246	SER	CA-CB	5.30	1.60	1.52
1	A	194	TYR	C-O	5.11	1.33	1.23
1	A	185	GLU	CG-CD	-5.00	1.44	1.51

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	254	LEU	CB-CG-CD1	-26.14	66.55	111.00
1	A	56	ARG	NE-CZ-NH2	-13.07	113.77	120.30
1	A	39	ARG	NE-CZ-NH2	-12.88	113.86	120.30
1	A	26	ARG	NE-CZ-NH2	-12.18	114.21	120.30
1	A	39	ARG	CA-CB-CG	11.12	137.86	113.40
1	A	254	LEU	CB-CG-CD2	10.68	129.15	111.00
1	A	201	ASP	CB-CG-OD2	10.44	127.70	118.30
1	A	189	ARG	NE-CZ-NH1	10.34	125.47	120.30
1	A	26	ARG	NE-CZ-NH1	10.34	125.47	120.30
1	A	134	ARG	NE-CZ-NH1	10.23	125.41	120.30
1	A	48	ARG	NE-CZ-NH1	8.97	124.78	120.30
1	A	189	ARG	CD-NE-CZ	8.89	136.05	123.60
1	A	228	SER	CB-CA-C	8.47	126.19	110.10
1	A	67	GLU	OE1-CD-OE2	-8.35	113.28	123.30
1	A	75	ASP	CB-CG-OD1	8.35	125.81	118.30
1	A	48	ARG	NE-CZ-NH2	-8.15	116.23	120.30
1	A	123	TYR	CD1-CE1-CZ	-8.05	112.56	119.80
1	A	31	ARG	NE-CZ-NH2	-7.79	116.40	120.30
1	A	145	GLU	OE1-CD-OE2	7.75	132.60	123.30
1	A	26	ARG	CD-NE-CZ	7.67	134.33	123.60
1	A	193	ARG	NE-CZ-NH2	-7.62	116.49	120.30
1	A	189	ARG	NE-CZ-NH2	-7.48	116.56	120.30
1	A	191	ARG	NE-CZ-NH1	7.40	124.00	120.30
1	A	26	ARG	CG-CD-NE	-7.07	96.95	111.80
1	A	197	ARG	NE-CZ-NH2	-6.81	116.89	120.30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	TYR	CG-CD1-CE1	-6.65	115.98	121.30
1	A	257	TYR	CZ-CE2-CD2	-6.33	114.10	119.80
1	A	196	ARG	CG-CD-NE	6.23	124.89	111.80
1	A	257	TYR	CE1-CZ-CE2	6.19	129.71	119.80
1	A	254	LEU	CB-CA-C	5.85	121.32	110.20
1	A	146	GLU	CG-CD-OE1	5.83	129.95	118.30
1	A	154	TYR	CG-CD2-CE2	-5.62	116.80	121.30
1	A	124	ASP	CB-CG-OD2	5.62	123.36	118.30
1	A	191	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	A	56	ARG	NH1-CZ-NH2	5.54	125.49	119.40
1	A	257	TYR	CB-CG-CD2	-5.49	117.71	121.00
1	A	145	GLU	CG-CD-OE2	-5.47	107.36	118.30
1	A	91	TYR	CZ-CE2-CD2	-5.19	115.13	119.80
1	A	194	TYR	CD1-CE1-CZ	5.17	124.46	119.80
1	A	201	ASP	OD1-CG-OD2	-5.11	113.59	123.30
1	A	29	ARG	NE-CZ-NH2	5.03	122.81	120.30
1	A	194	TYR	CB-CG-CD2	5.01	124.01	121.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	ALA	Mainchain
1	A	220	GLU	Sidechain
1	A	33	THR	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2076	0	2040	18	0
2	A	4	0	3	1	0
3	A	10	0	0	0	0
4	A	506	0	0	7	0
All	All	2596	0	2043	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:ARG:CB	1:A:39:ARG:CG	1.80	1.59
1:A:39:ARG:CB	1:A:39:ARG:CD	2.56	0.79
1:A:239:LYS:HD2	4:A:2454:HOH:O	1.86	0.76
1:A:239:LYS:HB3	4:A:2453:HOH:O	1.96	0.66
1:A:99:GLU:HG3	4:A:2270:HOH:O	1.97	0.64
1:A:64:ASN:HB2	1:A:145:GLU:CD	2.21	0.61
1:A:40:HIS:O	1:A:41:GLU:HB2	2.02	0.60
1:A:39:ARG:HB2	1:A:39:ARG:CG	2.22	0.51
1:A:189:ARG:NE	4:A:2399:HOH:O	2.46	0.49
1:A:48:ARG:HD2	4:A:2160:HOH:O	2.13	0.48
1:A:191:ARG:HG3	1:A:196:ARG:HH21	1.83	0.43
1:A:11:PHE:HB2	1:A:24:PHE:CD1	2.54	0.42
1:A:39:ARG:HB3	1:A:39:ARG:CG	2.22	0.42
1:A:227:ALA:HB2	4:A:2436:HOH:O	2.20	0.41
1:A:141:ASN:OD1	2:A:1268:ACT:OXT	2.39	0.41
1:A:194:TYR:HB2	1:A:196:ARG:HG3	2.03	0.41
1:A:211:TRP:CH2	1:A:256:VAL:HB	2.56	0.41
1:A:239:LYS:CD	4:A:2454:HOH:O	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	261/263 (99%)	258 (99%)	3 (1%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	221/221 (100%)	219 (99%)	2 (1%)	78 58

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	239	LYS
1	A	254	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	23	ASN
1	A	98	GLN
1	A	141	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	ACT	A	1268	-	1,3,3	2.22	1 (100%)	0,3,3	0.00	-
3	SO4	A	1270	-	4,4,4	1.14	0	6,6,6	1.54	2 (33%)
3	SO4	A	1269	-	4,4,4	0.20	0	6,6,6	0.82	0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1268	ACT	CH3-C	2.22	1.51	1.48

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1270	SO4	O4-S-O1	-2.45	96.53	109.31
3	A	1270	SO4	O4-S-O3	2.12	118.12	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1268	ACT	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/263 (100%)	-0.19	4 (1%) 73 72	8, 14, 26, 36	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	GLN	2.4
1	A	110	ASP	2.2
1	A	267	PHE	2.1
1	A	88	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	1269	5/5	0.96	0.10	40,49,51,51	0
2	ACT	A	1268	4/4	0.97	0.07	14,17,18,19	0
3	SO4	A	1270	5/5	0.99	0.05	16,22,26,29	0

## 6.5 Other polymers

There are no such residues in this entry.