

Full wwPDB X-ray Structure Validation Report (i)

Sep 5, 2023 – 01:20 PM EDT

PDB ID : 3UV3

Title: Ec IspH in complex with but-2-ynyl diphosphate (1086)

Authors: Span, I.; Wang, K.; Wang, W.; Zhang, Y.; Bacher, A.; Eisenreich, W.; Schulz,

C.; Oldfield, E.; Groll, M.

Deposited on : 2011-11-29

Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.35

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	324	81%	10%	• • 5%
1	В	324	82%	11%	



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5214 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 4-hydroxy-3-methylbut-2-enyl diphosphate reductase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	309	Total 2383	C 1487	N 427	O 459	S 10	0	0	0
1	В	310	Total 2390	C 1492	N 428	O 460	S 10	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	expression tag	UNP P62623
A	-6	HIS	-	expression tag	UNP P62623
A	-5	HIS	-	expression tag	UNP P62623
A	-4	HIS	-	expression tag	UNP P62623
A	-3	HIS	-	expression tag	UNP P62623
A	-2	HIS	-	expression tag	UNP P62623
A	-1	GLY	-	expression tag	UNP P62623
A	0	SER	-	expression tag	UNP P62623
В	-7	HIS	-	expression tag	UNP P62623
В	-6	HIS	-	expression tag	UNP P62623
В	-5	HIS	-	expression tag	UNP P62623
В	-4	HIS	-	expression tag	UNP P62623
В	-3	HIS	-	expression tag	UNP P62623
В	-2	HIS	-	expression tag	UNP P62623
В	-1	GLY	-	expression tag	UNP P62623
В	0	SER	-	expression tag	UNP P62623


• Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 8 4 4	0	0
2	В	1	Total Fe S 8 4 4	0	0

 \bullet Molecule 3 is but-2-yn-1-yl trihydrogen diphosphate (three-letter code: 0CM) (formula: $C_4H_8O_7P_2).$



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 13	C 4	O 7	P 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total C O P 13 4 7 2	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	197	Total O 197 197	0	0
4	В	202	Total O 202 202	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 4-hydroxy-3-methylbut-2-enyl diphosphate reductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	70.23Å 80.40Å 111.95Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 - 1.60	Depositor
Resolution (A)	47.82 - 1.60	EDS
% Data completeness	96.9 (40.00-1.60)	Depositor
(in resolution range)	96.9 (47.82-1.60)	EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.73 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.167 , 0.221	Depositor
it, it free	0.167 , 0.220	DCC
R_{free} test set	4079 reflections $(5.00%)$	wwPDB-VP
Wilson B-factor (\mathring{A}^2)	17.7	Xtriage
Anisotropy	0.634	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 48.6	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5214	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 47.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.2987e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 0CM, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
Mol Chain		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	1.08	6/2418 (0.2%)	1.41	$11/3273 \ (0.3\%)$	
1	В	1.07	5/2425 (0.2%)	1.29	$19/3283 \ (0.6\%)$	
All	All	1.07	11/4843 (0.2%)	1.35	$30/6556 \ (0.5\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
1	A	230	ARG	CZ-NH2	7.69	1.43	1.33
1	В	184	ARG	CZ-NH2	7.66	1.43	1.33
1	A	151	TRP	CD2-CE2	7.66	1.50	1.41
1	В	151	TRP	CD2-CE2	6.56	1.49	1.41
1	A	151	TRP	CG-CD1	6.50	1.45	1.36
1	A	230	ARG	CD-NE	-6.43	1.35	1.46
1	В	151	TRP	CG-CD1	6.00	1.45	1.36
1	В	38	GLU	CD-OE1	-5.55	1.19	1.25
1	В	230	ARG	CD-NE	-5.38	1.37	1.46
1	A	38	GLU	CD-OE1	-5.35	1.19	1.25
1	A	255	TRP	CD2-CE2	5.33	1.47	1.41

All (30) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\operatorname{Ideal}(^{o})$
1	A	230	ARG	NE-CZ-NH1	-39.57	100.52	120.30
1	A	230	ARG	NE-CZ-NH2	33.30	136.95	120.30
1	В	230	ARG	NE-CZ-NH1	-24.49	108.05	120.30
1	В	230	ARG	NE-CZ-NH2	21.91	131.25	120.30
1	В	184	ARG	NE-CZ-NH1	-13.55	113.53	120.30
1	A	230	ARG	CD-NE-CZ	12.65	141.30	123.60
1	В	52	ARG	NE-CZ-NH2	11.10	125.85	120.30
1	В	184	ARG	NE-CZ-NH2	10.81	125.71	120.30
1	В	281	ARG	NE-CZ-NH1	10.41	125.50	120.30
1	В	237	ARG	NE-CZ-NH2	9.63	125.12	120.30
1	В	131	MET	CG-SD-CE	8.57	113.92	100.20
1	A	36	ARG	NE-CZ-NH2	-8.57	116.02	120.30
1	В	230	ARG	CG-CD-NE	-8.27	94.44	111.80
1	В	230	ARG	CD-NE-CZ	7.93	134.70	123.60
1	A	230	ARG	CG-CD-NE	-7.53	95.99	111.80
1	В	155	VAL	CG1-CB-CG2	7.44	122.80	110.90
1	В	237	ARG	NE-CZ-NH1	-6.98	116.81	120.30
1	В	81	ARG	NE-CZ-NH2	-6.82	116.89	120.30
1	В	52	ARG	NE-CZ-NH1	-6.79	116.91	120.30
1	A	141	MET	CG-SD-CE	6.74	110.98	100.20
1	В	293	LEU	CB-CG-CD1	-6.29	100.31	111.00
1	В	281	ARG	NE-CZ-NH2	-6.26	117.17	120.30
1	A	161	LEU	CB-CG-CD2	6.09	121.36	111.00
1	В	149	ASP	CB-CG-OD1	5.93	123.63	118.30
1	В	16	ASP	CB-CG-OD2	5.83	123.55	118.30
1	A	179	ASP	CB-CG-OD2	5.74	123.47	118.30
1	A	302	PHE	CB-CG-CD2	5.54	124.67	120.80
1	A	87	ARG	NE-CZ-NH2	-5.49	117.55	120.30
1	В	155	VAL	CA-CB-CG1	5.42	119.03	110.90
1	A	302	PHE	CB-CG-CD1	-5.24	117.13	120.80

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	230	ARG	Sidechain
1	A	7	ASN	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2383	0	2395	32	0
1	В	2390	0	2404	24	1
2	A	8	0	0	0	0
2	В	8	0	0	0	0
3	A	13	0	6	1	0
3	В	13	0	8	0	0
4	A	197	0	0	7	2
4	В	202	0	0	7	1
All	All	5214	0	4813	56	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:ARG:HH21	1:A:36:ARG:HG2	1.20	1.02
1:A:36:ARG:HG2	1:A:36:ARG:NH2	1.77	0.91
1:A:103:HIS:HD2	4:A:448:HOH:O	1.67	0.78
1:A:36:ARG:HH21	1:A:36:ARG:CG	1.92	0.78
1:B:42:ASN:HD22	1:B:45:VAL:H	1.30	0.78
1:B:51:GLU:OE2	1:B:52:ARG:HD2	1.83	0.78
1:B:236:GLN:HG2	4:B:404:HOH:O	1.86	0.75
1:A:42:ASN:HD22	1:A:45:VAL:H	1.32	0.74
1:A:7:ASN:OD1	4:A:564:HOH:O	2.06	0.72
1:B:83:GLU:OE2	4:B:461:HOH:O	2.10	0.68
1:B:103:HIS:HD2	4:B:416:HOH:O	1.79	0.66
1:B:297:GLU:HG3	4:B:465:HOH:O	1.95	0.65
1:A:257:LYS:O	1:A:258:GLU:CB	2.45	0.64
1:B:136:ASN:ND2	1:B:138:GLU:H	1.96	0.64
1:A:257:LYS:O	1:A:258:GLU:HB3	1.99	0.61
1:A:111:ARG:NH1	4:A:422:HOH:O	2.33	0.61
1:A:112:ARG:HB3	1:A:114:GLU:HG3	1.84	0.60
1:A:4:LEU:HD22	1:A:261:CYS:SG	2.43	0.59
1:B:112:ARG:HG3	1:B:112:ARG:HH21	1.68	0.59
1:B:68:ILE:HD13	1:B:90:THR:HB	1.83	0.59
1:A:136:ASN:ND2	1:A:138:GLU:H	2.01	0.58
1:A:161:LEU:HD13	1:A:188:ILE:HD12	1.87	0.56
1:A:136:ASN:HD22	1:A:136:ASN:C	2.09	0.56



 $Continued\ from\ previous\ page...$

A + a m 1		Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:B:7:ASN:OD1	4:B:460:HOH:O	2.18	0.56
1:A:57:ILE:HD12	1:A:63:VAL:HG22	1.90	0.54
1:B:103:HIS:HE1	4:B:327:HOH:O	1.91	0.54
1:A:60:ILE:HD12	1:A:69:LEU:HD11	1.91	0.53
1:B:136:ASN:HD22	1:B:138:GLU:H	1.56	0.53
1:A:36:ARG:HD2	1:A:60:ILE:HD11	1.90	0.52
1:A:236:GLN:HG3	1:A:242:ALA:HB3	1.91	0.52
1:A:103:HIS:HE1	4:A:322:HOH:O	1.93	0.52
1:A:36:ARG:NH2	1:A:36:ARG:CG	2.53	0.52
1:B:148:ASP:HA	1:B:151:TRP:CD1	2.46	0.50
1:B:281:ARG:HD3	4:B:379:HOH:O	2.13	0.49
1:A:74:HIS:CD2	3:A:318:0CM:H6	2.47	0.49
1:A:1:MET:N	4:A:440:HOH:O	2.46	0.48
1:A:136:ASN:HD22	1:A:138:GLU:H	1.61	0.48
1:B:8:PRO:HG2	1:B:202:ASN:HB3	1.96	0.47
1:A:300:ILE:HD11	1:A:302:PHE:CE2	2.50	0.47
1:B:136:ASN:HD21	1:B:138:GLU:HB2	1.79	0.47
1:B:51:GLU:OE2	1:B:52:ARG:CD	2.61	0.47
1:B:161:LEU:HB3	1:B:188:ILE:HG13	1.97	0.45
1:A:60:ILE:CD1	1:A:69:LEU:HD11	2.47	0.45
1:A:15:VAL:HG13	1:A:40:VAL:HG11	1.99	0.45
1:A:42:ASN:ND2	1:A:45:VAL:H	2.09	0.44
4:A:424:HOH:O	1:B:189:VAL:HG21	2.17	0.44
1:A:42:ASN:HD21	1:A:44:TYR:HB3	1.83	0.43
1:A:136:ASN:HD22	1:A:137:PRO:N	2.17	0.42
1:A:159:GLU:O	1:A:160:LYS:HD3	2.19	0.42
1:A:5:LEU:HD12	1:A:264:VAL:HB	2.00	0.42
1:B:230:ARG:HD3	1:B:230:ARG:HA	1.69	0.42
1:B:299:ASN:HD22	1:B:299:ASN:C	2.22	0.42
1:A:59:GLN:HB3	4:A:450:HOH:O	2.21	0.41
1:B:112:ARG:HB3	1:B:114:GLU:HG3	2.02	0.41
1:B:310:VAL:HG12	1:B:310:VAL:O	2.21	0.41
1:B:42:ASN:ND2	1:B:45:VAL:H	2.07	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
4:A:345:HOH:O	4:B:358:HOH:O[4_555]	1.22	0.98
1:B:128:GLU:OE2	4:A:426:HOH:O[2_455]	1.88	0.32



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	307/324~(95%)	304 (99%)	3 (1%)	0	100	100
1	В	308/324~(95%)	305 (99%)	3 (1%)	0	100	100
All	All	$615/648 \; (95\%)$	609 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	es
1	A	$257/271 \ (95\%)$	245 (95%)	12 (5%)	26 7	
1	В	258/271 (95%)	242 (94%)	16 (6%)	18 4	
All	All	515/542~(95%)	487 (95%)	28 (5%)	22 5	

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	5	LEU
1	A	57	ILE
1	A	69	LEU
1	A	103	HIS
1	A	136	ASN
1	A	153	LEU
1	A	161	LEU



Continued from previous page...

Mol	Chain	Res	Type
1	A	187	LYS
1	A	230	ARG
1	A	258	GLU
1	A	294	GLU
1	В	5	LEU
1	В	136	ASN
1	В	138	GLU
1	В	148	ASP
1	В	153	LEU
1	В	155	VAL
1	В	159	GLU
1	В	161	LEU
1	В	230	ARG
1	В	249	LYS
1	В	253	GLU
1	В	273	ILE
1	В	285	LEU
1	В	299	ASN
1	В	300	ILE
1	В	306	LYS

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such side chains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	59	GLN
1	A	78	GLN
1	A	103	HIS
1	A	121	HIS
1	A	133	GLN
1	A	136	ASN
1	A	229	ASN
1	A	236	GLN
1	A	252	GLN
1	В	42	ASN
1	В	78	GLN
1	В	103	HIS
1	В	133	GLN
1	В	136	ASN
1	В	213	GLN
1	В	229	ASN
1	В	283	GLN



Continued from previous page...

Mol	Chain	Res	Type
1	В	299	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Dag	Dag	Dec	Dag	Res Link	Bond lengths			Bond angles		
MIOI	туре			LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2				
3	0CM	A	318	-	10,12,12	1.89	3 (30%)	12,17,17	0.88	0				
2	SF4	A	317	1,4	0,12,12	-	-	-						
3	0CM	В	318	-	10,12,12	1.42	2 (20%)	12,17,17	0.98	0				
2	SF4	В	317	1,4	0,12,12	-	-	-						

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0CM	A	318	-	-	5/11/12/12	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0CM	В	318	-	-	5/11/12/12	-
2	SF4	A	317	1,4	-	-	0/6/5/5
2	SF4	В	317	1,4	-	-	0/6/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
3	A	318	0CM	O5-C4	4.10	1.50	1.43
3	A	318	0CM	P6-O11	-2.40	1.44	1.55
3	В	318	0CM	O5-C4	2.26	1.47	1.43
3	В	318	0CM	P6-O10	-2.18	1.43	1.50
3	A	318	0CM	P8-O12	-2.13	1.46	1.54

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	318	0CM	C4-O5-P6-O7
3	A	318	0CM	C3-C4-O5-P6
3	В	318	0CM	C4-O5-P6-O7
3	В	318	0CM	C4-O5-P6-O10
3	В	318	0CM	C3-C4-O5-P6
3	В	318	0CM	C2-C3-C4-O5
3	A	318	0CM	C2-C3-C4-O5
3	A	318	0CM	C4-O5-P6-O10
3	A	318	0CM	C4-O5-P6-O11
3	В	318	0CM	C4-O5-P6-O11

There are no ring outliers.

1 monomer is involved in 1 short contact:

\mathbf{Mol}	Chain	Res	Type	Clashes	Symm-Clashes
3	A	318	0CM	1	0

5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q<0.9
1	A	309/324~(95%)	-0.41	3 (0%)	82 82	13, 22, 40, 62	0
1	В	310/324 (95%)	-0.35	2 (0%)	89 89	13, 23, 46, 69	0
All	All	619/648 (95%)	-0.38	5 (0%)	86 86	13, 23, 45, 69	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	310	VAL	3.5
1	A	88	ASP	3.3
1	A	309	ARG	2.6
1	A	86	SER	2.4
1	В	258	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	SF4	В	317	8/8	0.98	0.08	14,14,15,22	0
2	SF4	A	317	8/8	0.99	0.07	13,14,14,21	0
3	0CM	A	318	13/13	0.99	0.05	8,9,11,11	0
3	0CM	В	318	13/13	0.99	0.04	8,9,10,12	0

6.5 Other polymers (i)

There are no such residues in this entry.

