



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2018 – 12:13 pm GMT

PDB ID : 5UV8
Title : Interleukin-3 Receptor Complex
Authors : Broughton, S.E.; Parker, M.W.
Deposited on : 2017-02-19
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk31020
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk31020


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2449 (2.70-2.70)
Clashscore	122126	2756 (2.70-2.70)
Ramachandran outliers	120053	2716 (2.70-2.70)
Sidechain outliers	120020	2716 (2.70-2.70)
RSRZ outliers	108989	2376 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	 9% 87% 10% 2%
1	G	288	 2% 86% 8% 6%
2	B	122	 2% 83% 14% 1%
2	I	122	 2% 79% 11% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	BMA	A	403	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-3 receptor subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	Total	C	N	O	S	0	0	0
			2105	1331	378	381	15			
1	G	265	Total	C	N	O	S	0	0	0
			2143	1357	379	391	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	212	GLN	ASN	engineered mutation	UNP P26951
A	299	VAL	ALA	engineered mutation	UNP P26951
G	212	GLN	ASN	engineered mutation	UNP P26951
G	299	VAL	ALA	engineered mutation	UNP P26951

- Molecule 2 is a protein called Interleukin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	105	Total	C	N	O	S	0	0	0
			845	532	152	157	4			
2	I	108	Total	C	N	O	S	0	0	0
			871	551	155	161	4			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	13	TYR	TRP	engineered mutation	UNP P08700
I	13	TYR	TRP	engineered mutation	UNP P08700

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	G	1	Total 14	8	1	5	0	0
3	G	1	Total 14	8	1	5	0	0
3	G	1	Total 14	8	1	5	0	0
3	G	1	Total 14	8	1	5	0	0

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		
4	G	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	G	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	G	1	Total	C	O	0	0
			6	3	3		
6	G	1	Total	C	O	0	0
			6	3	3		
6	G	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	33	Total	O	0	0
			33	33		
7	B	9	Total	O	0	0
			9	9		
7	G	45	Total	O	0	0
			45	45		
7	I	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-3 receptor subunit alpha




- Molecule 1: Interleukin-3 receptor subunit alpha



- Molecule 2: Interleukin-3



- Molecule 2: Interleukin-3



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.04Å 132.04Å 210.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	114.35 – 2.70 48.10 – 2.70	Depositor EDS
% Data completeness (in resolution range)	100.0 (114.35-2.70) 100.0 (48.10-2.70)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.11 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.242 , 0.286 0.245 , 0.292	Depositor DCC
R_{free} test set	1535 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6261	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BMA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2157	0.69	0/2919
1	G	0.43	0/2198	0.71	0/2979
2	B	0.42	0/860	0.64	0/1164
2	I	0.42	0/888	0.66	0/1204
All	All	0.43	0/6103	0.69	0/8266

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2105	0	2031	0	0
1	G	2143	0	2065	7	0
2	B	845	0	848	4	0
2	I	871	0	875	12	0
3	A	70	0	60	0	0
3	G	56	0	49	0	0
4	A	22	0	20	0	0
4	G	11	0	10	0	0
5	A	10	0	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	G	10	0	10	0	0
6	G	18	0	24	0	0
7	A	33	0	0	0	0
7	B	9	0	0	0	0
7	G	45	0	0	0	0
7	I	13	0	0	0	0
All	All	6261	0	6002	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:87:ASN:CA	1:G:88:PRO:N	2.43	0.82
2:B:29:GLN:HG3	2:I:29:GLN:HG3	1.73	0.70
2:I:30:PRO:HG3	2:I:104:TRP:HE1	1.60	0.67
2:I:30:PRO:O	2:I:108:ARG:NH1	2.31	0.64
2:I:29:GLN:O	2:I:31:PRO:HD3	1.99	0.63
2:B:29:GLN:HG3	2:I:29:GLN:CG	2.33	0.57
1:G:224:MET:HE3	1:G:270:VAL:HG21	1.87	0.55
2:I:30:PRO:HG3	2:I:104:TRP:NE1	2.22	0.54
2:I:16:CYS:CB	2:I:84:CYS:HG	2.17	0.54
2:I:29:GLN:N	2:I:30:PRO:HD2	2.24	0.51
2:I:30:PRO:HG2	2:I:108:ARG:NH1	2.26	0.51
1:G:275:ARG:HD3	1:G:280:GLU:OE1	2.13	0.49
1:G:224:MET:CE	1:G:270:VAL:HG21	2.44	0.47
1:G:224:MET:HE1	1:G:240:LEU:HD23	1.97	0.46
2:B:55:ARG:HB3	2:B:56:PRO:HD3	1.97	0.46
2:I:25:THR:O	2:I:29:GLN:HG2	2.18	0.44
2:B:29:GLN:HG3	2:I:29:GLN:CD	2.40	0.42
2:I:55:ARG:HB3	2:I:56:PRO:HD3	2.01	0.42
1:G:214:THR:HG22	1:G:215:ALA:N	2.35	0.41
1:G:149:TYR:CD2	1:G:173:LEU:HD11	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/288 (86%)	238 (96%)	10 (4%)	1 (0%)	36	64
1	G	257/288 (89%)	246 (96%)	11 (4%)	0	100	100
2	B	101/122 (83%)	101 (100%)	0	0	100	100
2	I	104/122 (85%)	102 (98%)	2 (2%)	0	100	100
All	All	711/820 (87%)	687 (97%)	23 (3%)	1 (0%)	53	80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	265	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/257 (90%)	223 (96%)	8 (4%)	39	69
1	G	237/257 (92%)	232 (98%)	5 (2%)	56	83
2	B	95/110 (86%)	94 (99%)	1 (1%)	76	91
2	I	98/110 (89%)	96 (98%)	2 (2%)	58	84
All	All	661/734 (90%)	645 (98%)	16 (2%)	52	81

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ARG
1	A	45	ARG
1	A	57	ASP
1	A	171	SER
1	A	187	ARG
1	A	200	VAL
1	A	217	CYS
1	A	293	CYS
2	B	15	ASN
1	G	165	CYS
1	G	200	VAL
1	G	217	CYS
1	G	281	PHE
1	G	293	CYS
2	I	34	LEU
2	I	94	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	289	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	401	1,3	14,14,15	0.41	0	17,19,21	1.18	1 (5%)
3	NAG	A	402	3,4	14,14,15	0.69	0	17,19,21	1.46	1 (5%)
4	BMA	A	403	3	11,11,12	0.56	0	15,15,17	1.33	3 (20%)
3	NAG	A	404	1	14,14,15	0.76	0	17,19,21	1.62	2 (11%)
5	FUC	A	405	3	9,10,11	0.56	0	13,14,16	0.97	1 (7%)
3	NAG	A	406	1,3,5	14,14,15	0.43	0	17,19,21	1.84	5 (29%)
3	NAG	A	407	3,4	14,14,15	0.52	0	17,19,21	1.42	2 (11%)
4	BMA	A	408	3	11,11,12	0.45	0	15,15,17	0.89	1 (6%)
6	GOL	G	401	-	5,5,5	0.33	0	5,5,5	0.85	0
6	GOL	G	402	-	5,5,5	0.28	0	5,5,5	0.26	0
6	GOL	G	403	-	5,5,5	0.36	0	5,5,5	0.36	0
5	FUC	G	404	3	9,10,11	0.50	0	13,14,16	2.25	3 (23%)
3	NAG	G	405	1,3,5	14,14,15	0.45	0	17,19,21	1.20	2 (11%)
3	NAG	G	406	3,4	14,14,15	0.42	0	17,19,21	1.17	2 (11%)
4	BMA	G	407	3	11,11,12	0.29	0	15,15,17	0.81	1 (6%)
3	NAG	G	408	1	14,14,15	0.75	1 (7%)	17,19,21	1.35	3 (17%)
3	NAG	G	409	1	14,14,15	0.40	0	17,19,21	1.18	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	402	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	403	3	-	0/2/19/22	0/1/1/1
3	NAG	A	404	1	-	0/6/23/26	0/1/1/1
5	FUC	A	405	3	-	0/0/17/20	0/1/1/1
3	NAG	A	406	1,3,5	-	0/6/23/26	0/1/1/1
3	NAG	A	407	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	408	3	-	0/2/19/22	0/1/1/1
6	GOL	G	401	-	-	0/4/4/4	0/0/0/0
6	GOL	G	402	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	G	403	-	-	0/4/4/4	0/0/0/0
5	FUC	G	404	3	-	0/0/17/20	0/1/1/1
3	NAG	G	405	1,3,5	-	0/6/23/26	0/1/1/1
3	NAG	G	406	3,4	-	0/6/23/26	0/1/1/1
4	BMA	G	407	3	-	0/2/19/22	0/1/1/1
3	NAG	G	408	1	-	0/6/23/26	0/1/1/1
3	NAG	G	409	1	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	408	NAG	C1-C2	2.36	1.55	1.52

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	407	NAG	O5-C1-C2	-3.89	106.15	111.52
3	A	406	NAG	O5-C1-C2	-2.90	107.52	111.52
3	G	406	NAG	O5-C1-C2	-2.78	107.68	111.52
3	G	409	NAG	O5-C1-C2	-2.55	108.00	111.52
4	A	403	BMA	O5-C1-C2	-2.49	106.89	110.78
3	A	401	NAG	C3-C4-C5	-2.31	106.11	110.24
3	G	405	NAG	C1-C2-N2	-2.03	107.02	110.49
3	G	409	NAG	C1-O5-C5	2.04	114.99	112.19
4	A	403	BMA	C1-O5-C5	2.17	115.18	112.19
3	G	408	NAG	C2-N2-C7	2.19	126.14	122.94
4	A	408	BMA	O5-C5-C6	2.20	110.63	107.15
3	A	404	NAG	O5-C5-C6	2.22	110.66	107.15
3	G	408	NAG	C4-C3-C2	2.26	114.33	111.02
3	G	405	NAG	C4-C3-C2	2.29	114.37	111.02
3	A	406	NAG	C8-C7-N2	2.41	120.33	116.10
5	A	405	FUC	C1-O5-C5	2.43	117.76	112.39
4	G	407	BMA	C1-O5-C5	2.53	115.67	112.19
3	G	406	NAG	C3-C4-C5	2.61	114.90	110.24
3	A	406	NAG	C6-C5-C4	2.65	119.24	112.99
3	G	408	NAG	O5-C5-C6	2.73	111.47	107.15
3	A	406	NAG	C2-N2-C7	2.74	126.94	122.94
3	A	407	NAG	C1-O5-C5	3.35	116.79	112.19
3	A	406	NAG	C1-O5-C5	3.38	116.84	112.19
4	A	403	BMA	O5-C5-C6	3.42	112.57	107.15
5	G	404	FUC	O5-C1-C2	3.75	116.62	110.78
5	G	404	FUC	C1-O5-C5	4.26	121.80	112.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	NAG	C4-C3-C2	4.78	118.03	111.02
5	G	404	FUC	C1-C2-C3	4.99	115.98	109.66
3	A	404	NAG	C1-O5-C5	5.52	119.78	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/288 (89%)	0.61	26 (10%) 7 5	20, 43, 84, 97	0
1	G	265/288 (92%)	0.29	7 (2%) 56 56	19, 35, 71, 86	0
2	B	105/122 (86%)	0.25	3 (2%) 51 51	25, 43, 71, 102	0
2	I	108/122 (88%)	0.25	3 (2%) 53 53	22, 40, 66, 103	0
All	All	737/820 (89%)	0.39	39 (5%) 26 24	19, 40, 79, 103	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	42	ASP	4.8
1	A	41	TRP	4.4
1	A	28	THR	4.0
1	G	89	PRO	3.9
1	A	221	HIS	3.9
1	A	47	VAL	3.9
1	A	85	VAL	3.9
1	A	90	PHE	3.8
1	A	46	ASN	3.6
1	A	92	THR	3.5
1	A	223	PHE	3.4
1	A	245	ARG	3.3
2	I	35	LEU	3.2
1	A	266	GLY	3.2
1	A	29	ASN	2.9
1	G	44	ASN	2.9
1	A	45	ARG	2.9
1	A	249	VAL	2.9
1	G	43	LEU	2.8
1	A	91	SER	2.7
1	A	220	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	35	LEU	2.7
2	I	34	LEU	2.7
1	G	47	VAL	2.6
1	A	40	THR	2.5
1	A	250	ILE	2.5
1	A	39	LEU	2.5
1	A	65	ASN	2.5
2	B	29	GLN	2.4
1	A	240	LEU	2.4
1	G	90	PHE	2.4
1	A	222	SER	2.3
2	B	36	ASP	2.3
2	I	29	GLN	2.2
1	A	146	ARG	2.2
1	G	27	ILE	2.2
1	A	244	LYS	2.1
1	G	46	ASN	2.0
1	A	147	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	A	403	11/12	0.19	0.48	76,84,89,93	0
4	BMA	A	408	11/12	0.45	0.34	73,84,88,89	0
3	NAG	A	402	14/15	0.68	0.29	75,83,86,87	0
4	BMA	G	407	11/12	0.69	0.24	79,83,84,86	0
3	NAG	A	401	14/15	0.72	0.23	75,78,80,80	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	FUC	A	405	10/11	0.72	0.26	77,82,84,84	0
3	NAG	A	404	14/15	0.73	0.23	62,67,69,70	0
6	GOL	G	403	6/6	0.76	0.27	58,60,61,61	0
3	NAG	G	409	14/15	0.76	0.20	60,63,64,66	0
3	NAG	G	408	14/15	0.78	0.23	66,72,75,75	0
3	NAG	A	406	14/15	0.78	0.21	64,70,74,76	0
3	NAG	A	407	14/15	0.80	0.24	72,76,77,80	0
3	NAG	G	406	14/15	0.80	0.18	66,73,78,80	0
6	GOL	G	401	6/6	0.85	0.41	37,39,41,41	0
5	FUC	G	404	10/11	0.90	0.17	62,63,64,64	0
3	NAG	G	405	14/15	0.90	0.16	53,57,63,66	0
6	GOL	G	402	6/6	0.90	0.35	48,50,50,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.