



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 24, 2023 – 07:16 PM EDT

PDB ID : 5UWR
Title : Crystal Structure of CDC7 NES Peptide (extended) in complex with CRM1-Ran-RanBP1
Authors : Fung, H.Y.J.; Chook, Y.M.
Deposited on : 2017-02-21
Resolution : 2.24 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	237	 2% 84% 14%
2	B	143	 3% 86% 11%
3	C	1024	 3% 94% 2%
4	D	27	 22% 41% 56%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 22645 atoms, of which 11112 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	204	3314	1068	1661	283	296	6	0	4	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP P62826
A	-19	GLU	-	expression tag	UNP P62826
A	-18	THR	-	expression tag	UNP P62826
A	-17	GLY	-	expression tag	UNP P62826
A	-16	SER	-	expression tag	UNP P62826
A	-15	SER	-	expression tag	UNP P62826
A	-14	HIS	-	expression tag	UNP P62826
A	-13	HIS	-	expression tag	UNP P62826
A	-12	HIS	-	expression tag	UNP P62826
A	-11	HIS	-	expression tag	UNP P62826
A	-10	HIS	-	expression tag	UNP P62826
A	-9	HIS	-	expression tag	UNP P62826
A	-8	SER	-	expression tag	UNP P62826
A	-7	SER	-	expression tag	UNP P62826
A	-6	GLY	-	expression tag	UNP P62826
A	-5	LEU	-	expression tag	UNP P62826
A	-4	PRO	-	expression tag	UNP P62826
A	-3	ARG	-	expression tag	UNP P62826
A	-2	GLY	-	expression tag	UNP P62826
A	-1	SER	-	expression tag	UNP P62826
A	0	HIS	-	expression tag	UNP P62826

- Molecule 2 is a protein called Ran-specific GTPase-activating protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	127	2107	669	1052	186	195	5	0	2	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	59	GLY	-	expression tag	UNP P41920
B	60	GLY	-	expression tag	UNP P41920
B	61	SER	-	expression tag	UNP P41920

- Molecule 3 is a protein called Exportin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	C	1000	16402	5233	8252	1346	1529	42	0	19	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	GLY	-	expression tag	UNP P30822
C	-1	GLY	-	expression tag	UNP P30822
C	0	SER	-	expression tag	UNP P30822
C	441	ASP	VAL	conflict	UNP P30822
C	537	GLY	ASP	conflict	UNP P30822
C	539	CYS	THR	conflict	UNP P30822
C	540	GLU	VAL	conflict	UNP P30822
C	541	GLN	LYS	conflict	UNP P30822
C	1022	CYS	TYR	conflict	UNP P30822


- Molecule 4 is a protein called Cell division cycle 7-related protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
4	D	12	212	61	111	22	16	2	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	452	GLY	-	expression tag	UNP O00311
D	453	GLY	-	expression tag	UNP O00311
D	454	SER	-	expression tag	UNP O00311
D	455	TYR	-	expression tag	UNP O00311

- Molecule 5 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: $C_{10}H_{17}N_6O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
5	A	1	44	10	12	6	13	3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
6	A	1	1	1	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			14	3	8	3		
7	C	1	Total	C	H	O	0	0
			14	3	8	3		
7	C	1	Total	C	H	O	0	0
			14	3	8	3		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	1	Total	Cl	0	0
			1	1		



- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	121	Total	O	0	0
			121	121		
9	B	18	Total	O	0	0
			18	18		
9	C	383	Total	O	0	0
			383	383		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: GTP-binding nuclear protein Ran





- Molecule 2: Ran-specific GTPase-activating protein 1



- Molecule 3: Exportin-1



- Molecule 4: Cell division cycle 7-related protein kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.08Å 106.08Å 304.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.44 – 2.24 47.44 – 2.24	Depositor EDS
% Data completeness (in resolution range)	93.6 (47.44-2.24) 93.6 (47.44-2.24)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 2.24Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.187 , 0.222 0.187 , 0.222	Depositor DCC
R_{free} test set	2000 reflections (2.53%)	wwPDB-VP
Wilson B-factor (Å ²)	29.5	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	22645	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GNP, CL, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1711	0.46	0/2318
2	B	0.25	0/1086	0.44	0/1449
3	C	0.24	0/8386	0.41	0/11357
4	D	0.23	0/100	0.52	0/129
All	All	0.25	0/11283	0.42	0/15253

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1653	1661	1641	4	0
2	B	1055	1052	1042	2	0
3	C	8150	8252	8169	20	0
4	D	101	111	111	1	0
5	A	32	12	12	0	0
6	A	1	0	0	0	0
7	A	6	8	8	1	0
7	C	12	16	16	0	0
8	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	121	0	0	2	0
9	B	18	0	0	0	0
9	C	383	0	0	6	1
All	All	11533	11112	10999	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:653:SER:O	3:C:656:ARG:NH1	2.19	0.75
1:A:9:VAL:N	9:A:402:HOH:O	2.19	0.74
3:C:127:LYS:O	3:C:178:LYS:NZ	2.24	0.69
3:C:11:LEU:O	9:C:1201:HOH:O	2.11	0.68
1:A:77:ASP:OD2	9:A:401:HOH:O	2.11	0.67
3:C:851:GLU:OE1	9:C:1202:HOH:O	2.12	0.67
1:A:70:GLU:O	1:A:76:ARG:NH1	2.33	0.62
3:C:894:LYS:NZ	3:C:938:GLU:OE1	2.35	0.59
3:C:897:ASN:ND2	9:C:1213:HOH:O	2.36	0.58
3:C:992:SER:OG	9:C:1203:HOH:O	2.16	0.56
1:A:107:ASP:OD2	7:A:303:GOL:O3	2.23	0.56
3:C:439:GLU:O	3:C:440:GLU:HB2	2.10	0.52
3:C:262:LEU:O	3:C:325:ARG:NH1	2.38	0.48
2:B:116:ASN:OD1	2:B:118:LYS:N	2.42	0.47
3:C:264:ILE:O	3:C:266:GLN:NE2	2.45	0.47
3:C:90:ILE:HA	3:C:93:PHE:CE2	2.50	0.47
3:C:91:ARG:NH1	9:C:1225:HOH:O	2.47	0.46
2:B:92:LYS:HD2	2:B:94:PHE:CZ	2.51	0.46
3:C:74:LEU:HD11	3:C:82:LEU:HD11	1.97	0.45
3:C:138:ILE:HB	3:C:139:PRO:HD3	1.99	0.45
3:C:570:TRP:CG	3:C:614:GLN:HG2	2.51	0.45
3:C:72[B]:ASP:OD2	9:C:1204:HOH:O	2.21	0.44
3:C:75:ILE:O	3:C:79:TRP:HB2	2.18	0.43
3:C:540:GLU:OE2	4:D:464:ARG:NH1	2.53	0.41
3:C:981:VAL:HG13	3:C:982:PRO:HD2	2.03	0.41
3:C:460:ASP:N	3:C:460:ASP:OD1	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:C:1450:HOH:O	9:C:1522:HOH:O[6_455]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/237 (86%)	199 (98%)	5 (2%)	0	100	100
2	B	125/143 (87%)	119 (95%)	6 (5%)	0	100	100
3	C	1015/1024 (99%)	996 (98%)	19 (2%)	0	100	100
4	D	10/27 (37%)	9 (90%)	1 (10%)	0	100	100
All	All	1354/1431 (95%)	1323 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/203 (89%)	179 (99%)	1 (1%)	86	90
2	B	111/122 (91%)	111 (100%)	0	100	100
3	C	932/933 (100%)	926 (99%)	6 (1%)	86	90
4	D	11/24 (46%)	11 (100%)	0	100	100
All	All	1234/1282 (96%)	1227 (99%)	7 (1%)	86	90

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	TYR
3	C	35	GLN
3	C	100	SER
3	C	291	MET
3	C	656	ARG
3	C	940	PHE
3	C	992	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	GOL	A	303	-	5,5,5	0.37	0	5,5,5	0.17	0
5	GNP	A	301	6	29,34,34	4.43	9 (31%)	33,54,54	1.67	6 (18%)
7	GOL	C	1102	-	5,5,5	0.35	0	5,5,5	0.26	0
7	GOL	C	1101	-	5,5,5	0.37	0	5,5,5	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	303	-	-	2/4/4/4	-
5	GNP	A	301	6	-	2/14/38/38	0/3/3/3
7	GOL	C	1102	-	-	0/4/4/4	-
7	GOL	C	1101	-	-	2/4/4/4	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	301	GNP	C2'-C1'	-15.05	1.30	1.53
5	A	301	GNP	O4'-C1'	14.84	1.61	1.41
5	A	301	GNP	O4'-C4'	-6.30	1.30	1.45
5	A	301	GNP	C2-N2	4.11	1.42	1.33
5	A	301	GNP	PG-O1G	3.49	1.51	1.46
5	A	301	GNP	PB-O1B	3.35	1.51	1.46
5	A	301	GNP	O2'-C2'	2.95	1.49	1.43
5	A	301	GNP	O3'-C3'	-2.81	1.36	1.43
5	A	301	GNP	PG-O3G	-2.00	1.51	1.56

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	301	GNP	N3-C2-N1	-5.10	120.42	127.22
5	A	301	GNP	C2-N3-C4	4.02	119.95	115.36
5	A	301	GNP	C5-C6-N1	-2.70	119.74	123.43
5	A	301	GNP	C2-N1-C6	2.59	120.05	115.93
5	A	301	GNP	PB-O3A-PA	-2.15	125.06	132.62
5	A	301	GNP	N2-C2-N1	2.10	120.53	117.25

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	301	GNP	PG-N3B-PB-O1B
5	A	301	GNP	PG-N3B-PB-O3A
7	A	303	GOL	C1-C2-C3-O3
7	A	303	GOL	O2-C2-C3-O3
7	C	1101	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	C	1101	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	303	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	204/237 (86%)	0.01	4 (1%) 65 66	17, 32, 75, 113	0
2	B	127/143 (88%)	0.18	5 (3%) 39 39	35, 52, 80, 104	0
3	C	1000/1024 (97%)	-0.04	26 (2%) 56 57	17, 38, 72, 138	0
4	D	12/27 (44%)	1.95	6 (50%) 0 0	90, 93, 107, 111	0
All	All	1343/1431 (93%)	0.01	41 (3%) 49 49	17, 39, 76, 138	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	264	ILE	6.4
3	C	543	ARG	5.8
1	A	195	ALA	4.7
4	D	465	LEU	3.8
3	C	262	LEU	3.5
3	C	978	GLU	3.4
2	B	168	GLU	3.4
3	C	539	CYS	3.3
3	C	545	LYS	3.2
2	B	79	MET	3.1
3	C	270	LEU	3.0
3	C	226	TYR	3.0
2	B	199	LYS	3.0
4	D	460	LYS	3.0
3	C	967	TYR	2.9
4	D	468	MET	2.9
4	D	459	ARG	2.9
3	C	977	GLN	2.9
3	C	588	HIS	2.9
3	C	544	GLY	2.8
3	C	591	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	182	LEU	2.7
3	C	541	GLN	2.7
1	A	197	TYR	2.7
3	C	589	GLU	2.4
3	C	682	GLN	2.4
3	C	538	LEU	2.3
3	C	690	LEU	2.3
4	D	461	LEU	2.3
3	C	583	PHE	2.3
3	C	502	TRP	2.2
3	C	969	ASN	2.2
3	C	540	GLU	2.1
3	C	587	THR	2.1
2	B	122	VAL	2.1
3	C	616	ARG	2.1
1	A	179	MET	2.0
2	B	196	GLU	2.0
3	C	503	HIS	2.0
4	D	458	LEU	2.0
3	C	546	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	GOL	C	1102	6/6	0.77	0.19	47,57,61,64	0
7	GOL	C	1101	6/6	0.80	0.17	46,55,59,62	0
7	GOL	A	303	6/6	0.84	0.18	45,54,59,59	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	MG	A	302	1/1	0.96	0.10	23,23,23,23	0
8	CL	C	1103	1/1	0.96	0.12	52,52,52,52	0
5	GNP	A	301	32/32	0.98	0.14	20,25,33,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.