

Full wwPDB X-ray Structure Validation Report (i)

May 29, 2020 – 09:15 am BST

PDB ID : 5UXZ

Title: X-ray crystal structure of Halotag bound to the P9 benzothiadiazole fluoro-

genic ligand

Authors : Dunham, N.P.; Boal, A.K.

Deposited on : 2017-02-23

Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	307	92%	• 6%			
1	В	307	24% 89%	7% •			



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4827 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Haloalkane dehalogenase.

Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	A	290	Total 2328	C 1515	± 1	O 412	S 9	0	0	0
1	В	295	Total 2363	C 1537		O 420	S 9	0	0	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Chain Residue M		Actual	Comment	Reference
A	47	VAL	LEU	conflict	UNP P0A3G2
A	58	THR	SER	conflict	UNP P0A3G2
A	78	GLY	ASP	conflict	UNP P0A3G2
A	87	PHE	TYR	conflict	UNP P0A3G2
A	88	MET	LEU	conflict	UNP P0A3G2
A	128	PHE	CYS	conflict	UNP P0A3G2
A	155	THR	ALA	conflict	UNP P0A3G2
A	160	LYS	GLU	conflict	UNP P0A3G2
A	167	VAL	ALA	conflict	UNP P0A3G2
A	172	THR	ALA	conflict	UNP P0A3G2
A	175	MET	LYS	conflict	UNP P0A3G2
A	176	GLY	CYS	conflict	UNP P0A3G2
A	195	ASN	LYS	conflict	UNP P0A3G2
A	224	GLU	ALA	conflict	UNP P0A3G2
A	227	ASP	ASN	conflict	UNP P0A3G2
A	257	LYS	GLU	conflict	UNP P0A3G2
A	264	ALA	THR	conflict	UNP P0A3G2
A	272	ASN	HIS	conflict	UNP P0A3G2
A	273	LEU	TYR	conflict	UNP P0A3G2
A	291	SER	=	expression tag	UNP P0A3G2
A	293	THR	=	expression tag	UNP P0A3G2
A	294	LEU	-	expression tag	UNP P0A3G2
A	295	GLU	=	expression tag	UNP P0A3G2
A	296	ILE	=	expression tag	UNP P0A3G2
A	297	SER	_	expression tag	UNP P0A3G2



Continued from previous page...


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	298	GLY	-	expression tag	UNP P0A3G2
A	299	LEU	-	expression tag	UNP P0A3G2
A	300	GLU	-	expression tag	UNP P0A3G2
A	301	ILE	-	expression tag	UNP P0A3G2
A	302	SER	-	expression tag	UNP P0A3G2
A	303	GLY	-	expression tag	UNP P0A3G2
A	304	LEU	-	expression tag	UNP P0A3G2
A	305	GLU	-	expression tag	UNP P0A3G2
A	306	HIS	-	expression tag	UNP P0A3G2
A	307	HIS	-	expression tag	UNP P0A3G2
A	308	HIS	-	expression tag	UNP P0A3G2
A	309	HIS	-	expression tag	UNP P0A3G2
A	310	HIS	-	expression tag	UNP P0A3G2
A	311	HIS	-	expression tag	UNP P0A3G2
В	47	VAL	LEU	conflict	UNP P0A3G2
В	58	THR	SER	conflict	UNP P0A3G2
В	78	GLY	ASP	conflict	UNP P0A3G2
В	87	PHE	TYR	conflict	UNP P0A3G2
В	88	MET	LEU	conflict	UNP P0A3G2
В	128	PHE	CYS	conflict	UNP P0A3G2
В	155	THR	ALA	conflict	UNP P0A3G2
В	160	LYS	GLU	conflict	UNP P0A3G2
В	167	VAL	ALA	conflict	UNP P0A3G2
В	172	THR	ALA	conflict	UNP P0A3G2
В	175	MET	LYS	conflict	UNP P0A3G2
В	176	GLY	CYS	conflict	UNP P0A3G2
В	195	ASN	LYS	conflict	UNP P0A3G2
В	224	GLU	ALA	conflict	UNP P0A3G2
В	227	ASP	ASN	conflict	UNP P0A3G2
В	257	LYS	GLU	conflict	UNP P0A3G2
В	264	ALA	THR	conflict	UNP P0A3G2
В	272	ASN	HIS	conflict	UNP P0A3G2
В	273	LEU	TYR	conflict	UNP P0A3G2
В	291	SER	-	expression tag	UNP P0A3G2
В	292	THR	_	expression tag	UNP P0A3G2
В	293	LEU	-	expression tag	UNP P0A3G2
В	294	GLU	-	expression tag	UNP P0A3G2
В	295	ILE	-	expression tag	UNP P0A3G2
В	296	SER	-	expression tag	UNP P0A3G2
В	297	GLY	-	expression tag	UNP P0A3G2
В	298	LEU	-	expression tag	UNP P0A3G2
В	299	GLU	-	expression tag	UNP P0A3G2
	l	<u> </u>	<u> </u>		l on mort nage



~ · · ·	e		
Continued	trom	previous	paae

Chain	Residue	Modelled	Actual	Comment	Reference
В	300	ILE	_	expression tag	UNP P0A3G2
В	301	SER	-	expression tag	UNP P0A3G2
В	302	GLY	_	expression tag	UNP P0A3G2
В	303	LEU	-	expression tag	UNP P0A3G2
В	304	GLU	_	expression tag	UNP P0A3G2
В	305	HIS	_	expression tag	UNP P0A3G2
В	306	HIS	_	expression tag	UNP P0A3G2
В	307	HIS	_	expression tag	UNP P0A3G2
В	308	HIS	-	expression tag	UNP P0A3G2
В	309	HIS	-	expression tag	UNP P0A3G2
В	310	HIS	-	expression tag	UNP P0A3G2

• Molecule 2 is N 2 -{[7-(dimethylamino)-2,1,3-benzothiadiazol-4-yl]sulfonyl}-N-hexyl-N 2 -m ethylglycinamide (three-letter code: 8PM) (formula: $C_{17}H_{27}N_5O_3S_2$).



Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf
2	Λ	1	Total	С	N	О	S	0	0
	A	1	27	17	5	3	2	0	
9	D	1	Total	С	N	О	S	0	0
	D		27	17	5	3	2		

• Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Cl 1 1	0	0



Continued from previous page...

Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

 \bullet Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

• Molecule 5 is water.


\mathbf{Mol}	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
5	A	53	Total O 53 53	0	0
5	В	19	Total O 19 19	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Haloalkane dehalogenase







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	49.94Å 69.12Å 83.90Å	Depositor
a, b, c, α , β , γ	90.00° 94.67° 90.00°	Depositor
Resolution (Å)	50.00 - 1.92	Depositor
resolution (A)	49.77 - 1.92	EDS
% Data completeness	81.9 (50.00-1.92)	Depositor
(in resolution range)	81.9 (49.77-1.92)	EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.44 (at 1.92Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
P. P.	0.269 , 0.287	Depositor
R, R_{free}	0.274 , 0.293	DCC
R_{free} test set	1806 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 44.9	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4827	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 8PM, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.33	$1/2407 \ (0.0\%)$	0.46	0/3292	
1	В	0.33	$1/2442 \ (0.0\%)$	0.45	0/3339	
All	All	0.33	$2/4849 \ (0.0\%)$	0.45	0/6631	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathbf{Ideal}(\mathbf{\mathring{A}})$
1	В	106	ASP	CG-OD1	8.96	1.46	1.25
1	A	106	ASP	CG-OD1	8.70	1.45	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2328	0	2271	2	0
1	В	2363	0	2307	9	0
2	A	27	0	0	0	0
2	В	27	0	0	1	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	8	0	12	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	53	0	0	0	0
5	В	19	0	0	0	0
All	All	4827	0	4590	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{\AA}) \end{array}$	Clash overlap (Å)
1:B:285:GLU:OE1	1:B:288:ARG:NH1	2.37	0.57
1:A:275:GLN:H	1:A:275:GLN:HE21	1.53	0.56
1:B:275:GLN:HE21	1:B:275:GLN:H	1.59	0.51
1:B:200:GLU:N	1:B:201:PRO:HD2	2.29	0.46
1:B:245:VAL:HG11	2:B:401:8PM:O24	2.17	0.44
1:B:193:PHE:CZ	1:B:202:LEU:HD21	2.53	0.44
1:B:56:ALA:N	1:B:57:PRO:CD	2.82	0.43
1:B:64:PRO:HB3	1:B:91:PHE:CZ	2.54	0.42
1:B:42:PRO:HB3	1:B:202:LEU:HD22	2.02	0.42
1:B:107:TRP:CD2	1:B:132:ILE:CD1	3.05	0.40
1:A:200:GLU:N	1:A:201:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	A	288/307~(94%)	275 (96%)	13 (4%)	0	100	100
1	В	293/307~(95%)	278 (95%)	14 (5%)	1 (0%)	41	31
All	All	581/614 (95%)	553 (95%)	27 (5%)	1 (0%)	47	38



All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	166	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outliers		Percentiles	
1	A	250/265~(94%)	246 (98%)	4 (2%)	62 58
1	В	254/265~(96%)	249 (98%)	5 (2%)	55 49
All	All	$504/530 \ (95\%)$	495 (98%)	9 (2%)	59 53

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	ARG
1	A	156	ASP
1	A	275	GLN
1	A	278	ASN
1	В	139	ASP
1	В	168	PHE
1	В	275	GLN
1	В	278	ASN
1	В	298	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	272	ASN
1	A	275	GLN
1	A	278	ASN
1	В	217	ASN
1	В	272	ASN
1	В	275	GLN
1	В	278	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain Re		Res	Link	Bond lengths			Bond angles															
10101	Type	Chain		Chain	Chain	Chain	Chain	Chain	Chain	nes	nes	nes	nes	res	res Li	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	404	-	3,3,3	0.45	0	2,2,2	0.26	0												
2	8PM	A	401	1	27,28,28	3.20	5 (18%)	32,39,39	2.06	8 (25%)												
2	8PM	В	401	1	27,28,28	3.14	5 (18%)	32,39,39	2.04	9 (28%)												
4	EDO	A	403	-	3,3,3	0.50	0	2,2,2	0.13	0												

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	${f Res}$	Link	Chirals	Torsions	Rings
4	EDO	A	404	-	1	1/1/1/1	-
2	8PM	A	401	1	-	11/27/27/27	0/2/2/2
2	8PM	В	401	1	-	7/27/27/27	0/2/2/2
4	EDO	A	403	_	-	0/1/1/1	-



All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	${ m Observed}({ m \AA})$	$\mathbf{Ideal}(\mathbf{\AA})$
2	A	401	8PM	C2-S1	-12.20	1.60	1.77
2	В	401	8PM	C2-S1	-11.63	1.61	1.77
2	В	401	8PM	C2-C3	9.59	1.49	1.42
2	A	401	8PM	C2-C3	9.50	1.49	1.42
2	В	401	8PM	C5-C4	4.45	1.50	1.42
2	A	401	8PM	C5-C4	4.43	1.50	1.42
2	В	401	8PM	C7-C2	2.45	1.40	1.37
2	В	401	8PM	C4-C3	2.37	1.51	1.43
2	A	401	8PM	C4-C3	2.29	1.51	1.43
2	A	401	8PM	C7-C2	2.14	1.40	1.37

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^o)$
2	В	401	8PM	O23-S1-O22	-7.42	107.50	119.52
2	A	401	8PM	O23-S1-O22	-7.24	107.79	119.52
2	A	401	8PM	O22-S1-N14	4.63	112.83	107.05
2	В	401	8PM	C2-S1-N14	4.38	117.12	106.52
2	A	401	8PM	O23-S1-N14	3.45	111.35	107.05
2	В	401	8PM	C10-N8-C5	2.63	121.94	114.12
2	A	401	8PM	C7-C2-S1	2.63	120.83	116.97
2	В	401	8PM	C3-N12-S13	2.45	107.17	104.97
2	A	401	8PM	C4-N11-S13	2.45	107.17	104.97
2	В	401	8PM	C4-N11-S13	2.42	107.14	104.97
2	A	401	8PM	C9-N8-C5	2.38	121.20	114.12
2	A	401	8PM	C10-N8-C5	2.32	121.03	114.12
2	A	401	8PM	C3-N12-S13	2.23	106.97	104.97
2	В	401	8PM	O24-C16-N17	-2.23	118.81	123.01
2	В	401	8PM	C15-C16-N17	2.18	120.48	115.52
2	В	401	8PM	C9-N8-C5	2.14	120.49	114.12
2	В	401	8PM	C18-N17-C16	2.09	126.71	122.84

There are no chirality outliers.

All (19) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	401	8PM	C16-C15-N14-C21
2	A	401	8PM	C21-N14-S1-O22
2	В	401	8PM	O24-C16-N17-C18
2	A	401	8PM	N17-C18-C19-C20
2	В	401	8PM	C15-C16-N17-C18



Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	401	8PM	C4-C5-N8-C9
2	A	401	8PM	C21-N14-S1-C2
2	В	401	8PM	C18-C19-C20-C25
2	В	401	8PM	C4-C5-N8-C10
2	A	401	8PM	C6-C5-N8-C9
2	В	401	8PM	C6-C5-N8-C10
2	A	401	8PM	C19-C20-C25-C26
4	A	404	EDO	O1-C1-C2-O2
2	В	401	8PM	C20-C25-C26-C27
2	A	401	8PM	C15-N14-S1-O23
2	A	401	8PM	C18-C19-C20-C25
2	A	401	8PM	C15-N14-S1-C2
2	A	401	8PM	C15-N14-S1-O22
2	A	401	8PM	C21-N14-S1-O23


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	401	8PM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





Rings

5.7 Other polymers (i)

There are no such residues in this entry.

Torsions

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	290/307~(94%)	0.81	32 (11%) 5 6	5, 17, 37, 41	35 (12%)
1	В	295/307~(96%)	1.44	75 (25%) 0 0	15, 32, 51, 57	35 (11%)
All	All	585/614 (95%)	1.13	107 (18%) 1 1	5, 25, 48, 57	70 (11%)

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	193	PHE	7.8
1	В	194	LEU	6.6
1	В	203	TRP	6.1
1	В	140	GLU	5.3
1	A	157	VAL	5.2
1	В	298	LEU	5.0
1	В	74	LYS	5.0
1	В	156	ASP	4.5
1	В	137	THR	4.4
1	В	211	ILE	4.4
1	В	212	ALA	4.3
1	A	261	ASN	4.3
1	В	157	VAL	4.2
1	A	144	PHE	4.1
1	В	77	LEU	4.0
1	В	31	ASP	3.8
1	В	165	GLN	3.8
1	A	256	ALA	3.7
1	В	76	ASP	3.7
1	В	17	VAL	3.5
1	В	167	VAL	3.5
1	В	80	PHE	3.4
1	A	203	TRP	3.3
1	A	148	THR	3.3



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	В	21	ARG	3.3
1	В	152	PHE	3.3
1	В	215	PRO	3.3
1	A	135	ILE	3.2
1	В	138	TRP	3.2
1	В	155	THR	3.2
1	В	161	LEU	3.2
1	В	143	GLU	3.1
1	В	160	LYS	3.1
1	В	56	ALA	3.1
1	В	196	PRO	3.1
1	В	14	TYR	3.1
1	A	227	ASP	3.1
1	В	162	ILE	3.1
1	В	198	ASP	3.0
1	A	221	LEU	3.0
1	В	90	ALA	3.0
1	В	134	PRO	3.0
1	В	154	THR	2.9
1	В	79	TYR	2.9
1	В	58	THR	2.9
1	A	254	ARG	2.9
1	В	144	PHE	2.9
1	В	19	GLY	2.9
1	В	207	ASN	2.9
1	В	12	PRO	2.8
1	В	188	HIS	2.8
1	A	215	PRO	2.8
1	A	155	THR	2.8
1	В	94	ALA	2.7
1	В	288	ARG	2.7
1	A	139	ASP	2.7
1	В	158	GLY	2.7
1	В	142	PRO	2.7
1	В	139	ASP	2.6
1	В	141	TRP	2.6
1	A	288	ARG	2.6
1	В	29	PRO	2.6
1	A	156	ASP	2.5
1	A	151	ALA	2.5
1	A	213	GLY	2.5
1	A	229	LEU	2.5



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	RSRZ
1	В	192	PRO	2.5
1	В	151	ALA	2.5
1	A	138	TRP	2.4
1	В	27	VAL	2.4
1	A	160	LYS	2.4
1	В	135	ILE	2.4
1	A	226	MET	2.3
1	В	250	ALA	2.3
1	A	211	ILE	2.3
1	A	231	GLN	2.3
1	В	254	ARG	2.3
1	В	33	THR	2.3
1	В	136	PRO	2.3
1	A	188	HIS	2.3
1	В	297	GLY	2.3
1	A	219	VAL	2.3
1	A	294	LEU	2.3
1	В	149	PHE	2.2
1	В	75	PRO	2.2
1	В	30	ARG	2.2
1	В	57	PRO	2.2
1	В	197	VAL	2.2
1	A	293	THR	2.2
1	A	152	PHE	2.2
1	A	163	ILE	2.2
1	В	28	GLY	2.2
1	В	95	LEU	2.1
1	В	15	VAL	2.1
1	В	71	LYS	2.1
1	В	169	ILE	2.1
1	В	230	HIS	2.1
1	В	16	GLU	2.1
1	В	213	GLY	2.1
1	В	227	ASP	2.1
1	В	287	ALA	2.1
1	В	209	LEU	2.1
1	В	246	LEU	2.1
1	A	141	TRP	2.0
1	В	146	ARG	2.0
1	A	224	GLU	2.0
1	A	214	GLU	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	8PM	В	401	27/27	0.84	0.18	20,31,33,33	0
2	8PM	A	401	27/27	0.86	0.17	14,23,25,25	0
4	EDO	A	403	4/4	0.89	0.14	17,18,18,18	0
4	EDO	A	404	4/4	0.93	0.24	16,16,16,16	0
3	CL	В	402	1/1	0.94	0.10	26,26,26,26	0
3	CL	A	402	1/1	0.97	0.09	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 8PM B 401: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

