



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 28, 2024 – 06:53 pm BST

PDB ID : 1UYS
Title : Acetyl-CoA carboxylase carboxyltransferase domain in complex with inhibitor haloxyfop
Authors : Zhang, H.; Tweel, B.; Tong, L.
Deposited on : 2004-03-02
Resolution : 2.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	737	
1	B	737	
1	C	737	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 16130 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYL-COA CARBOXYLASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	670	Total 5326	C 3396	N 914	O 998	S 2	Se 16	0	0	1
1	B	670	Total 5326	C 3396	N 914	O 998	S 2	Se 16	0	0	1
1	C	660	Total 5248	C 3343	N 903	O 984	S 2	Se 16	0	0	1

- Molecule 2 is (2R)-2-(4-{[3-chloro-5-(trifluoromethyl)pyridin-2-yl]oxy}phenoxy)propanoic acid (three-letter code: H1L) (formula: C₁₅H₁₁ClF₃NO₄).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	Cl	F	N	O		
2	A	1	Total 24	C 15	Cl 1	F 3	N 1	O 4	0	0
2	B	1	Total 24	C 15	Cl 1	F 3	N 1	O 4	0	0
2	C	1	Total 24	C 15	Cl 1	F 3	N 1	O 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	69	Total 69	O 69	0	0
3	B	50	Total 50	O 50	0	0
3	C	39	Total 39	O 39	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	246.21Å 125.58Å 146.88Å 90.00° 94.08° 90.00°	Depositor
Resolution (Å)	27.42 – 2.80 27.42 – 2.80	Depositor EDS
% Data completeness (in resolution range)	87.6 (27.42-2.80) 94.3 (27.42-2.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.07 (at 2.80Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.218 , 0.252 0.225 , 0.258	Depositor DCC
R_{free} test set	10734 reflections (9.99%)	wwPDB-VP
Wilson B-factor (Å ²)	40.4	Xtrriage
Anisotropy	0.460	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	16130	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: H1L

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/5429	0.67	4/7332 (0.1%)
1	B	0.47	0/5429	0.61	0/7332
1	C	0.46	0/5347	0.63	3/7218 (0.0%)
All	All	0.48	0/16205	0.64	7/21882 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1745	ARG	NE-CZ-NH1	-12.57	114.02	120.30
1	C	1745	ARG	NE-CZ-NH2	11.36	125.98	120.30
1	A	1618	ARG	NE-CZ-NH1	8.45	124.53	120.30
1	C	1745	ARG	CD-NE-CZ	-5.82	115.46	123.60
1	A	1618	ARG	NH1-CZ-NH2	-5.23	113.65	119.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5326	0	5269	199	1
1	B	5326	0	5269	215	0
1	C	5248	0	5187	231	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	24	0	10	0	0
2	B	24	0	10	4	0
2	C	24	0	10	6	0
3	A	69	0	0	4	0
3	B	50	0	0	3	0
3	C	39	0	0	4	0
All	All	16130	0	15755	618	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 618 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1618:ARG:HG2	1:C:1618:ARG:HH11	1.25	1.01
1:C:1637:PRO:HG2	1:C:1638:LEU:HD23	1.40	1.00
1:C:1823:THR:HG22	1:C:1824:LYS:H	1.33	0.93
1:A:2045:ASN:HB2	1:A:2046:ARG:CZ	1.99	0.92
1:A:1582:PHE:HA	1:A:1616:ILE:HG23	1.54	0.90

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1970:TYR:OH	1:A:1970:TYR:OH[2_555]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	664/737 (90%)	602 (91%)	57 (9%)	5 (1%)	16 44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	664/737 (90%)	597 (90%)	57 (9%)	10 (2%)	8	29
1	C	654/737 (89%)	576 (88%)	66 (10%)	12 (2%)	7	24
All	All	1982/2211 (90%)	1775 (90%)	180 (9%)	27 (1%)	9	30

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1991	PRO
1	B	1534	ASP
1	B	1650	ASP
1	B	1731	ARG
1	C	1632	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	567/611 (93%)	530 (94%)	37 (6%)	14	40
1	B	567/611 (93%)	523 (92%)	44 (8%)	10	31
1	C	558/611 (91%)	527 (94%)	31 (6%)	17	47
All	All	1692/1833 (92%)	1580 (93%)	112 (7%)	14	39

5 of 112 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1824	LYS
1	C	2114	GLU
1	B	2043	THR
1	C	2081	GLU
1	C	1801	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 61 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1752	GLN
1	C	1922	GLN
1	B	1944	GLN
1	C	1815	ASN
1	C	2142	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	H1L	C	3000	-	24,25,25	0.94	0	32,36,36	1.42	6 (18%)
2	H1L	B	3000	-	24,25,25	1.25	3 (12%)	32,36,36	1.66	8 (25%)
2	H1L	A	3000	-	24,25,25	1.07	1 (4%)	32,36,36	1.55	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	H1L	C	3000	-	-	5/18/18/18	0/2/2/2
2	H1L	B	3000	-	-	5/18/18/18	0/2/2/2
2	H1L	A	3000	-	-	4/18/18/18	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3000	H1L	C7-C6	2.48	1.43	1.38
2	B	3000	H1L	C12-C11	2.26	1.54	1.49
2	B	3000	H1L	C5-C8	2.24	1.43	1.38
2	A	3000	H1L	O2-C3	-2.11	1.34	1.38

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	3000	H1L	C11-C10-N9	-3.78	119.73	123.34
2	B	3000	H1L	C13-C11-C10	3.65	121.07	118.99
2	B	3000	H1L	C11-C10-N9	-3.33	120.16	123.34
2	C	3000	H1L	C11-C10-N9	-3.19	120.29	123.34
2	C	3000	H1L	O8-C9-N9	3.08	123.56	119.49

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	3000	H1L	O1A-C1-C2-O2
2	A	3000	H1L	C1-C2-O2-C3
2	B	3000	H1L	C1-C2-O2-C3
2	C	3000	H1L	C1-C2-O2-C3
2	B	3000	H1L	O1B-C1-C2-O2


There are no ring outliers.


2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3000	H1L	6	0
2	B	3000	H1L	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	654/737 (88%)	0.28	84 (12%) 9 8	13, 32, 98, 100	0
1	B	654/737 (88%)	0.44	97 (14%) 7 6	15, 37, 99, 100	0
1	C	644/737 (87%)	0.41	91 (14%) 7 6	14, 37, 100, 100	0
All	All	1952/2211 (88%)	0.38	272 (13%) 7 7	13, 36, 100, 100	0

The worst 5 of 272 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2047	LEU	8.6
1	C	2047	LEU	7.7
1	B	2047	LEU	7.7
1	C	1913	ALA	6.7
1	B	1908	ALA	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	H1L	B	3000	24/24	0.86	0.17	44,49,59,71	0
2	H1L	C	3000	24/24	0.86	0.17	52,55,60,71	0
2	H1L	A	3000	24/24	0.90	0.13	29,38,45,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around H1L C 3000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around H1L A 3000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.