



wwPDB EM Validation Summary Report ⓘ

Feb 27, 2024 – 08:09 AM EST

PDB ID : 6UZZ
EMDB ID : EMD-20965
Title : structure of human KCNQ1-CaM complex
Authors : Mackinnon, R.; Sun, J.
Deposited on : 2019-11-16
Resolution : 3.10 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	
1	C	557	
1	E	557	
1	G	557	
2	B	149	
2	D	149	
2	F	149	
2	H	149	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 15832 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium voltage-gated channel subfamily KQT member 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	352	2848	1867	497	473	11	0	0
1	C	352	2848	1867	497	473	11	0	0
1	E	352	2848	1867	497	473	11	0	0
1	G	352	2848	1867	497	473	11	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	75	MET	-	expression tag	UNP P51787
A	621	SER	-	expression tag	UNP P51787
A	622	ASN	-	expression tag	UNP P51787
A	623	SER	-	expression tag	UNP P51787
A	624	LEU	-	expression tag	UNP P51787
A	625	GLU	-	expression tag	UNP P51787
A	626	VAL	-	expression tag	UNP P51787
A	627	LEU	-	expression tag	UNP P51787
A	628	PHE	-	expression tag	UNP P51787
A	629	GLN	-	expression tag	UNP P51787
A	630	GLY	-	expression tag	UNP P51787
A	631	PRO	-	expression tag	UNP P51787
C	75	MET	-	expression tag	UNP P51787
C	621	SER	-	expression tag	UNP P51787
C	622	ASN	-	expression tag	UNP P51787
C	623	SER	-	expression tag	UNP P51787
C	624	LEU	-	expression tag	UNP P51787
C	625	GLU	-	expression tag	UNP P51787
C	626	VAL	-	expression tag	UNP P51787
C	627	LEU	-	expression tag	UNP P51787
C	628	PHE	-	expression tag	UNP P51787
C	629	GLN	-	expression tag	UNP P51787

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	630	GLY	-	expression tag	UNP P51787
C	631	PRO	-	expression tag	UNP P51787
E	75	MET	-	expression tag	UNP P51787
E	621	SER	-	expression tag	UNP P51787
E	622	ASN	-	expression tag	UNP P51787
E	623	SER	-	expression tag	UNP P51787
E	624	LEU	-	expression tag	UNP P51787
E	625	GLU	-	expression tag	UNP P51787
E	626	VAL	-	expression tag	UNP P51787
E	627	LEU	-	expression tag	UNP P51787
E	628	PHE	-	expression tag	UNP P51787
E	629	GLN	-	expression tag	UNP P51787
E	630	GLY	-	expression tag	UNP P51787
E	631	PRO	-	expression tag	UNP P51787
G	75	MET	-	expression tag	UNP P51787
G	621	SER	-	expression tag	UNP P51787
G	622	ASN	-	expression tag	UNP P51787
G	623	SER	-	expression tag	UNP P51787
G	624	LEU	-	expression tag	UNP P51787
G	625	GLU	-	expression tag	UNP P51787
G	626	VAL	-	expression tag	UNP P51787
G	627	LEU	-	expression tag	UNP P51787
G	628	PHE	-	expression tag	UNP P51787
G	629	GLN	-	expression tag	UNP P51787
G	630	GLY	-	expression tag	UNP P51787
G	631	PRO	-	expression tag	UNP P51787

- Molecule 2 is a protein called Calmodulin-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	144	Total	C	N	O	S	0	0
			1108	682	180	237	9		
2	D	144	Total	C	N	O	S	0	0
			1108	682	180	237	9		
2	F	144	Total	C	N	O	S	0	0
			1108	682	180	237	9		
2	H	144	Total	C	N	O	S	0	0
			1108	682	180	237	9		


- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
3	B	2	Total 2	Ca 2	0
3	D	2	Total 2	Ca 2	0
3	F	2	Total 2	Ca 2	0
3	H	2	Total 2	Ca 2	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Potassium voltage-gated channel subfamily KQT member 1



- Molecule 1: Potassium voltage-gated channel subfamily KQT member 1



ARG
LEU
ALA
SER
LEU
ILE
THR
ASP
MET
LEU
HIS
GLN
LEU
LEU
SER
LEU
HIS
ASN
SER
LEU
GLU
VAL
THR
ARG
LEU
PHE
GLN
GLY
PRO

- Molecule 1: Potassium voltage-gated channel subfamily KQT member 1

Chain E: 

MET
ALA
SER
LEU
GLY
PRO
ARG
PRO
PRO
VAL
VAL
SER
LEU
ASP
ARG
VAL
ARG
TYR
SER
THR
ARG
ARG
PRO
VAL
VAL
ALA
ARG
ARG
T104
R116
E160
F167
E170
R174
K218
GLY
GLN
VAL
PHE
A223
T224
S225
Q234
R237
H240
R243
T247

E261
T264
F296
T312
K318
I337
F351
Q376
N386
P387
D388
Y395
I396
ARG
LYS
ALA
PRO
ARG
SER
HIS
THR
ILE
VAL
ARG
LEU
SER
PRO
SER
PRO
PRO
LEU
LEU
LEU
VAL
SER
MET
PRO
VAL
VAL
PHE
MET
THR
LYS
PHE
ASN
PHE
SER
LEU
ASP
ASP
ASN
GLY
THR
LEU
GLY
GLU

GLU
LYS
MET
THR
VAL
PRO
HIS
THR
ILE
THR
SER
CYS
ASP
PRO
PRO
GLU
GLU
ARG
ARG
LEU
ASP
HIS
PHE
SER
VAL
VAL
ASP
GLY
TYR
ASP
SER
SER
THR
VAL
ARG
LEU
SER
SER
GLU
LYS
THR
LEU
LEU
GLU
VAL
SER
MET
PRO
HIS
PHE
MET
THR
LYS
ASN
SER
PHE
ALA
GLU
ASP
LYS
LEU
VAL
THR
LEU
GLY
GLU

THR
LEU
THR
THR
PRO
ILE
THR
HIS
ILE
SER
GLN
L506
R507
E508
H509
H510
T513
R561
R565
G568
LYS
PRO
SER
LEU
PHE
ILE
SER
VAL
SER
GLU
LYS
SER
THR
LEU
ASP
ARG
GLY
SER
ASN
THR
ILE
GLY
ALA
ARG
LEU
ASN
VAL
VAL
GLU
ASP
LYS
VAL
THR
GLN
ASP
GLN

ARG
LEU
ALA
SER
LEU
ILE
THR
ASP
MET
LEU
HIS
GLN
LEU
SER
LEU
HIS
ASN
SER
LEU
VAL
THR
ARG
PHE
GLN
GLY
PRO

- Molecule 1: Potassium voltage-gated channel subfamily KQT member 1

Chain G: 

MET
ALA
SER
LEU
GLY
PRO
ARG
PRO
PRO
VAL
VAL
SER
LEU
ASP
ARG
VAL
ARG
TYR
SER
THR
ARG
ARG
PRO
VAL
VAL
ALA
ARG
ARG
T104
R116
E160
F167
E170
R174
K218
GLY
GLN
VAL
PHE
A223
T224
S225
Q234
R237
H240
R243
T247


E261
T264
F296
T312
K318
I337
F351
Q376
N386
P387
D388
Y395
I396
ARG
LYS
ALA
PRO
ARG
SER
HIS
THR
ILE
VAL
ARG
LEU
SER
PRO
SER
PRO
PRO
LEU
LEU
LEU
VAL
SER
MET
PRO
VAL
VAL
PHE
MET
THR
LYS
PHE
ASN
PHE
SER
LEU
ASP
ASP
ASN
GLY
THR
LEU
GLY
GLU

GLU
LYS
MET
THR
VAL
PRO
HIS
THR
ILE
SER
GLN
L506
R507
E508
H509
H510
T513
R561
R565
G568
LYS
PRO
SER
LEU
PHE
ILE
SER
VAL
SER
GLU
LYS
SER
THR
LEU
ASP
ARG
GLY
SER
ASN
THR
ILE
GLY
ALA
ARG
LEU
ASN
VAL
VAL
GLU
ASP
LYS
VAL
THR
GLN
ASP
GLN

THR
LEU
THR
THR
PRO
ILE
THR
HIS
ILE
SER
GLN
L506
R507
E508
H509
H510
T513
R561
R565
G568
LYS
PRO
SER
LEU
PHE
ILE
SER
VAL
SER
GLU
LYS
SER
THR
LEU
ASP
ARG
GLY
SER
ASN
THR
ILE
GLY
ALA
ARG
LEU
ASN
VAL
VAL
GLU
ASP
LYS
VAL
THR
GLN
ASP
GLN


ARG
LEU
ALA
SER
LEU
ILE
THR
ASP
MET
LEU
HIS
GLN
LEU
SER
LEU
HIS
ASN
SER
LEU
VAL
THR
ARG
PHE
GLN
GLY
PRO

- Molecule 2: Calmodulin-1



Chain B: 

MET
ALA
ASP
GLN
LEU
T6
E7
E8
Q9
I10
E15
L19
T30
I53
L70
S82
A89
E128
A129
D130
I131
E132
G133
D134
G135
Q136
V137
K149


- Molecule 2: Calmodulin-1



• Molecule 2: Calmodulin-1



• Molecule 2: Calmodulin-1



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C4	Depositor
Number of particles used	66746	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	94	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	42.193	Depositor
Minimum map value	-21.386	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	5.0	Depositor
Map size (\AA)	309.0, 309.0, 309.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.03, 1.03, 1.03	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2918	0.56	0/3949
1	C	0.47	0/2918	0.56	0/3949
1	E	0.47	0/2918	0.56	0/3949
1	G	0.47	0/2918	0.56	0/3949
2	B	0.38	0/1120	0.55	0/1507
2	D	0.38	0/1120	0.55	0/1507
2	F	0.38	0/1120	0.55	0/1507
2	H	0.38	0/1120	0.55	0/1507
All	All	0.45	0/16152	0.56	0/21824

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2848	0	2918	16	0
1	C	2848	0	2918	15	0
1	E	2848	0	2918	15	0
1	G	2848	0	2918	15	0
2	B	1108	0	1022	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1108	0	1022	7	0
2	F	1108	0	1022	7	0
2	H	1108	0	1022	7	0
3	B	2	0	0	0	0
3	D	2	0	0	0	0
3	F	2	0	0	0	0
3	H	2	0	0	0	0
All	All	15832	0	15760	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 72 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:133:GLY:O	2:D:134:ASP:CB	2.40	0.69
2:F:133:GLY:O	2:F:134:ASP:CB	2.40	0.69
2:H:133:GLY:O	2:H:134:ASP:CB	2.40	0.68
2:B:133:GLY:O	2:B:134:ASP:CB	2.40	0.68
1:A:296:PHE:HE1	1:A:318:LYS:HB3	1.65	0.61

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	346/557 (62%)	340 (98%)	6 (2%)	0	100	100
1	C	346/557 (62%)	340 (98%)	6 (2%)	0	100	100
1	E	346/557 (62%)	340 (98%)	6 (2%)	0	100	100
1	G	346/557 (62%)	340 (98%)	6 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	142/149 (95%)	138 (97%)	3 (2%)	1 (1%)	22	57
2	D	142/149 (95%)	138 (97%)	3 (2%)	1 (1%)	22	57
2	F	142/149 (95%)	138 (97%)	3 (2%)	1 (1%)	22	57
2	H	142/149 (95%)	138 (97%)	3 (2%)	1 (1%)	22	57
All	All	1952/2824 (69%)	1912 (98%)	36 (2%)	4 (0%)	50	79

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	134	ASP
2	D	134	ASP
2	F	134	ASP
2	H	134	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/490 (61%)	298 (99%)	2 (1%)	84	93
1	C	300/490 (61%)	298 (99%)	2 (1%)	84	93
1	E	300/490 (61%)	298 (99%)	2 (1%)	84	93
1	G	300/490 (61%)	298 (99%)	2 (1%)	84	93
2	B	115/127 (91%)	115 (100%)	0	100	100
2	D	115/127 (91%)	115 (100%)	0	100	100
2	F	115/127 (91%)	115 (100%)	0	100	100
2	H	115/127 (91%)	115 (100%)	0	100	100
All	All	1660/2468 (67%)	1652 (100%)	8 (0%)	89	94

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	386	ASN
1	G	116	ARG
1	E	116	ARG
1	C	386	ASN
1	E	386	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	363	HIS
1	E	386	ASN
1	G	386	ASN
1	G	240	HIS
1	C	240	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20965. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 146



Y Index: 154




Z Index: 121


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 5.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 107 nm³; this corresponds to an approximate mass of 97 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20965 and PDB model 6UZZ. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 5.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (5.0).



















9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (5.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7980	 0.5260
A	 0.8200	 0.5350
B	 0.7450	 0.5060
C	 0.8200	 0.5350
D	 0.7420	 0.5060
E	 0.8200	 0.5340
F	 0.7450	 0.5050
G	 0.8200	 0.5350
H	 0.7450	 0.5050

