



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 10, 2023 – 07:04 pm GMT

PDB ID : 1V08  
Title : Crystal structure of the Zea mae beta-glucosidase-1 in complex with glucotetrazole  
Authors : Moriniere, J.; Verdoucq, L.; Bevan, D.R.; Esen, A.; Henrissat, B.; Czjzek, M.  
Deposited on : 2004-03-25  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	512	
1	B	512	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8829 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BETA-GLUCOSIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	491	Total 3967	C 2543	N 657	O 749	S 18	0	0	1
1	B	491	Total 3967	C 2543	N 657	O 749	S 18	0	0	1

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	191	ASP	GLU	engineered mutation	UNP P49235

- Molecule 2 is NOJIRIMYCINE TETRAZOLE (three-letter code: NTZ) (formula: C<sub>6</sub>H<sub>10</sub>N<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 14	C 6	N 4	O 4	0	0
2	A	1	Total 14	C 6	N 4	O 4	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			14	6	4	4		
2	B	1	Total	C	N	O	0	0
			14	6	4	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	450	Total	O	0	0
			450	450		
3	B	389	Total	O	0	0
			389	389		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BETA-GLUCOSIDASE



- Molecule 1: BETA-GLUCOSIDASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.18Å 104.80Å 119.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.75 – 1.90 29.69 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.75-1.90) 96.3 (29.69-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.89 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.173 , 0.205 0.183 , 0.211	Depositor DCC
$R_{free}$ test set	4604 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtrriage
Anisotropy	0.638	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 45.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8829	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NTZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/4088	0.77	14/5551 (0.3%)
1	B	0.43	0/4088	0.75	15/5551 (0.3%)
All	All	0.43	0/8176	0.76	29/11102 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	297	ASP	CB-CG-OD2	7.28	124.85	118.30
1	A	27	ASP	CB-CG-OD2	6.84	124.46	118.30
1	B	54	ASP	CB-CG-OD2	6.80	124.42	118.30
1	B	297	ASP	CB-CG-OD2	6.77	124.39	118.30
1	A	461	ASP	CB-CG-OD2	6.65	124.29	118.30
1	A	411	ASP	CB-CG-OD2	6.62	124.26	118.30
1	B	411	ASP	CB-CG-OD2	6.53	124.17	118.30
1	A	158	ASP	CB-CG-OD2	6.32	123.99	118.30
1	B	158	ASP	CB-CG-OD2	6.29	123.96	118.30
1	B	144	ASP	CB-CG-OD2	6.27	123.94	118.30
1	B	177	ASP	CB-CG-OD2	6.15	123.84	118.30
1	B	27	ASP	CB-CG-OD2	6.12	123.81	118.30
1	B	461	ASP	CB-CG-OD2	6.12	123.80	118.30
1	B	274	ASP	CB-CG-OD2	6.09	123.78	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	144	ASP	CB-CG-OD2	6.05	123.74	118.30
1	A	253	ASP	CB-CG-OD2	6.00	123.70	118.30
1	B	243	ASP	CB-CG-OD2	5.99	123.69	118.30
1	B	355	ASP	CB-CG-OD2	5.88	123.59	118.30
1	B	117	ASP	CB-CG-OD2	5.88	123.59	118.30
1	A	54	ASP	CB-CG-OD2	5.69	123.42	118.30
1	A	479	ASP	CB-CG-OD2	5.57	123.31	118.30
1	B	167	ASP	CB-CG-OD2	5.56	123.30	118.30
1	A	117	ASP	CB-CG-OD2	5.47	123.22	118.30
1	A	46	ASP	CB-CG-OD2	5.40	123.16	118.30
1	A	177	ASP	CB-CG-OD2	5.35	123.11	118.30
1	B	479	ASP	CB-CG-OD2	5.26	123.04	118.30
1	B	252	ASP	CB-CG-OD2	5.24	123.02	118.30
1	A	366	ASP	CB-CG-OD2	5.14	122.93	118.30
1	A	446	ASP	CB-CG-OD2	5.10	122.89	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	458	SER	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3967	0	3757	26	0
1	B	3967	0	3757	23	1
2	A	28	0	20	1	0
2	B	28	0	20	0	0
3	A	450	0	0	14	1
3	B	389	0	0	9	0
All	All	8829	0	7554	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:458:SER:HA	3:B:2355:HOH:O	1.46	1.15
1:B:75:ASN:ND2	3:B:2063:HOH:O	2.25	0.69
1:A:184:LYS:NZ	3:A:2193:HOH:O	2.32	0.63
1:B:73:GLY:O	3:B:2061:HOH:O	2.16	0.62
1:A:75:ASN:ND2	3:A:2082:HOH:O	2.34	0.60
1:A:304:SER:O	1:A:307:ARG:HD3	2.04	0.56
1:A:52:ASN:HD21	1:A:144:ASP:HA	1.71	0.55
1:A:279:GLU:HG3	3:A:2265:HOH:O	2.06	0.55
1:B:90:MET:HE1	1:B:475:ILE:HD12	1.88	0.55
1:B:52:ASN:HD22	1:B:52:ASN:N	2.05	0.54
1:B:79:MET:O	1:B:82:THR:HG22	2.07	0.54
1:B:414:THR:HG23	1:B:417:THR:H	1.72	0.53
1:A:442:LYS:NZ	3:A:2410:HOH:O	2.41	0.53
1:A:123:ARG:HD3	3:A:2185:HOH:O	2.09	0.53
1:A:452:GLN:HG3	3:A:2418:HOH:O	2.09	0.52
1:A:452:GLN:CD	3:A:2418:HOH:O	2.49	0.51
1:B:52:ASN:HD21	1:B:144:ASP:HA	1.76	0.51
1:A:307:ARG:HG3	3:A:2285:HOH:O	2.11	0.50
1:B:457:TRP:O	1:B:458:SER:HB2	2.11	0.50
1:A:90:MET:HE1	1:A:492:ALA:HA	1.93	0.50
1:A:52:ASN:N	1:A:52:ASN:HD22	2.10	0.49
1:A:247:LYS:NZ	3:A:2230:HOH:O	2.45	0.49
1:A:419:LEU:HD13	1:A:424:ALA:HB2	1.93	0.49
1:B:279:GLU:HG3	3:B:2088:HOH:O	2.12	0.49
1:A:295:ARG:HD2	1:B:273:LEU:HD11	1.97	0.46
1:B:480:ARG:HD3	3:B:2061:HOH:O	2.15	0.46
1:A:414:THR:HG23	1:A:417:THR:H	1.81	0.46
1:A:111:GLU:HG2	1:A:112:GLY:N	2.30	0.46
1:A:101:TRP:HB3	1:A:102:PRO:HD3	1.98	0.45
1:A:224:LEU:HD13	1:A:350:VAL:O	2.17	0.45
1:A:502:LYS:N	3:A:2447:HOH:O	2.49	0.45
1:A:82:THR:HG22	3:A:2030:HOH:O	2.18	0.44
1:B:85:ARG:CZ	3:B:2075:HOH:O	2.65	0.44
1:B:320:LYS:HE2	3:B:2177:HOH:O	2.18	0.44
3:A:2282:HOH:O	1:B:307:ARG:CZ	2.64	0.44
1:A:90:MET:HE1	1:A:475:ILE:HD12	2.00	0.43
3:A:2286:HOH:O	1:B:307:ARG:HG3	2.17	0.43
1:A:376:ASN:HB2	1:A:377:PRO:CD	2.49	0.43
1:B:75:ASN:HD21	1:B:78:HIS:HD2	1.67	0.43
1:A:452:GLN:CG	3:A:2418:HOH:O	2.65	0.42
1:A:194:THR:HG21	2:A:1503:NTZ:H2	2.02	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:376:ASN:HB2	1:B:377:PRO:CD	2.49	0.41
1:A:232:HIS:CE1	1:A:313:PHE:CE2	3.08	0.41
1:B:101:TRP:N	1:B:102:PRO:HD2	2.36	0.41
1:B:75:ASN:ND2	3:B:2062:HOH:O	2.54	0.40
1:B:143:TRP:CZ3	1:B:205:PHE:CE2	3.09	0.40
1:B:429:LYS:HD3	3:B:2340:HOH:O	2.22	0.40
1:B:141:PHE:CZ	1:B:143:TRP:HA	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:79:MET:SD	3:A:2413:HOH:O 2_464	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	489/512 (96%)	470 (96%)	18 (4%)	1 (0%)	47	38
1	B	489/512 (96%)	471 (96%)	15 (3%)	3 (1%)	25	15
All	All	978/1024 (96%)	941 (96%)	33 (3%)	4 (0%)	34	24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	458	SER
1	A	458	SER
1	B	158	ASP
1	B	482	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	423/441 (96%)	402 (95%)	21 (5%)	24	15
1	B	423/441 (96%)	408 (96%)	15 (4%)	36	27
All	All	846/882 (96%)	810 (96%)	36 (4%)	29	19

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	MET
1	A	52	ASN
1	A	70	SER
1	A	75	ASN
1	A	82	THR
1	A	85	ARG
1	A	86	LEU
1	A	111	GLU
1	A	224	LEU
1	A	285	ASN
1	A	330	LEU
1	A	344	SER
1	A	346	ASN
1	A	348	SER
1	A	386	LEU
1	A	419	LEU
1	A	442	LYS
1	A	458	SER
1	A	459	LEU
1	A	462	ASN
1	A	482	ASN
1	B	52	ASN
1	B	86	LEU
1	B	111	GLU
1	B	224	LEU
1	B	252	ASP
1	B	285	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	316	GLU
1	B	328	LEU
1	B	330	LEU
1	B	348	SER
1	B	386	LEU
1	B	419	LEU
1	B	459	LEU
1	B	462	ASN
1	B	482	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	52	ASN
1	A	132	ASN
1	A	178	ASN
1	A	246	ASN
1	A	276	GLN
1	A	452	GLN
1	A	462	ASN
1	A	481	ASN
1	B	52	ASN
1	B	78	HIS
1	B	132	ASN
1	B	246	ASN
1	B	452	GLN
1	B	462	ASN
1	B	481	ASN
1	B	482	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NTZ	B	1503	-	13,15,15	2.63	4 (30%)	12,22,22	3.04	4 (33%)
2	NTZ	B	1502	-	13,15,15	2.41	4 (30%)	12,22,22	2.60	3 (25%)
2	NTZ	A	1502	-	13,15,15	2.52	4 (30%)	12,22,22	2.43	3 (25%)
2	NTZ	A	1503	-	13,15,15	2.54	4 (30%)	12,22,22	3.16	6 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NTZ	B	1503	-	-	1/2/22/22	0/1/2/2
2	NTZ	B	1502	-	-	0/2/22/22	0/1/2/2
2	NTZ	A	1502	-	-	0/2/22/22	0/1/2/2
2	NTZ	A	1503	-	-	1/2/22/22	0/1/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1502	NTZ	N21-N18	-6.35	1.25	1.34
2	B	1502	NTZ	N21-N18	-6.04	1.26	1.34
2	B	1503	NTZ	N21-N18	-5.91	1.26	1.34
2	A	1503	NTZ	N21-N18	-5.72	1.26	1.34
2	A	1503	NTZ	N17-N18	4.91	1.41	1.30
2	B	1503	NTZ	N17-N18	4.74	1.41	1.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1502	NTZ	C1-C2	-4.15	1.44	1.51
2	B	1503	NTZ	N17-N1	4.12	1.42	1.34
2	B	1502	NTZ	N17-N18	3.90	1.39	1.30
2	A	1503	NTZ	N17-N1	3.88	1.41	1.34
2	A	1502	NTZ	N17-N18	3.71	1.39	1.30
2	B	1503	NTZ	C1-C2	-3.64	1.45	1.51
2	B	1502	NTZ	C1-C2	-3.52	1.45	1.51
2	A	1503	NTZ	C1-C2	-3.13	1.46	1.51
2	B	1502	NTZ	N17-N1	3.05	1.40	1.34
2	A	1502	NTZ	N17-N1	2.85	1.40	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1503	NTZ	C1-N21-N18	9.49	113.99	105.28
2	B	1502	NTZ	C1-N21-N18	7.90	112.53	105.28
2	A	1503	NTZ	C1-N21-N18	7.66	112.31	105.28
2	A	1502	NTZ	C1-N21-N18	7.04	111.74	105.28
2	A	1503	NTZ	C3-C4-C5	-4.96	102.98	111.37
2	A	1502	NTZ	C3-C4-C5	-3.26	105.86	111.37
2	B	1502	NTZ	C3-C4-C5	-3.24	105.90	111.37
2	A	1503	NTZ	N21-N18-N17	-3.06	106.64	110.09
2	A	1503	NTZ	O2-C2-C3	2.97	114.80	108.55
2	A	1503	NTZ	O4-C4-C5	2.68	114.52	109.77
2	B	1503	NTZ	O3-C3-C2	-2.57	103.24	108.98
2	B	1503	NTZ	N21-N18-N17	-2.53	107.24	110.09
2	A	1502	NTZ	O3-C3-C2	-2.21	104.05	108.98
2	B	1502	NTZ	N21-N18-N17	-2.07	107.75	110.09
2	A	1503	NTZ	C4-C3-C2	-2.05	107.12	110.24
2	B	1503	NTZ	C4-C3-C2	2.04	113.33	110.24

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1503	NTZ	N1-C5-C6-O6
2	B	1503	NTZ	N1-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1503	NTZ	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	491/512 (95%)	-0.03	16 (3%) 46 49	16, 22, 34, 44	1 (0%)
1	B	491/512 (95%)	0.13	20 (4%) 37 40	16, 26, 37, 44	1 (0%)
All	All	982/1024 (95%)	0.05	36 (3%) 41 44	16, 24, 36, 44	2 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	346	ASN	5.5
1	B	502	LYS	4.5
1	B	501	ALA	4.3
1	A	501	ALA	3.9
1	B	415	LYS	3.5
1	A	502	LYS	3.4
1	A	346	ASN	3.4
1	B	457	TRP	3.4
1	B	416	GLU	3.3
1	A	457	TRP	3.3
1	A	458	SER	3.2
1	B	111	GLU	3.1
1	B	252	ASP	3.1
1	A	416	GLU	3.0
1	A	252	ASP	2.9
1	A	428	TYR	2.7
1	A	422	GLU	2.7
1	B	345	PRO	2.6
1	A	26	SER	2.6
1	A	347	TYR	2.6
1	A	415	LYS	2.6
1	B	428	TYR	2.5
1	B	458	SER	2.5
1	A	366	ASP	2.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	15	PRO	2.4
1	B	152	LYS	2.4
1	A	12	MET	2.4
1	B	33	ALA	2.3
1	B	12	MET	2.3
1	B	107	LYS	2.2
1	B	253	ASP	2.1
1	B	384	GLU	2.1
1	B	456	ALA	2.0
1	B	14	SER	2.0
1	A	456	ALA	2.0
1	B	459	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NTZ	B	1503	14/14	0.49	0.28	37,42,44,44	0
2	NTZ	A	1503	14/14	0.68	0.27	39,44,45,46	0
2	NTZ	B	1502	14/14	0.94	0.14	18,20,23,24	0
2	NTZ	A	1502	14/14	0.97	0.15	18,20,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.