



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 7, 2022 – 12:34 PM EDT

PDB ID : 3V4D
Title : Crystal structure of RutC protein a member of the YjgF family from E.coli
Authors : Knapik, A.A.; Petkowski, J.J.; Otwinowski, Z.; Cymborowski, M.T.; Cooper, D.R.; Chruszcz, M.; Porebski, P.J.; Niedzialkowska, E.; Almo, S.C.; Minor, W.; New York Structural Genomics Research Consortium (NYSGRC)
Deposited on : 2011-12-14
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.27
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	 11% 81% 8% • 10%
1	B	134	 11% 89% •• 7%
1	C	134	 7% 90% •• 6%
1	D	134	 10% 93% •••
1	E	134	 4% 93% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	134	 <p>A horizontal bar chart representing the quality of the chain. The bar is divided into three segments: a red segment on the left labeled '17%', a large green segment in the middle labeled '95%', and a small yellow/orange segment on the right. To the right of the bar, there are two small black dots.</p>

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 6007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminoacrylate peracid reductase RutC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	120	890	579	144	164	2	1	0	0	0
1	C	126	954	617	157	175	2	3	0	2	0
1	F	133	984	639	163	177	2	3	0	1	0
1	B	125	934	605	151	173	2	3	0	1	0
1	D	130	965	624	158	179	2	2	0	0	0
1	E	125	938	607	153	174	2	2	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	LEU	-	expression tag	UNP P0AFQ6
A	-4	TYR	-	expression tag	UNP P0AFQ6
A	-3	PHE	-	expression tag	UNP P0AFQ6
A	-2	GLN	-	expression tag	UNP P0AFQ6
A	-1	GLY	-	expression tag	UNP P0AFQ6
A	0	HIS	-	expression tag	UNP P0AFQ6
C	-5	LEU	-	expression tag	UNP P0AFQ6
C	-4	TYR	-	expression tag	UNP P0AFQ6
C	-3	PHE	-	expression tag	UNP P0AFQ6
C	-2	GLN	-	expression tag	UNP P0AFQ6
C	-1	GLY	-	expression tag	UNP P0AFQ6
C	0	HIS	-	expression tag	UNP P0AFQ6
F	-5	LEU	-	expression tag	UNP P0AFQ6
F	-4	TYR	-	expression tag	UNP P0AFQ6
F	-3	PHE	-	expression tag	UNP P0AFQ6
F	-2	GLN	-	expression tag	UNP P0AFQ6
F	-1	GLY	-	expression tag	UNP P0AFQ6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	HIS	-	expression tag	UNP P0AFQ6
B	-5	LEU	-	expression tag	UNP P0AFQ6
B	-4	TYR	-	expression tag	UNP P0AFQ6
B	-3	PHE	-	expression tag	UNP P0AFQ6
B	-2	GLN	-	expression tag	UNP P0AFQ6
B	-1	GLY	-	expression tag	UNP P0AFQ6
B	0	HIS	-	expression tag	UNP P0AFQ6
D	-5	LEU	-	expression tag	UNP P0AFQ6
D	-4	TYR	-	expression tag	UNP P0AFQ6
D	-3	PHE	-	expression tag	UNP P0AFQ6
D	-2	GLN	-	expression tag	UNP P0AFQ6
D	-1	GLY	-	expression tag	UNP P0AFQ6
D	0	HIS	-	expression tag	UNP P0AFQ6
E	-5	LEU	-	expression tag	UNP P0AFQ6
E	-4	TYR	-	expression tag	UNP P0AFQ6
E	-3	PHE	-	expression tag	UNP P0AFQ6
E	-2	GLN	-	expression tag	UNP P0AFQ6
E	-1	GLY	-	expression tag	UNP P0AFQ6
E	0	HIS	-	expression tag	UNP P0AFQ6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	44	Total O 44 44	0	0
2	C	62	Total O 62 62	0	0
2	F	47	Total O 47 47	0	0
2	B	70	Total O 70 70	0	0
2	D	45	Total O 45 45	0	0
2	E	74	Total O 74 74	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aminoacrylate peracid reductase RutC




- Molecule 1: Aminoacrylate peracid reductase RutC




- Molecule 1: Aminoacrylate peracid reductase RutC




- Molecule 1: Aminoacrylate peracid reductase RutC




- Molecule 1: Aminoacrylate peracid reductase RutC



- Molecule 1: Aminoacrylate peracid reductase RutC

Chain E:  4% 93% 7%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.25Å 179.18Å 45.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 33.58 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (50.00-1.95) 99.7 (33.58-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.193 , 0.217 0.199 , 0.222	Depositor DCC
R_{free} test set	2816 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtrriage
Anisotropy	0.713	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6007	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.39 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3766e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	1/910 (0.1%)	0.82	0/1243
1	B	0.91	0/956	0.86	1/1303 (0.1%)
1	C	0.95	0/980	0.85	1/1334 (0.1%)
1	D	0.85	0/987	0.81	0/1347
1	E	0.92	0/958	0.82	0/1303
1	F	0.85	0/1010	0.80	0/1379
All	All	0.90	1/5801 (0.0%)	0.83	2/7909 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	83	TRP	CD2-CE2	5.35	1.47	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	49	LYS	CD-CE-NZ	5.29	123.87	111.70
1	B	37	ASP	CB-CA-C	-5.20	100.00	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	890	0	866	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	934	0	913	6	0
1	C	954	0	944	4	0
1	D	965	0	943	7	0
1	E	938	0	922	0	0
1	F	984	0	958	5	0
2	A	44	0	0	2	0
2	B	70	0	0	2	0
2	C	62	0	0	1	0
2	D	45	0	0	1	0
2	E	74	0	0	0	0
2	F	47	0	0	1	0
All	All	6007	0	5546	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:16:ALA:HB1	1:F:17:PRO:HD2	1.57	0.86
1:A:47:ASP:OD1	1:A:50:ALA:HB3	1.78	0.84
1:D:90:ASN:HD21	1:D:104:ARG:HH22	1.32	0.77
1:F:16:ALA:HB1	1:F:17:PRO:CD	2.15	0.76
1:D:16:ALA:HB1	1:D:17:PRO:HD2	1.67	0.75
1:D:16:ALA:HB1	1:D:17:PRO:CD	2.22	0.70
1:A:5:VAL:HG11	1:A:19:VAL:HG11	1.74	0.68
1:B:37:ASP:HB3	1:B:39:HIS:H	1.61	0.66
1:C:1[A]:MSE:HA	1:C:1[A]:MSE:CE	2.29	0.63
1:D:90:ASN:ND2	1:D:104:ARG:HH12	1.99	0.59
1:C:16:ALA:HB2	2:C:252:HOH:O	2.03	0.57
1:A:5:VAL:CG1	1:A:19:VAL:HG11	2.36	0.56
1:C:1[A]:MSE:HA	1:C:1[A]:MSE:HE3	1.88	0.55
1:D:90:ASN:HD21	1:D:104:ARG:NH2	2.01	0.55
1:B:1[A]:MSE:HE1	2:B:338:HOH:O	2.07	0.55
1:A:44:PHE:HB3	1:A:47:ASP:OD1	2.08	0.53
1:B:1[A]:MSE:HE1	1:B:25:ASP:OD1	2.11	0.51
1:A:70:MSE:HE3	1:A:97:PHE:CZ	2.46	0.50
1:A:5:VAL:HG12	1:A:7:ILE:HG13	1.96	0.48
1:D:14:PRO:HB3	2:D:282:HOH:O	2.15	0.47
1:A:47:ASP:OD1	1:A:50:ALA:CB	2.58	0.46
1:F:16:ALA:CB	1:F:17:PRO:CD	2.87	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:-4:TYR:CB	1:F:1[B]:MSE:CE	2.95	0.45
1:B:1[B]:MSE:HG3	1:B:2:PRO:HD2	1.98	0.45
1:B:105:PHE:C	1:B:105:PHE:CD1	2.90	0.45
1:A:3:LYS:NZ	2:A:393:HOH:O	2.41	0.44
1:A:69:THR:HG22	1:D:65:THR:HA	1.99	0.44
1:C:5:VAL:O	1:C:6:ILE:HD13	2.19	0.43
1:F:46:ASP:CB	2:F:365:HOH:O	2.66	0.42
1:B:1[A]:MSE:HE3	2:B:229:HOH:O	2.19	0.41
1:A:98:PRO:HD2	2:A:135:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/134 (87%)	115 (99%)	1 (1%)	0	100	100
1	B	122/134 (91%)	121 (99%)	1 (1%)	0	100	100
1	C	124/134 (92%)	122 (98%)	2 (2%)	0	100	100
1	D	128/134 (96%)	126 (98%)	2 (2%)	0	100	100
1	E	121/134 (90%)	120 (99%)	1 (1%)	0	100	100
1	F	132/134 (98%)	126 (96%)	5 (4%)	1 (1%)	19	9
All	All	743/804 (92%)	730 (98%)	12 (2%)	1 (0%)	51	43

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	0	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/106 (85%)	89 (99%)	1 (1%)	73	71
1	B	95/106 (90%)	93 (98%)	2 (2%)	53	46
1	C	99/106 (93%)	95 (96%)	4 (4%)	31	19
1	D	98/106 (92%)	97 (99%)	1 (1%)	76	74
1	E	97/106 (92%)	97 (100%)	0	100	100
1	F	97/106 (92%)	95 (98%)	2 (2%)	53	46
All	All	576/636 (91%)	566 (98%)	10 (2%)	76	55

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ASP
1	C	1[A]	MSE
1	C	1[B]	MSE
1	C	53[A]	ARG
1	C	53[B]	ARG
1	F	1[A]	MSE
1	F	1[B]	MSE
1	B	1[A]	MSE
1	B	1[B]	MSE
1	D	90	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	90	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	119/134 (88%)	0.75	15 (12%) 3 6	23, 48, 86, 111	0
1	B	123/134 (91%)	0.35	15 (12%) 4 6	20, 38, 79, 96	0
1	C	124/134 (92%)	0.22	9 (7%) 15 23	19, 37, 59, 71	0
1	D	128/134 (95%)	0.69	14 (10%) 5 9	21, 44, 69, 86	0
1	E	123/134 (91%)	0.11	6 (4%) 29 39	20, 34, 55, 77	0
1	F	131/134 (97%)	0.85	23 (17%) 1 1	21, 44, 90, 109	6 (4%)
All	All	748/804 (93%)	0.50	82 (10%) 5 9	19, 40, 77, 111	6 (0%)

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	13	ALA	7.8
1	D	12	SER	6.5
1	D	0	HIS	6.2
1	F	13	ALA	5.8
1	A	7	ILE	5.8
1	D	-1	GLY	5.6
1	F	-3	PHE	5.6
1	F	15	LEU	5.4
1	F	16	ALA	5.3
1	F	-1	GLY	5.0
1	A	38	GLN	4.8
1	A	50	ALA	4.1
1	A	44	PHE	4.0
1	F	39	HIS	3.8
1	A	9	ALA	3.7
1	F	79	PHE	3.7
1	F	45	ALA	3.6
1	A	39	HIS	3.5
1	A	105	PHE	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	78	ILE	3.5
1	A	2	PRO	3.4
1	A	10	GLY	3.4
1	C	105	PHE	3.4
1	B	45	ALA	3.3
1	A	40	ASN	3.2
1	D	39	HIS	3.2
1	F	105	PHE	3.2
1	F	-4	TYR	3.2
1	B	9	ALA	3.1
1	E	0	HIS	3.1
1	F	-2	GLN	3.1
1	D	15	LEU	3.1
1	F	-5	LEU	3.1
1	B	0	HIS	3.0
1	E	105	PHE	3.0
1	F	44	PHE	2.9
1	D	45	ALA	2.9
1	C	78	ILE	2.9
1	F	107	ILE	2.9
1	D	128	LYS	2.9
1	F	42	VAL	2.8
1	D	14	PRO	2.8
1	D	121	ALA	2.8
1	A	41	ASN	2.7
1	B	120	ILE	2.7
1	B	75	PHE	2.7
1	F	37	ASP	2.6
1	E	77	SER	2.6
1	C	127	ALA	2.6
1	B	121	ALA	2.6
1	E	46	ASP	2.5
1	D	98	PRO	2.5
1	C	16	ALA	2.5
1	C	45	ALA	2.5
1	B	78	ILE	2.5
1	B	123	ILE	2.5
1	C	79	PHE	2.5
1	D	123	ILE	2.4
1	B	105	PHE	2.4
1	E	78	ILE	2.4
1	B	44	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	46	ASP	2.3
1	B	37	ASP	2.3
1	F	40	ASN	2.3
1	E	75	PHE	2.3
1	A	45	ALA	2.3
1	D	120	ILE	2.3
1	A	120	ILE	2.2
1	F	14	PRO	2.2
1	D	105	PHE	2.1
1	F	5	VAL	2.1
1	C	128	LYS	2.1
1	A	79	PHE	2.1
1	F	77	SER	2.1
1	B	77	SER	2.1
1	A	43	LEU	2.1
1	C	77	SER	2.0
1	F	38	GLN	2.0
1	C	121	ALA	2.0
1	F	0	HIS	2.0
1	B	10	GLY	2.0
1	B	107	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.