

Full wwPDB X-ray Structure Validation Report (i)

Aug 8, 2020 – 01:29 PM BST

PDB ID	:	5V4E
Title	:	Engineered human IgG Fc domain glyco801 (Fc801)
Authors	:	Yan, W.; Marshall, N.; Zhang, Y.J.
Deposited on	:	2017-03-09
Resolution	:	3.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries}, { m resolution\ range}({ m \AA}))$		
R _{free}	130704	1335 (3.24-3.20)		
Clashscore	141614	1460 (3.24-3.20)		
Ramachandran outliers	138981	1437 (3.24-3.20)		
Sidechain outliers	138945	1436 (3.24-3.20)		
RSRZ outliers	127900	1291 (3.24-3.20)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	226	86%	10% •
1	В	226	4% 75% 7	% 18%
1	С	226	% 60% 12%	28%
1	D	226	% 	8% 12%
1	Е	226	3% 82%	10% 8%
1	F	226	3% 64% 15%	22%



Conti	nued fron	ı previous	page									
Mol	Chain	Length		Quality of chain								
1	G	226		85%		12% •						
1	Н	226	%	87%		9% •						
2	Ι	8	13%	75%		13%						
2	М	8	25%	50%		25%						
3	J	3	33%		67%							
4	K	3		67%	339	⁄₀						
5	L	7	14%	71%		14%						
6	Ν	8	50%		38%	13%						

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	\mathbf{Res}	Chirality	Geometry	Clashes	Electron density
2	NAG	Ι	5	-	-	-	Х



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 11948 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	218	Total	С	Ν	Ο	S	0	0	0
	A	210	1590	1006	265	312	7	0	0	0
1	р	186	Total	С	Ν	Ο	S	0	0	0
	D 100	100	1281	812	217	247	5	0	0	0
1	C	169	Total	С	Ν	Ο	S	0	0	0
		102	1186	741	205	234	6	0	0	0
1	П	200	Total	С	Ν	Ο	S	0	0	0
L T			1431	907	237	282	5	0	0	0
1	F	20.8	Total	С	Ν	Ο	S	0	0	0
L T		200	1507	955	254	293	5	0	0	0
1	Б	177	Total	С	Ν	Ο	S	0	0	0
	Г	111	1252	786	210	251	5	0	0	0
1	C	210	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	G	219	1614	1024	264	318	8	0	0	0
1	II 016	216	Total	С	Ν	Ο	S	0	0	0
	11	210	1559	995	251	306	7		U	0

• Molecule 1 is a protein called Ig gamma-1 chain C region.

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	320	GLU	LYS	conflict	UNP P01857
А	386	ARG	GLN	conflict	UNP P01857
В	320	GLU	LYS	$\operatorname{conflict}$	UNP P01857
В	386	ARG	GLN	$\operatorname{conflict}$	UNP P01857
С	320	GLU	LYS	$\operatorname{conflict}$	UNP P01857
С	386	ARG	GLN	$\operatorname{conflict}$	UNP P01857
D	320	GLU	LYS	$\operatorname{conflict}$	UNP P01857
D	386	ARG	GLN	$\operatorname{conflict}$	UNP P01857
Е	320	GLU	LYS	$\operatorname{conflict}$	UNP P01857
E	386	ARG	GLN	$\operatorname{conflict}$	UNP P01857
F	320	GLU	LYS	conflict	UNP P01857
F	386	ARG	GLN	conflict	UNP P01857
G	320	GLU	LYS	$\operatorname{conflict}$	UNP P01857



0000000		ere ae page			
Chain	Residue	Modelled	Actual	Comment	Reference
G	386	ARG	GLN	$\operatorname{conflict}$	UNP P01857
Н	320	GLU	LYS	$\operatorname{conflict}$	UNP P01857
Н	386	ARG	GLN	$\operatorname{conflict}$	UNP P01857

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alp ha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-man nopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	Ι	8	Total 99	$\begin{array}{c} \mathrm{C} \\ 56 \end{array}$	N 4	O 39	0	0	0
2	М	8	Total 99	C 56	N 4	O 39	0	0	0

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alp ha-D-mannopyranose-(1-3)-beta-D-mannopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	J	3	Total 36	C 20	N 1	O 15	0	0	0

• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alp ha-D-mannopyranose-(1-6)-beta-D-mannopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	K	3	Total 36	C 20	N 1	O 15	0	0	0



• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alp ha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-de oxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	L	7	Total 85	C 48	N 3	О 34	0	0	0

• Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alp ha-D-mannopyranose-(1-2)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-man nopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
6	Ν	8	Total 99	C N 56 4	O 39	0	0	0

• Molecule 7 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	1	Total K 1 1	0	0
7	D	1	Total K 1 1	0	0

• Molecule 8 is alpha-D-mannopyranose (three-letter code: MAN) (formula: $C_6H_{12}O_6$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	Total C O 11 6 5	0	0
8	В	1	Total C O 11 6 5	0	0

• Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	В	1	Total 14	C 8	N 1	O 5	0	0



• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	7	Total O 7 7	0	0
10	В	3	Total O 3 3	0	0
10	С	3	Total O 3 3	0	0
10	D	4	Total O 4 4	0	0
10	Е	3	Total O 3 3	0	0
10	F	5	Total O 5 5	0	0
10	G	6	Total O 6 6	0	0
10	Н	5	Total O 5 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Ig gamma-1 chain C region





• Molecule 1: Ig gamma-1 chain C region



 $\label{eq:2} \bullet \mbox{Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)] 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-$

13%

75%

NAG1 NAG2 BMA3 BMA3 NAN4 NAN6 MAN6 NAG7 NAG7

 $\label{eq:2} \bullet \mbox{Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] - 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] - 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)] - 2-$

Chain M:	25%	50%	25%
NAG1 NAG2 BMA3 MAN4 NAG5 MAN6 FUG8			

Chain J:	33%	67%
BMA1 MAN2 NAG3		

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-bet a-D-mannopyranose

Chain K:	67%	33%

BMA1 MAN2 NAG3

• Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



 $\label{eq:main_optimal_states} \bullet \ Molecule \ 6: \ 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)] \\ beta-D-mannopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] \\ 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)] \\ 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)] \\ 2$

Chain N:	50%	38%	13%
AG 1 AG 2 AN 4 AN 4 AG 5 AG 6 AG 7 AG 7 AG 7 AG 7 AG 7 AG 7 AG 7 AG 7			



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	93.61Å 141.02Å 98.87Å	Deperitor
a, b, c, α , β , γ	90.00° 117.33° 90.00°	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	39.89 - 3.22	Depositor
Resolution (A)	39.88 - 3.22	EDS
% Data completeness	99.6 (39.89-3.22)	Depositor
(in resolution range)	99.6 (39.88-3.22)	EDS
R _{merge}	0.15	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.15 (at 3.18 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
D D .	0.236 , 0.281	Depositor
Π, Π_{free}	0.238 , 0.287	DCC
R_{free} test set	1944 reflections (5.26%)	wwPDB-VP
Wilson B-factor $(Å^2)$	83.4	Xtriage
Anisotropy	0.406	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31 , 60.8	EDS
L-test for twinning ²	$< L > = 0.48, < L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	11948	wwPDB-VP
Average B, all atoms $(Å^2)$	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, K, BMA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
10101	Cham	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.24	0/1638	0.44	0/2259
1	В	0.24	0/1314	0.44	0/1809
1	С	0.24	0/1215	0.45	0/1660
1	D	0.24	0/1470	0.43	0/2026
1	Е	0.24	0/1553	0.43	0/2142
1	F	0.25	0/1282	0.45	0/1758
1	G	0.25	0/1663	0.43	0/2294
1	Н	0.24	0/1607	0.42	0/2220
All	All	0.24	0/11742	0.44	0/16168

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1590	0	1375	17	0
1	В	1281	0	1027	9	0
1	С	1186	0	1022	15	0
1	D	1431	0	1205	12	0
1	Е	1507	0	1270	15	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1252	0	1041	19	0
1	G	1614	0	1418	19	0
1	Н	1559	0	1329	17	0
2	Ι	99	0	85	5	0
2	М	99	0	85	2	0
3	J	36	0	31	0	0
4	Κ	36	0	31	1	0
5	L	85	0	73	3	0
6	Ν	99	0	85	2	0
7	А	1	0	0	0	0
7	D	1	0	0	0	0
8	В	22	0	20	0	0
9	В	14	0	13	0	0
10	А	7	0	0	1	0
10	В	3	0	0	1	0
10	С	3	0	0	0	0
10	D	4	0	0	1	0
10	Ε	3	0	0	1	0
10	F	5	0	0	4	0
10	G	6	0	0	1	0
10	Н	5	0	0	4	0
All	All	11948	0	10110	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (118) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:249:ASP:OD1	10:G:601:HOH:O	1.97	0.81
1:C:346:PRO:HB3	1:C:372:PHE:HB3	1.69	0.74
1:A:301:ARG:HE	2:I:2:NAG:H81	1.52	0.74
1:B:389:ASN:ND2	10:B:601:HOH:O	2.22	0.71
1:E:367:CYS:O	10:E:601:HOH:O	2.08	0.71
1:E:346:PRO:HB3	1:E:372:PHE:HB3	1.73	0.70
1:A:354:SER:O	10:A:601:HOH:O	2.09	0.70
1:C:351:LEU:HB2	1:C:366:THR:HB	1.77	0.67
1:F:346:PRO:HB3	1:F:372:PHE:HB3	1.77	0.67
1:G:236:GLY:HA3	1:H:235:LEU:HB2	1.76	0.67
1:B:351:LEU:HB2	1:B:366:THR:HB	1.79	0.65
1:A:248:LYS:HE2	1:A:255:ARG:HE	1.62	0.65



		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:318:GLU:HA	1:D:337:SER:HB3	1.78	0.65
1:G:238:PRO:HG3	1:G:325:ASN:HD22	1.62	0.64
1:C:418:GLN:HA	1:C:443:LEU:HD22	1.80	0.63
1:A:265:ASP:OD1	2:I:1:NAG:N2	2.27	0.62
1:E:365:LEU:HD12	1:E:410:LEU:HD23	1.81	0.62
2:I:4:MAN:H61	2:I:5:NAG:H83	1.82	0.62
1:D:419:GLN:NE2	10:D:602:HOH:O	2.33	0.61
1:D:346:PRO:HB3	1:D:372:PHE:HB3	1.83	0.60
1:G:365:LEU:HD12	1:G:410:LEU:HD23	1.82	0.60
1:C:243:PHE:N	1:C:260:THR:O	2.34	0.60
1:C:399:ASP:OD2	1:D:409:LYS:NZ	2.34	0.60
1:F:351:LEU:HB2	1:F:366:THR:HB	1.84	0.59
1:G:226:CYS:N	1:H:226:CYS:SG	2.76	0.59
1:A:248:LYS:NZ	1:A:380:GLU:OE2	2.30	0.58
1:G:293:GLU:OE2	1:G:301:ARG:NH1	2.37	0.58
1:D:346:PRO:HD3	1:D:429:HIS:CD2	2.39	0.58
1:H:346:PRO:HB3	1:H:372:PHE:HB3	1.85	0.58
1:H:351:LEU:HB2	1:H:366:THR:HB	1.86	0.58
1:B:346:PRO:HB3	1:B:372:PHE:HB3	1.85	0.57
1:B:423:PHE:O	1:B:441:LEU:N	2.36	0.57
1:H:302:VAL:O	10:H:601:HOH:O	2.17	0.57
1:G:368:LEU:HD13	1:G:407:TYR:CZ	2.41	0.56
1:G:346:PRO:HB3	1:G:372:PHE:HB3	1.87	0.56
1:F:358:LEU:HD23	1:F:363:VAL:HG11	1.87	0.55
1:E:348:VAL:O	1:E:439:LYS:NZ	2.30	0.55
1:B:323:VAL:O	1:B:332:ILE:N	2.39	0.55
1:E:393:THR:HA	1:E:408:SER:HA	1.88	0.54
1:H:302:VAL:O	10:H:602:HOH:O	2.19	0.54
1:A:245:PRO:HD3	1:A:259:VAL:HG22	1.90	0.54
1:F:422:VAL:HA	1:F:442:SER:HB2	1.89	0.53
1:F:429:HIS:H	1:F:432:LEU:HD12	1.73	0.52
1:F:357:GLU:N	10:F:501:HOH:O	2.42	0.52
1:D:368:LEU:HD13	1:D:407:TYR:CZ	2.45	0.52
1:D:388:GLU:OE1	1:D:416:ARG:NH2	2.43	0.52
1:G:264:VAL:HG21	2:M:2:NAG:H61	1.92	0.52
1:D:339:ALA:HB3	1:D:374:PRO:HB3	1.91	0.52
1:F:249:ASP:OD1	1:F:255:ARG:HD2	2.10	0.51
1:H:263:VAL:N	10:H:602:HOH:O	2.43	0.51
1:H:368:LEU:HD13	1:H:407:TYR:CZ	2.46	0.51
1:C:370:LYS:HE2	1:D:409:LYS:HE3	1.92	0.51
1:F:443:LEU:O	10:F:502:HOH:O	2.19	0.50



		Interatomic	Clash
Atom-1	Atom-2	distance $(Å)$	overlap (Å)
1:E:368:LEU:HD13	1:E:407:TYR:CZ	2.46	0.50
1:H:240:VAL:HG22	1:H:263:VAL:HG22	1.93	0.50
1:B:424:SER:HA	1:B:440:SER:HA	1.93	0.49
1:F:358:LEU:HG	10:F:501:HOH:O	2.13	0.49
1:G:365:LEU:HB2	1:G:410:LEU:HB3	1.94	0.49
1:G:245:PRO:HD3	1:G:259:VAL:HG22	1.95	0.49
1:G:276:ASN:HB2	1:G:322:LYS:HB3	1.94	0.49
1:H:263:VAL:HB	10:H:601:HOH:O	2.12	0.49
1:C:368:LEU:HD13	1:C:407:TYR:CZ	2.47	0.49
1:G:271:PRO:HB2	1:G:292:ARG:HH11	1.79	0.48
1:H:301:ARG:HH11	6:N:2:NAG:H81	1.78	0.48
1:E:377:ILE:HG21	1:E:406:LEU:HD21	1.95	0.48
1:E:258:GLU:HB3	1:E:305:VAL:HG13	1.96	0.47
1:H:266:VAL:HG13	1:H:300:TYR:HB2	1.97	0.47
1:C:242:LEU:HA	1:C:261:CYS:HA	1.96	0.47
1:E:349:TYR:O	1:E:368:LEU:N	2.38	0.47
1:E:264:VAL:HG11	5:L:2:NAG:H83	1.95	0.47
1:C:248:LYS:HG3	1:C:428:MET:HE1	1.97	0.46
5:L:5:NAG:H3	5:L:5:NAG:H83	1.97	0.46
1:A:301:ARG:NE	2:I:2:NAG:H81	2.26	0.46
1:H:241:PHE:CZ	6:N:2:NAG:H61	2.50	0.46
1:F:403:SER:OG	1:F:404:PHE:N	2.48	0.46
1:H:339:ALA:HB3	1:H:374:PRO:HB3	1.98	0.46
1:A:368:LEU:HD13	1:A:407:TYR:CZ	2.51	0.46
1:F:424:SER:HA	1:F:440:SER:HA	1.96	0.46
1:A:341:GLY:HA3	1:A:373:TYR:CE2	2.51	0.46
1:C:389:ASN:N	1:C:389:ASN:OD1	2.45	0.46
1:A:418:GLN:HA	1:A:443:LEU:HD22	1.99	0.45
1:C:339:ALA:HB3	1:C:374:PRO:HB3	1.97	0.45
1:E:268:HIS:ND1	1:E:300:TYR:HE1	2.14	0.45
1:F:319:TYR:O	1:F:335:THR:HA	2.17	0.45
1:H:293:GLU:HG2	1:H:301:ARG:O	2.17	0.45
1:A:341:GLY:HA3	1:A:373:TYR:HE2	1.81	0.44
1:G:238:PRO:HD2	1:G:328:LEU:HD13	1.99	0.44
1:A:272:GLU:O	1:A:325:ASN:ND2	2.48	0.44
1:H:293:GLU:HG3	1:H:301:ARG:HB3	2.00	0.44
1:A:383:SER:HB3	1:A:388:GLU:OE2	2.17	0.44
1:D:351:LEU:HB2	1:D:366:THR:HB	1.98	0.44
1:A:258:GLU:HB3	1:A:305:VAL:HG13	2.00	0.43
1:F:368:LEU:HD13	1:F:407:TYR:CZ	2.53	0.43
1:G:242:LEU:HD23	1:G:242:LEU:HA	1.92	0.43



		Interatomic	Clash	
Atom-1	Atom-2	distance (\AA)	overlap (Å)	
1:G:378:ALA:HB3	1:G:428:MET:HB2	2.01	0.43	
1:F:423:PHE:O	1:F:441:LEU:N	2.50	0.43	
1:F:338:LYS:NZ	1:F:430:GLU:OE2	2.47	0.43	
1:A:344:ARG:HB2	1:A:373:TYR:N	2.34	0.43	
1:F:253:ILE:HA	1:F:310:HIS:CD2	2.54	0.43	
1:F:355:ARG:C	10:F:501:HOH:O	2.56	0.42	
1:D:433:HIS:NE2	1:G:387:PRO:HG3	2.34	0.42	
1:C:319:TYR:O	1:C:336:ILE:N	2.51	0.42	
1:E:351:LEU:HB2	1:E:366:THR:HB	2.01	0.42	
1:H:318:GLU:HA	1:H:337:SER:HB3	2.01	0.42	
1:F:389:ASN:HB2	1:G:293:GLU:HB2	2.02	0.42	
1:D:243:PHE:CE1	4:K:2:MAN:H2	2.54	0.42	
1:E:287:ALA:HB2	1:E:306:LEU:HD13	2.00	0.42	
1:C:247:PRO:O	1:C:251:LEU:HG	2.20	0.42	
1:C:367:CYS:HB2	1:C:381:TRP:CZ2	2.55	0.42	
1:E:264:VAL:CG1	5:L:2:NAG:H83	2.50	0.42	
1:A:260:THR:HG21	2:I:7:NAG:O6	2.20	0.42	
1:E:365:LEU:HB2	1:E:410:LEU:HB3	2.03	0.41	
1:C:390:ASN:O	1:C:410:LEU:HD12	2.21	0.41	
1:F:248:LYS:HG3	1:F:428:MET:HE1	2.03	0.41	
1:G:296:TYR:HD2	2:M:8:FUC:H61	1.86	0.41	
1:A:397:VAL:HG21	1:B:394:THR:HA	2.03	0.40	
1:B:289:THR:HG22	1:B:304:SER:HB2	2.03	0.40	
1:B:278:TYR:HB2	1:B:320:GLU:HB3	2.02	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowe		Outliers	Percentiles	
1	А	216/226~(96%)	213~(99%)	3 (1%)	0	100 10)0



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	178/226~(79%)	176~(99%)	2(1%)	0	100	100
1	С	154/226~(68%)	150~(97%)	4(3%)	0	100	100
1	D	192/226~(85%)	189~(98%)	3(2%)	0	100	100
1	Ε	206/226~(91%)	203~(98%)	3(2%)	0	100	100
1	F	169/226~(75%)	164~(97%)	5(3%)	0	100	100
1	G	217/226~(96%)	212~(98%)	5(2%)	0	100	100
1	Н	212/226~(94%)	$206 \ (97\%)$	6 (3%)	0	100	100
All	All	1544/1808~(85%)	1513 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	159/209~(76%)	159~(100%)	0	100 100
1	В	109/209~(52%)	109~(100%)	0	100 100
1	С	119/209~(57%)	119 (100%)	0	100 100
1	D	139/209~(66%)	139~(100%)	0	100 100
1	Ε	144/209~(69%)	144~(100%)	0	100 100
1	F	118/209~(56%)	117~(99%)	1 (1%)	81 92
1	G	168/209~(80%)	168~(100%)	0	100 100
1	Н	154/209~(74%)	$154 \ (100\%)$	0	100 100
All	All	1110/1672~(66%)	1109 (100%)	1 (0%)	93 98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	433	HIS



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

37 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tuno	Chain	Dog	Tink	Bo	ond leng	ths	В	ond ang	les
	туре	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	Ι	1	1,2	$14,\!14,\!15$	0.27	0	$17,\!19,\!21$	0.41	0
2	NAG	Ι	2	2	$14,\!14,\!15$	0.21	0	$17,\!19,\!21$	0.43	0
2	BMA	I	3	2	11,11,12	0.84	0	$15,\!15,\!17$	1.13	1(6%)
2	MAN	I	4	2	11,11,12	0.86	0	$15,\!15,\!17$	1.34	3 (20%)
2	NAG	Ι	5	2	14, 14, 15	0.25	0	$17,\!19,\!21$	0.54	0
2	MAN	Ι	6	2	11,11,12	0.62	0	$15,\!15,\!17$	1.15	2 (13%)
2	NAG	Ι	7	2	14, 14, 15	0.25	0	17,19,21	0.38	0
2	FUC	Ι	8	2	10, 10, 11	0.80	0	$14,\!14,\!16$	0.88	0
3	BMA	J	1	3	11,11,12	0.58	0	$15,\!15,\!17$	0.80	0
3	MAN	J	2	3	11, 11, 12	0.65	0	$15,\!15,\!17$	1.25	2(13%)
3	NAG	J	3	3	14,14,15	0.86	1 (7%)	$17,\!19,\!21$	0.87	1 (5%)
4	BMA	K	1	4	11,11,12	0.64	0	$15,\!15,\!17$	0.80	0
4	MAN	K	2	4	11,11,12	0.70	0	$15,\!15,\!17$	1.32	3 (20%)
4	NAG	K	3	4	14, 14, 15	0.27	0	$17,\!19,\!21$	0.49	0
5	NAG	L	1	1,5	14,14,15	0.36	0	$17,\!19,\!21$	0.75	1 (5%)
5	NAG	L	2	5	14, 14, 15	0.43	0	17,19,21	0.55	0
5	BMA	L	3	5	11,11,12	0.66	0	$15,\!15,\!17$	1.20	1 (6%)



Mal	Tuno	Chain	Dog	Tink	Bo	ond leng	$_{\rm ths}$	Bond angles		
	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	MAN	L	4	5	11,11,12	1.58	3 (27%)	15,15,17	1.54	3 (20%)
5	NAG	L	5	5	14,14,15	0.96	1 (7%)	17,19,21	1.78	3 (17%)
5	MAN	L	6	5	11,11,12	1.05	1 (9%)	15,15,17	2.22	2 (13%)
5	FUC	L	7	5	10,10,11	0.80	0	14,14,16	0.74	0
2	NAG	M	1	1,2	14,14,15	0.18	0	17,19,21	0.76	1 (5%)
2	NAG	М	2	2	14,14,15	0.92	1 (7%)	17,19,21	1.26	1 (5%)
2	BMA	М	3	2	11,11,12	0.72	0	$15,\!15,\!17$	1.41	3 (20%)
2	MAN	М	4	2	11,11,12	1.22	2 (18%)	$15,\!15,\!17$	1.48	2 (13%)
2	NAG	М	5	2	14,14,15	0.26	0	17,19,21	0.43	0
2	MAN	M	6	2	11,11,12	0.50	0	15,15,17	1.17	2 (13%)
2	NAG	М	7	2	14,14,15	0.29	0	17,19,21	0.45	0
2	FUC	М	8	2	10,10,11	0.79	0	14,14,16	1.07	1 (7%)
6	NAG	N	1	1,6	14,14,15	0.30	0	17,19,21	0.55	0
6	NAG	N	2	6	14,14,15	0.28	0	17,19,21	0.88	1 (5%)
6	BMA	N	3	6	11,11,12	1.97	2 (18%)	15,15,17	1.63	3 (20%)
6	MAN	N	4	6	11,11,12	0.94	1 (9%)	15,15,17	1.20	2 (13%)
6	NAG	N	5	6	14,14,15	0.38	0	17,19,21	0.61	0
6	MAN	N	6	6	11,11,12	0.77	0	$1\overline{5,15,17}$	0.83	1(6%)
6	NAG	N	7	6	14,14,15	0.22	0	17,19,21	0.44	0
6	FUC	N	8	6	10,10,11	0.57	0	14,14,16	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	Ι	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	Ι	2	2	-	3/6/23/26	0/1/1/1
2	BMA	Ι	3	2	-	2/2/19/22	0/1/1/1
2	MAN	Ι	4	2	-	2/2/19/22	0/1/1/1
2	NAG	Ι	5	2	-	2/6/23/26	0/1/1/1
2	MAN	Ι	6	2	-	0/2/19/22	0/1/1/1
2	NAG	Ι	7	2	-	2/6/23/26	0/1/1/1
2	FUC	Ι	8	2	-	-	0/1/1/1
3	BMA	J	1	3	-	0/2/19/22	0/1/1/1
3	MAN	J	2	3	-	0/2/19/22	0/1/1/1
3	NAG	J	3	3	-	2/6/23/26	0/1/1/1



Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
4	BMA	K	1	4	-	2/2/19/22	0/1/1/1
4	MAN	K	2	4	-	2/2/19/22	0/1/1/1
4	NAG	K	3	4	-	1/6/23/26	0/1/1/1
5	NAG	L	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	L	2	5	-	4/6/23/26	0/1/1/1
5	BMA	L	3	5	-	0/2/19/22	0/1/1/1
5	MAN	L	4	5	-	2/2/19/22	0/1/1/1
5	NAG	L	5	5	-	5/6/23/26	0/1/1/1
5	MAN	L	6	5	-	1/2/19/22	0/1/1/1
5	FUC	L	7	5	-	-	0/1/1/1
2	NAG	М	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	М	2	2	-	0/6/23/26	0/1/1/1
2	BMA	М	3	2	-	0/2/19/22	0/1/1/1
2	MAN	М	4	2	-	0/2/19/22	0/1/1/1
2	NAG	М	5	2	-	2/6/23/26	0/1/1/1
2	MAN	М	6	2	-	1/2/19/22	0/1/1/1
2	NAG	М	7	2	-	0/6/23/26	0/1/1/1
2	FUC	М	8	2	-	-	0/1/1/1
6	NAG	N	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	Ν	2	6	-	4/6/23/26	0/1/1/1
6	BMA	N	3	6	-	0/2/19/22	0/1/1/1
6	MAN	N	4	6	-	0/2/19/22	0/1/1/1
6	NAG	Ν	5	6	-	4/6/23/26	0/1/1/1
6	MAN	N	6	6	-	0/2/19/22	0/1/1/1
6	NAG	N	7	6	-	0/6/23/26	0/1/1/1
6	FUC	N	8	6	-	-	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	N	3	BMA	C1-C2	4.99	1.63	1.52
6	N	3	BMA	C2-C3	3.93	1.58	1.52
2	М	2	NAG	O5-C1	3.25	1.48	1.43
5	L	4	MAN	C2-C3	3.25	1.57	1.52
5	L	5	NAG	O5-C1	3.25	1.48	1.43
5	L	6	MAN	O5-C5	2.91	1.49	1.43
3	J	3	NAG	O5-C1	2.52	1.47	1.43
2	М	4	MAN	C1-C2	2.40	1.57	1.52
6	N	4	MAN	C1-C2	2.38	1.57	1.52
2	М	4	MAN	O2-C2	2.37	1.48	1.43



$\alpha \cdot \cdot \cdot \cdot$	e		
Continued	from	previous	page

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
5	L	4	MAN	C1-C2	2.28	1.57	1.52
5	L	4	MAN	C4-C3	2.06	1.57	1.52

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
5	L	6	MAN	C1-O5-C5	7.28	122.05	112.19
5	L	5	NAG	C1-O5-C5	5.12	119.13	112.19
2	М	2	NAG	C1-O5-C5	4.88	118.80	112.19
5	L	5	NAG	C2-N2-C7	4.32	129.06	122.90
2	М	4	MAN	O2-C2-C1	3.71	116.75	109.15
6	N	3	BMA	O3-C3-C2	3.66	117.00	109.99
3	J	2	MAN	C1-O5-C5	3.61	117.08	112.19
2	М	3	BMA	C1-C2-C3	3.44	113.89	109.67
2	Ι	4	MAN	O2-C2-C3	-3.27	103.59	110.14
3	J	3	NAG	C1-O5-C5	3.24	116.58	112.19
2	М	4	MAN	C1-O5-C5	3.23	116.57	112.19
4	К	2	MAN	C1-O5-C5	3.22	116.55	112.19
2	Ι	6	MAN	C1-O5-C5	3.20	116.53	112.19
2	М	6	MAN	C1-O5-C5	3.11	116.41	112.19
6	N	3	BMA	O2-C2-C1	3.06	115.42	109.15
5	L	4	MAN	C2-C3-C4	3.06	116.19	110.89
5	L	4	MAN	C1-C2-C3	3.00	113.35	109.67
6	Ν	4	MAN	C1-O5-C5	2.92	116.15	112.19
4	К	2	MAN	C1-C2-C3	-2.74	106.30	109.67
2	М	1	NAG	C1-O5-C5	2.61	115.73	112.19
6	Ν	4	MAN	O2-C2-C3	-2.59	104.95	110.14
3	J	2	MAN	O2-C2-C3	-2.52	105.09	110.14
6	N	3	BMA	C3-C4-C5	-2.51	105.77	110.24
2	Ι	4	MAN	C1-O5-C5	2.51	115.59	112.19
2	Ι	3	BMA	C1-C2-C3	2.50	112.73	109.67
5	L	1	NAG	C1-O5-C5	2.44	115.50	112.19
5	L	6	MAN	O5-C1-C2	2.41	114.50	110.77
2	М	3	BMA	O2-C2-C3	-2.37	105.40	110.14
2	М	6	MAN	O2-C2-C3	-2.31	105.50	110.14
5	L	4	MAN	O2-C2-C1	2.29	113.83	109.15
6	N	2	NAG	C1-O5-C5	2.27	115.27	112.19
2	Ι	4	MAN	C1-C2-C3	-2.27	106.88	109.67
5	L	3	BMA	C1-O5-C5	2.26	115.25	112.19
6	N	6	MAN	02-C2-C3	-2.20	105.73	110.14
4	K	2	MAN	O2-C2-C3	-2.15	105.82	110.14
2	М	3	BMA	C1-O5-C5	2.14	115.08	112.19



Mol	Chain	\mathbf{Res}	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	Ι	6	MAN	O2-C2-C3	-2.10	105.93	110.14
2	М	8	FUC	O5-C5-C4	2.08	113.25	109.52
5	L	5	NAG	C1-C2-N2	2.02	113.94	110.49

There are no chirality outliers.

All (45) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	N	5	NAG	C4-C5-C6-O6
5	L	2	NAG	O5-C5-C6-O6
6	N	5	NAG	O5-C5-C6-O6
2	М	5	NAG	O5-C5-C6-O6
5	L	2	NAG	C4-C5-C6-O6
2	Ι	1	NAG	O5-C5-C6-O6
2	Ι	7	NAG	O5-C5-C6-O6
6	N	2	NAG	O5-C5-C6-O6
2	М	5	NAG	C4-C5-C6-O6
5	L	5	NAG	C4-C5-C6-O6
5	L	4	MAN	O5-C5-C6-O6
2	Ι	1	NAG	C4-C5-C6-O6
6	N	2	NAG	C4-C5-C6-O6
2	Ι	7	NAG	C4-C5-C6-O6
6	N	2	NAG	C8-C7-N2-C2
6	N	2	NAG	O7-C7-N2-C2
5	L	2	NAG	C8-C7-N2-C2
5	L	2	NAG	O7-C7-N2-C2
3	J	3	NAG	C8-C7-N2-C2
3	J	3	NAG	O7-C7-N2-C2
2	Ι	2	NAG	C8-C7-N2-C2
2	Ι	2	NAG	O7-C7-N2-C2
5	L	5	NAG	C8-C7-N2-C2
5	L	5	NAG	O7-C7-N2-C2
4	K	2	MAN	O5-C5-C6-O6
2	Ι	4	MAN	C4-C5-C6-O6
2	Ι	3	BMA	O5-C5-C6-O6
2	Ι	5	NAG	O5-C5-C6-O6
5	L	5	NAG	O5-C5-C6-O6
4	K	3	NAG	O5-C5-C6-O6
2	Ι	3	BMA	C4-C5-C6-O6
2	I	4	MAN	O <u>5-C5-C6-</u> O6
4	K	2	MAN	C4-C5-C6-O6
4	K	1	BMA	C4-C5-C6-O6



Mol	Chain	Res	Type	Atoms
5	L	6	MAN	O5-C5-C6-O6
4	K	1	BMA	O5-C5-C6-O6
5	L	4	MAN	C4-C5-C6-O6
5	L	1	NAG	O5-C5-C6-O6
6	N	5	NAG	C1-C2-N2-C7
5	L	1	NAG	C4-C5-C6-O6
2	Ι	2	NAG	O5-C5-C6-O6
2	М	6	MAN	O5-C5-C6-O6
2	Ι	5	NAG	C3-C2-N2-C7
6	N	5	NAG	C3-C2-N2-C7
5	L	5	NAG	C3-C2-N2-C7

Continued from previous page...

There are no ring outliers.

11 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	М	2	NAG	1	0
2	Ι	1	NAG	1	0
6	N	2	NAG	2	0
5	L	2	NAG	2	0
4	Κ	2	MAN	1	0
2	Ι	5	NAG	1	0
2	М	8	FUC	1	0
2	Ι	4	MAN	1	0
2	Ι	2	NAG	2	0
2	Ι	7	NAG	1	0
5	L	5	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





























5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type Chain Res	Chain	Dog	Timl	Bond lengths			Bond angles		
		LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2		
8	MAN	В	502	-	11,11,12	0.95	0	15,15,17	1.19	2 (13%)
8	MAN	В	501	-	11,11,12	0.67	0	15,15,17	1.06	2 (13%)



Mol Typ	Tuno	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Dog	Link	Bo	ond leng	\mathbf{ths}	B	ond ang	les
	туре		nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2						
9	NAG	В	503	-	14,14,15	0.22	0	17,19,21	0.41	0						

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	MAN	В	502	-	-	1/2/19/22	0/1/1/1
8	MAN	В	501	-	-	0/2/19/22	0/1/1/1
9	NAG	В	503	-	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
8	В	501	MAN	C1-O5-C5	2.40	115.44	112.19
8	В	502	MAN	O2-C2-C3	-2.38	105.36	110.14
8	В	501	MAN	O2-C2-C3	-2.23	105.67	110.14
8	В	502	MAN	C1-O5-C5	2.00	114.91	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	Atoms
8	В	502	MAN	O5-C5-C6-O6
9	В	503	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2			$\mathbf{OWAB}(\mathrm{\AA}^2)$	$Q{<}0.9$
1	А	218/226~(96%)	-0.05	1 (0%)	91	86	$39,\ 70,\ 92,\ 103$	0
1	В	186/226~(82%)	0.24	9 (4%)	30	19	45, 86, 114, 127	0
1	С	162/226~(71%)	-0.09	3 (1%)	66	54	38, 66, 124, 136	0
1	D	200/226~(88%)	0.02	2(1%)	82	73	40, 77, 110, 120	0
1	Е	208/226~(92%)	-0.08	6 (2%)	51	37	39, 80, 108, 126	0
1	F	177/226~(78%)	0.10	6 (3%)	45	31	45, 75, 125, 136	0
1	G	219/226~(96%)	-0.21	1 (0%)	91	86	41,66,86,110	0
1	Н	216/226~(95%)	0.02	3 (1%)	75	64	42, 79, 101, 125	0
All	All	1586/1808~(87%)	-0.01	31 (1%)	65	52	38, 74, 112, 136	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	239	SER	6.2
1	В	276	ASN	4.6
1	F	263	VAL	4.1
1	С	321	CYS	3.9
1	А	324	SER	3.8
1	F	239	SER	3.6
1	G	361	ASN	3.2
1	С	275	PHE	3.1
1	Н	278	TYR	3.0
1	Е	324	SER	3.0
1	Е	268	HIS	2.9
1	Н	275	PHE	2.9
1	Е	275	PHE	2.8
1	В	273	VAL	2.8
1	В	316	GLY	2.6
1	В	323	VAL	2.6



Mol	Chain	Res	Type	RSRZ
1	В	293	GLU	2.6
1	Е	267	SER	2.5
1	В	261	CYS	2.5
1	D	419	GLN	2.5
1	В	277	TRP	2.4
1	В	287	ALA	2.4
1	С	276	ASN	2.4
1	F	240	VAL	2.3
1	В	289	THR	2.3
1	F	262	VAL	2.3
1	Н	235	LEU	2.3
1	F	313	TRP	2.1
1	F	264	VAL	2.1
1	Е	271	PRO	2.0
1	Е	270	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
3	BMA	J	1	11/12	0.62	0.22	$103,\!120,\!122,\!130$	0
3	NAG	J	3	14/15	0.70	0.35	$107,\!116,\!120,\!123$	14
2	NAG	Ι	5	14/15	0.78	0.40	$94,\!101,\!107,\!109$	14
5	MAN	L	6	11/12	0.81	0.19	$99,\!111,\!113,\!113$	0
5	NAG	L	1	14/15	0.82	0.42	$100,\!103,\!113,\!113$	14
2	NAG	М	5	14/15	0.82	0.29	101,104,111,111	14
2	MAN	Ι	4	11/12	0.83	0.14	$76,\!97,\!103,\!105$	0
6	MAN	N	4	11/12	0.85	0.30	84,87,92,95	11
3	MAN	J	2	11/12	0.85	0.27	114,119,122,122	0
4	BMA	K	1	11/12	0.86	0.17	$97,\!106,\!116,\!121$	0
6	NAG	N	2	14/15	0.86	0.25	83,86,92,94	0
5	NAG	L	5	14/15	0.86	0.22	$83,\!96,\!99,\!102$	0
4	NAG	K	3	14/15	0.87	0.29	$102,\!10\overline{5},\!116,\!118$	0



5V4E	
------	--

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	FUC	Ι	8	10/11	0.87	0.26	$93,\!99,\!102,\!104$	0
6	NAG	N	7	14/15	0.88	0.25	$73,\!87,\!98,\!98$	0
6	NAG	N	5	14/15	0.89	0.20	$83,\!93,\!98,\!101$	0
2	MAN	Ι	6	11/12	0.89	0.18	78,83,87,89	0
5	MAN	L	4	11/12	0.89	0.21	$97,\!100,\!109,\!114$	0
2	NAG	Ι	1	14/15	0.89	0.16	$80,\!87,\!91,\!95$	0
5	FUC	L	7	10/11	0.89	0.16	$103,\!108,\!110,\!111$	0
4	MAN	Κ	2	11/12	0.90	0.26	$96,\!110,\!114,\!117$	0
2	NAG	Ι	7	14/15	0.90	0.22	$67,\!83,\!90,\!92$	0
6	FUC	N	8	10/11	0.90	0.18	$94,\!101,\!105,\!109$	0
2	BMA	Ι	3	11/12	0.91	0.12	$79,\!82,\!90,\!99$	0
6	BMA	N	3	11/12	0.91	0.13	82,84,89,94	0
2	NAG	М	7	14/15	0.91	0.20	$51,\!65,\!73,\!78$	0
5	BMA	L	3	11/12	0.91	0.12	$98,\!108,\!112,\!112$	0
2	MAN	М	4	11/12	0.91	0.19	$88,\!93,\!96,\!104$	0
2	FUC	М	8	10/11	0.92	0.17	$76,\!81,\!85,\!89$	0
6	NAG	N	1	14/15	0.92	0.24	82,89,99,104	0
5	NAG	L	2	14/15	0.92	0.24	94,104,111,118	0
2	NAG	Ι	2	14/15	0.92	0.13	74,85,88,89	0
2	NAG	М	1	14/15	0.94	0.14	66,73,87,88	0
2	BMA	М	3	11/12	0.94	0.15	$58,\!66,\!76,\!85$	0
2	MAN	М	6	11/12	0.96	0.13	$56,\!60,\!68,\!69$	0
6	MAN	N	6	11/12	0.96	0.19	$84,\!87,\!94,\!102$	0
2	NAG	М	2	14/15	0.96	0.16	59,70,74,76	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	MAN	В	502	11/12	0.66	0.30	107,113,123,127	0
9	NAG	В	503	14/15	0.83	0.20	102,109,113,115	0
8	MAN	В	501	11/12	0.86	0.16	85,103,112,114	0
7	K	А	509	1/1	0.92	0.19	96,96,96,96	0
7	K	D	504	1/1	0.94	0.23	77,77,77,77	0



6.5 Other polymers (i)

There are no such residues in this entry.

