

Full wwPDB X-ray Structure Validation Report (i)

Oct 4, 2023 – 09:14 PM EDT

PDB ID	:	6VCI
Title	:	Lipophilic envelope-spanning tunnel protein (LetB), domains MCE2-MCE3
Authors	:	Isom, G.L.; Coudray, N.; MacRae, M.R.; McManus, C.T.; Ekiert, D.C.;
		Bhabha, G.
Deposited on	:	2019-12-21
Resolution	:	2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 2.15 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3311 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	1 C	214	Total	С	Ν	0	S	0	0	0
	214	1619	1027	272	317	3	0	0	0	
1	1 A	A 215	Total	С	Ν	0	S	0	2	0
	210	1644	1040	276	325	3	0	2	0	

• Molecule 1 is a protein called Lipophilic envelope-spanning tunnel protein LetB.

Chain	Residue	Modelled	Actual	Comment	Reference
С	158	MET	-	expression tag	UNP P76272
С	384	GLY	-	expression tag	UNP P76272
С	385	GLU	-	expression tag	UNP P76272
С	386	THR	-	expression tag	UNP P76272
С	387	HIS	-	expression tag	UNP P76272
С	388	HIS	-	expression tag	UNP P76272
С	389	HIS	-	expression tag	UNP P76272
С	390	HIS	-	expression tag	UNP P76272
С	391	HIS	-	expression tag	UNP P76272
С	392	HIS	-	expression tag	UNP P76272
А	158	MET	-	expression tag	UNP P76272
А	384	GLY	-	expression tag	UNP P76272
А	385	GLU	-	expression tag	UNP P76272
А	386	THR	-	expression tag	UNP P76272
А	387	HIS	-	expression tag	UNP P76272
А	388	HIS	-	expression tag	UNP P76272
А	389	HIS	-	expression tag	UNP P76272
А	390	HIS	-	expression tag	UNP P76272
А	391	HIS	-	expression tag	UNP P76272
А	392	HIS	-	expression tag	UNP P76272

There are 20 discrepancies between the modelled and reference sequences:

• Molecule 2 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	4	Total O 4 4	0	0
2	А	44	Total O 44 44	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 65	Depositor	
Cell constants	87.56Å 87.56Å 116.99Å	Depositor	
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor	
Resolution (Å)	41.00 - 2.15	Depositor	
% Data completeness	99.8 (41.00-2.15)	Depositor	
(in resolution range)		-	
R_{merge}	(Not available)	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$1.30 (at 2.16 \text{\AA})$	Xtriage	
Refinement program	PHENIX 1.10.1_2155	Depositor	
R, R_{free}	0.214 , 0.244	Depositor	
Wilson B-factor $(Å^2)$	57.6	Xtriage	
Anisotropy	0.117	Xtriage	
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.33$	Xtriage	
Estimated twinning fraction	0.047 for h,-h-k,-l	Xtriage	
Total number of atoms	3311	wwPDB-VP	
Average B, all atoms $(Å^2)$	83.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

