

wwPDB X-ray Structure Validation Summary Report (i)

Oct 5, 2023 – 12:21 AM EDT

PDB ID : 6VDW

Title: Crystal Structure of Dehaloperoxidase B in Complex with cofactor Man-

ganese(III) Porphyrin and Substrate 4-nitrophenol

Authors: Ghiladi, R.A.; de Serrano, V.S.; McGuire, A.; Malewschik, T.

Deposited on : 2019-12-27

Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)


There are 7 unique types of molecules in this entry. The entry contains 2900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dehaloperoxidase B.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	137	Total 1241			O 247	S 7	0	25	0
1	В	137	Total 1256			O 254	S 7	0	28	0


• Molecule 2 is PROTOPORPHYRIN IX CONTAINING MN (three-letter code: MNR) (formula: $C_{34}H_{32}MnN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Λ	1	Total	С	Mn	N	О	0	0
	A	1	43	34	1	4	4	0	0
9	2 B	1	Total	С	Mn	N	О	0	0
2		1	43	34	1	4	4	U	


• Molecule 3 is P-NITROPHENOL (three-letter code: NPO) (formula: $C_6H_5NO_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 10 6 1 3	0	0
3	В	1	Total C N O 10 6 1 3	0	0

 \bullet Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	В	1	Total C O 4 2 2	0	0
4	В	1	Total C O 4 2 2	0	0

• Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	A	1	Total C O 7 4 3	0	0
5	В	1	Total C O 7 4 3	0	0

 \bullet Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	Total O S 5 4 1	0	0
6	В	1	Total O S 5 4 1	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	114	Total O 114 114	0	0
7	В	117	Total O 117 117	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	59.30Å 67.77Å 67.61Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.58 - 1.50	Depositor
% Data completeness	99.8 (44.58-1.50)	Depositor
(in resolution range)	` , ,	
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.35 (at 1.50Å)	Xtriage
Refinement program	PHENIX dev_3714	Depositor
R, R_{free}	0.164 , 0.199	Depositor
Wilson B-factor (\mathring{A}^2)	15.6	Xtriage
Anisotropy	0.192	Xtriage
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	0.079 for -h,l,k	Xtriage
Total number of atoms	2900	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.97% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	gths	В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	EDO	A	207	-	3,3,3	0.83	0	2,2,2	0.18	0
4	EDO	В	203	_	3,3,3	0.83	0	2,2,2	0.50	0
2	MNR	В	201	1	36,50,50	4.67	23 (63%)	34,82,82	3.54	16 (47%)
4	EDO	A	203	-	3,3,3	0.60	0	2,2,2	0.57	0
4	EDO	В	204	-	3,3,3	0.47	0	2,2,2	0.46	0
4	EDO	A	208	-	3,3,3	0.31	0	2,2,2	0.55	0
4	EDO	A	205	-	3,3,3	0.73	0	2,2,2	0.57	0
6	SO4	В	207	-	4,4,4	0.51	0	6,6,6	2.54	3 (50%)
5	PEG	A	206	-	6,6,6	0.50	0	5,5,5	0.88	0
5	PEG	A	209	-	6,6,6	0.58	0	5,5,5	0.76	0
4	EDO	A	204	-	3,3,3	0.63	0	2,2,2	0.70	0
3	NPO	В	202	-	9,10,10	1.25	2 (22%)	11,13,13	1.52	0
3	NPO	A	202	-	9,10,10	1.49	0	11,13,13	2.63	5 (45%)
6	SO4	В	206	-	4,4,4	0.47	0	6,6,6	2.45	3 (50%)
5	PEG	A	210	-	6,6,6	0.37	0	5,5,5	0.55	0
5	PEG	В	205	-	6,6,6	0.45	0	5,5,5	0.84	0
2	MNR	A	201	1	36,50,50	4.81	28 (77%)	34,82,82	3.78	16 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	207	-	-	0/1/1/1	-
4	EDO	В	203	-	-	0/1/1/1	-
2	MNR	В	201	1	-	4/12/94/94	-
4	EDO	A	203	-	-	0/1/1/1	-
4	EDO	В	204	-	-	0/1/1/1	-
4	EDO	A	208	-	-	0/1/1/1	-
4	EDO	A	205	_	-	0/1/1/1	-
5	PEG	A	206	-	-	3/4/4/4	-
5	PEG	A	209	-	-	2/4/4/4	-
4	EDO	A	204	-	-	0/1/1/1	-
3	NPO	В	202	-	-	0/2/4/4	0/1/1/1
3	NPO	A	202	-	-	0/2/4/4	0/1/1/1
5	PEG	A	210	_	_	4/4/4/4	_

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	В	205	-	-	1/4/4/4	-
2	MNR	A	201	1	-	4/12/94/94	-

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\mathring{\mathrm{A}})$	$Ideal(\AA)$
2	A	201	MNR	C3B-CAB	-10.89	1.25	1.47
2	В	201	MNR	C3B-CAB	-10.54	1.26	1.47
2	A	201	MNR	CMD-C2D	8.93	1.72	1.51
2	A	201	MNR	CAD-C3D	8.56	1.67	1.52
2	В	201	MNR	C3C-C2C	8.45	1.54	1.37

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	201	MNR	C4A-C3A-C2A	-9.67	100.57	108.61
2	В	201	MNR	CHB-C4A-NA	8.31	128.19	120.84
2	В	201	MNR	CHA-C1A-NA	-8.03	113.73	120.84
2	A	201	MNR	CMA-C3A-C4A	7.84	139.53	124.73
2	В	201	MNR	C4A-C3A-C2A	-7.30	102.54	108.61

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	MNR	C2D-C3D-CAD-CBD
2	A	201	MNR	C4D-C3D-CAD-CBD
5	A	210	PEG	O2-C3-C4-O4
5	A	210	PEG	C1-C2-O2-C3
5	A	206	PEG	O1-C1-C2-O2


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the




average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

