

# Full wwPDB X-ray Structure Validation Report (i)

#### Nov 5, 2023 – 10:19 pm GMT

PDB ID	:	2VE6
Title	:	Crystal structure of a Murine MHC class I H2-Db molecule in complex with a
		photocleavable peptide
Authors	:	Grotenbreg, G.M.; Roan, N.R.; Guillen, E.; Meijers, R.; Wang, J.H.; Bell,
		G.W.; Starnbach, M.N.; Ploegh, H.L.
Deposited on	:	2007-10-17
Resolution	:	2.65  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374(2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	277	.% <b>8</b> 6%	14%
1	D	277	87%	13%
1	G	277	7%	16% ·
1	J	277	4% 87%	12%
2	В	99	89%	11%



Mol	Chain	Length	Quality of chain					
2	Е	99	.% <b>88%</b>		12%			
2	Н	99	8%		19% ••			
2	Κ	99	5%		16%			
3	С	9	78%		22%			
3	F	9	67%	22%	11%			
3	Ι	9	67%	22%	11%			
3	L	9	56%	33%	11%			



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 12856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN D-B AL-PHA CHAIN.

Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	Trace
1	Λ	276	Total	С	Ν	0	S	0	2	0
	Л	210	2291	1446	407	429	9	0	5	0
1	Л	276	Total	С	Ν	0	S	0	2	0
	D	270	2284	1441	406	428	9	0		0
1	С	276	Total	С	Ν	0	S	0	0	0
	G		2264	1430	400	425	9	0	0	0
1	1 T	276	Total	С	Ν	0	S	0	0	0
1	J		2264	1430	400	425	9	0	0	0

• Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues		At	$\mathbf{oms}$			ZeroOcc	AltConf	Trace
2	В	00	Total	С	Ν	0	S	0	0	0
	D	99	818	522	138	150	8	0		
0	F	00	Total	С	Ν	0	S	0	1	0
	Ľ	99	826	527	139	151	9	0	1	0
0	Ц	99	Total	С	Ν	0	S	0	0	0
	11		818	522	138	150	8	0	0	0
2 K	K	99	Total	С	Ν	0	S	0	0	0
	ſ		818	522	138	150	8		U	U

• Molecule 3 is a protein called SENDAI VIRUS EPITOPE RESIDUES 324-332 MODIFIED AT P7.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	С	9	Total C N O 75 50 11 14	0	0	0
3	F	9	Total   C   N   O	0	0	0
5	Ľ	3	75 50 11 14	0	0	0
3	T	9	Total C N O	0	0	0
Э	1	9	$75  ext{ } 50  ext{ } 11  ext{ } 14$			0



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
3	L	9	Total 75	C 50	N 11	0 14	0	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	49	Total         O           49         49	0	0
4	В	27	TotalO2727	0	0
4	С	1	Total O 1 1	0	0
4	D	38	Total         O           38         38	0	0
4	Е	11	Total         O           11         11	0	0
4	F	1	Total O 1 1	0	0
4	G	21	TotalO2121	0	0
4	Н	4	Total O 4 4	0	0
4	Ι	1	Total O 1 1	0	0
4	J	15	Total         O           15         15	0	0
4	К	5	Total O 5 5	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: H-2 CLASS I HISTOCOMPATIBILITY ANTIGEN D-B ALPHA CHAIN





• Molecule 2: BETA-2-MICROGLOBULIN

Chain B:	89%	11%	
M1 R12 122 128 128 128 128	K45 K43 K60 F70 F98 F98 F98		
• Molecule 2:	BETA-2-MICROGLOBULIN		
Chain E:	88%	12%	
M1 14 17 12 128 135	L40 K45 F163 F164 K93 K99		
• Molecule 2:	BETA-2-MICROGLOBULIN		
Chain H:	78%	19% ••	
M1 Q2 P5 R12 N21 N21 122	V27 128 135 135 135 135 140 140 144 140 144 144 144 144 144 144	66X	
• Molecule 2:	BETA-2-MICROGLOBULIN		
Chain K:	84%	16%	
M1 17 08 09 018 018 018 018 018 0123	N24 V27 F30 F30 F36 F36 F36 F36 F47 F70 F70 F70 F70 F70 F70 F70 F70 F70 F7		
• Molecule 3:	SENDAI VIRUS EPITOPE RESIDUES 324-3	32 MODIFIED	AT P7
Chain C:	78%	22%	
F1 P807 L9			
• Molecule 3:	SENDAI VIRUS EPITOPE RESIDUES 324-3	332 MODIFIED	AT P7
Chain F:	67% 22	2% 11%	
F1 Y6 A8 L9			
• Molecule 3:	SENDAI VIRUS EPITOPE RESIDUES 324-3	32 MODIFIED	AT P7
Chain I:	67% 229	% 11%	





• Molecule 3: SENDAI VIRUS EPITOPE RESIDUES 324-332 MODIFIED AT P7

Chain L: 56% 33% 11%





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	52.24Å 103.87Å 168.81Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.83^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	19.66 - 2.65	Depositor
Resolution (A)	19.66 - 2.65	EDS
% Data completeness	96.9 (19.66-2.65)	Depositor
(in resolution range)	96.9(19.66-2.65)	EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.56 (at 2.67 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
P. P.	0.234 , $0.291$	Depositor
$n, n_{free}$	0.234 , $0.286$	DCC
$R_{free}$ test set	2558 reflections $(5.06%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	51.6	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.30, 43.8	EDS
L-test for twinning <sup>2</sup>	$< L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	0.043 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12856	wwPDB-VP
Average B, all atoms $(Å^2)$	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PRQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond lengths		Bond angles	
MOI	Ullaill	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.30	0/2358	0.48	0/3200
1	D	0.30	0/2351	0.49	0/3192
1	G	0.29	0/2331	0.46	0/3166
1	J	0.28	0/2331	0.46	0/3166
2	В	0.28	0/844	0.47	0/1143
2	Е	0.30	0/852	0.48	0/1153
2	Н	0.30	0/844	0.61	2/1143~(0.2%)
2	Κ	0.27	0/844	0.43	0/1143
3	С	0.41	0/62	0.43	0/81
3	F	0.40	0/62	0.39	0/81
3	Ι	0.44	0/62	0.32	0/81
3	L	0.41	0/62	0.34	0/81
All	All	0.29	0/13003	0.48	2/17630~(0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	Н	0	1
3	С	0	1
3	F	0	2
3	Ι	0	4
3	L	0	4
All	All	0	12

There are no bond length outliers.

All (2) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Н	46	ILE	C-N-CD	-9.04	100.72	120.60
2	Н	46	ILE	C-N-CA	6.64	149.91	122.00

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	С	7	PRQ	Mainchain
3	F	6	TYR	Mainchain
3	F	7	PRQ	Mainchain
2	Н	46	ILE	Peptide
3	Ι	6	TYR	Mainchain,Peptide
3	Ι	7	PRQ	Mainchain,Peptide
3	L	6	TYR	Mainchain,Peptide
3	L	7	PRQ	Mainchain,Peptide

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2291	0	2162	27	0
1	D	2284	0	2155	26	0
1	G	2264	0	2136	43	0
1	J	2264	0	2136	21	0
2	В	818	0	795	7	0
2	Ε	826	0	803	7	0
2	Н	818	0	795	19	0
2	Κ	818	0	795	8	0
3	С	75	0	60	2	0
3	F	75	0	60	2	0
3	Ι	75	0	60	0	0
3	L	75	0	60	2	0
4	А	49	0	0	0	0
4	В	27	0	0	1	0
4	С	1	0	0	0	0
4	D	38	0	0	0	0
4	Е	11	0	0	0	0
4	F	1	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	21	0	0	1	0
4	Н	4	0	0	0	0
4	Ι	1	0	0	0	0
4	J	15	0	0	0	0
4	Κ	5	0	0	0	0
All	All	12856	0	12017	152	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (152) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:228:MET:HE2	1:D:245:ALA:HB1	1.43	1.00
2:H:46:ILE:HG22	2:H:47:PRO:C	1.87	0.92
1:D:215:LEU:HD22	1:D:261:VAL:HG22	1.56	0.87
1:G:187:ALA:HB3	1:G:272:LEU:HD11	1.64	0.79
1:D:228:MET:CE	1:D:245:ALA:HB1	2.14	0.77
2:H:46:ILE:HG22	2:H:47:PRO:O	1.91	0.71
1:G:199:VAL:HG23	1:G:251:LEU:HD12	1.71	0.70
2:H:46:ILE:CG2	2:H:47:PRO:C	2.60	0.69
1:G:201:LEU:HD12	1:G:249:VAL:HG21	1.75	0.69
1:D:187:ALA:HB3	1:D:272:LEU:HD21	1.75	0.67
1:A:25:VAL:HG21	4:B:2016:HOH:O	1.93	0.67
1:G:191:HIS:CE1	1:G:199:VAL:HG13	2.29	0.67
1:G:187:ALA:CB	1:G:272:LEU:HD11	2.24	0.67
1:D:77:SER:HB3	3:F:9:LEU:HD12	1.78	0.66
1:A:103:LEU:HD21	1:A:165:VAL:HG13	1.79	0.64
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.32	0.63
1:G:215:LEU:HD22	1:G:261:VAL:HG22	1.80	0.63
1:G:274:TRP:CD1	1:G:275:GLU:O	2.51	0.63
1:G:103:LEU:HD21	1:G:165:VAL:HG13	1.81	0.63
1:G:199:VAL:HG23	1:G:251:LEU:CD1	2.28	0.62
1:D:191:HIS:NE2	1:D:199:VAL:HG11	2.14	0.61
1:G:141:GLN:NE2	4:G:2014:HOH:O	2.33	0.61
1:D:231:VAL:HG13	1:D:244:TRP:CZ2	2.36	0.60
1:J:33:PHE:CD2	1:J:34:VAL:HG13	2.36	0.60
1:J:219:LEU:HD13	1:J:257:TYR:CE2	2.37	0.60
1:A:28:VAL:HG23	1:A:33:PHE:CE1	2.37	0.60
2:B:12:ARG:NH1	2:B:22:ILE:HD12	2.18	0.59
1:G:230:LEU:HD13	1:G:245:ALA:HB2	1.84	0.58



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
2:K:46:ILE:HG23	2:K:47:PRO:HD2	1.86	0.58
1:J:81:LEU:HD13	1:J:118:TYR:CD1	2.39	0.58
1:G:86:ASN:HA	1:G:87:GLN:HB3	1.86	0.58
1:J:28:VAL:HG23	1:J:33:PHE:CE1	2.39	0.57
1:A:28:VAL:HG23	1:A:33:PHE:CD1	2.40	0.56
1:A:147:TRP:CZ2	3:C:9:LEU:HD12	2.41	0.56
1:A:114:LEU:HD22	1:A:156:TYR:CG	2.41	0.56
1:G:249:VAL:HG13	1:G:257:TYR:CE1	2.40	0.56
2:B:28:THR:HG22	2:B:63:TYR:HB2	1.88	0.56
1:G:274:TRP:CG	1:G:275:GLU:O	2.59	0.56
1:A:258:THR:HG22	1:A:273:ARG:HB3	1.89	0.55
2:H:12:ARG:NH2	2:H:22:ILE:HD12	2.22	0.55
2:B:40:LEU:HD23	2:B:45:LYS:HA	1.89	0.55
1:J:117:ALA:HB2	2:K:60:TRP:CE2	2.43	0.54
1:J:28:VAL:HG23	1:J:33:PHE:CD1	2.42	0.54
1:G:28:VAL:HG23	1:G:33:PHE:CD1	2.42	0.54
1:A:191:HIS:NE2	1:A:199:VAL:HG11	2.23	0.54
1:D:81:LEU:HD13	1:D:118:TYR:CD1	2.42	0.54
1:G:33:PHE:CD2	1:G:34:VAL:HG13	2.43	0.53
2:H:28:THR:HG22	2:H:63:TYR:HB2	1.90	0.53
2:E:28:THR:HG22	2:E:63:TYR:HB2	1.90	0.53
2:K:24:ASN:HB3	2:K:65:LEU:HD11	1.91	0.53
1:A:255:GLN:HE22	1:A:273:ARG:HD3	1.74	0.52
1:G:235:PRO:HG2	2:H:65:LEU:HD22	1.92	0.52
1:A:138:MET:HE3	1:A:141:GLN:HB2	1.90	0.52
1:D:16:GLY:C	1:D:17:LEU:HD22	2.30	0.52
1:G:228:MET:CE	1:G:245:ALA:HB1	2.39	0.52
2:H:46:ILE:CG2	2:H:49:VAL:HG23	2.40	0.51
1:J:78:LEU:O	1:J:82:LEU:HD13	2.11	0.51
1:J:230:LEU:HD13	1:J:245:ALA:HB2	1.93	0.51
2:H:70:PHE:HA	2:H:71:THR:HB	1.94	0.50
1:D:123:TYR:CE2	3:F:9:LEU:HD22	2.47	0.49
1:J:45:TYR:CE2	1:J:67:ALA:HB2	2.46	0.49
1:A:109:LEU:HD22	1:A:161:GLU:HG2	1.93	0.49
1:D:176:ASN:HB2	1:D:180:LEU:HD12	1.93	0.49
2:H:70:PHE:HA	2:H:71:THR:CB	2.42	0.49
2:H:35:ILE:HD11	2:H:82:VAL:CG1	2.42	0.49
1:J:114:LEU:HD11	1:J:116:PHE:CZ	2.48	0.49
2:K:7:ILE:HB	2:K:93:VAL:HG21	1.95	0.49
1:A:215:LEU:HD22	1:A:261:VAL:HG22	1.95	0.49
2:H:54:MET:HG3	2:H:64:ILE:HD12	1.94	0.48



	A h o	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
3:L:5:ASN:HD22	3:L:5:ASN:N	2.11	0.48
2:E:7:ILE:HB	2:E:93:VAL:HG21	1.95	0.48
2:K:35:ILE:HD13	2:K:37:ILE:HD11	1.94	0.48
1:D:258:THR:HG22	1:D:273[A]:ARG:HG3	1.96	0.48
2:H:46:ILE:HG23	2:H:47:PRO:HB2	1.95	0.48
1:G:25:VAL:HG22	1:G:35:ARG:HD2	1.96	0.48
1:J:163:GLU:N	1:J:163:GLU:OE2	2.47	0.48
1:G:82:LEU:HA	1:G:87:GLN:HG2	1.96	0.47
1:G:86:ASN:CA	1:G:87:GLN:CB	2.92	0.47
1:G:175:GLY:O	1:G:177:ALA:N	2.47	0.47
1:A:135:ALA:HB1	1:A:140:ALA:HB3	1.96	0.47
1:D:231:VAL:CG1	1:D:244:TRP:CZ2	2.96	0.47
1:J:191:HIS:NE2	1:J:199:VAL:HG11	2.29	0.47
1:D:45:TYR:CE2	1:D:67:ALA:HB2	2.50	0.47
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.50	0.47
1:A:114:LEU:HD23	1:A:126:LEU:HB2	1.96	0.47
1:J:219:LEU:HD13	1:J:257:TYR:CZ	2.50	0.47
1:D:258:THR:HG22	1:D:273[A]:ARG:CG	2.45	0.47
1:A:191:HIS:CD2	1:A:199:VAL:HG11	2.50	0.46
2:E:27:VAL:HG11	2:E:35:ILE:CD1	2.45	0.46
1:J:147:TRP:CZ2	3:L:9:LEU:HD12	2.50	0.46
2:H:40:LEU:HD21	2:H:81:ARG:NH2	2.29	0.46
1:G:211:ALA:HB2	1:G:241:PHE:CE1	2.51	0.46
1:G:249:VAL:HG12	1:G:250:PRO:O	2.15	0.46
1:G:228:MET:HE2	1:G:245:ALA:HB1	1.98	0.46
1:A:100:GLY:O	1:A:160:LEU:HD22	2.15	0.46
1:D:28:VAL:HG23	1:D:33:PHE:CE1	2.50	0.46
1:G:86:ASN:N	1:G:87:GLN:HB2	2.30	0.46
1:A:87:GLN:NE2	1:A:118:TYR:OH	2.48	0.45
1:D:228:MET:CE	1:D:245:ALA:CB	2.90	0.45
1:A:114:LEU:HD22	1:A:156:TYR:CB	2.47	0.45
1:D:191:HIS:CD2	1:D:199:VAL:HG11	2.52	0.45
1:D:272:LEU:HD12	1:D:272:LEU:N	2.31	0.45
2:E:40:LEU:HD23	2:E:45:LYS:HA	1.99	0.45
1:J:135:ALA:HB1	1:J:140:ALA:HB3	1.98	0.45
1:J:272:LEU:HD23	1:J:272:LEU:N	2.31	0.45
2:E:27:VAL:HG11	2:E:35:ILE:HD11	1.98	0.45
1:G:77:SER:O	1:G:81:LEU:HB2	2.17	0.45
2:B:12:ARG:CZ	2:B:22:ILE:HD12	2.47	0.45
1:G:197:GLY:C	1:G:251:LEU:HD13	2.37	0.45
1:A:25:VAL:HG22	1:A:35:ARG:HG3	1.98	0.45



		Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
1:A:73:TRP:CH2	3:C:9:LEU:HD13	2.52	0.45
1:G:81:LEU:HD22	1:G:118:TYR:CE1	2.51	0.45
2:H:5:PRO:HB2	2:H:27:VAL:HG13	1.98	0.45
1:D:235:PRO:HG2	2:E:65:LEU:HD22	2.00	0.44
1:G:82:LEU:O	1:G:87:GLN:HB3	2.17	0.44
2:H:71:THR:N	2:H:72:PRO:HD3	2.33	0.44
1:G:191:HIS:HE1	1:G:199:VAL:HG22	1.82	0.44
1:A:135:ALA:HB1	1:A:140:ALA:CB	2.48	0.43
1:G:138:MET:O	1:G:139:ALA:HB3	2.18	0.43
1:G:103:LEU:CD2	1:G:165:VAL:HG13	2.46	0.43
1:A:137:ASP:O	1:A:141:GLN:HG2	2.19	0.43
1:A:109:LEU:HD22	1:A:161:GLU:CG	2.48	0.43
2:H:46:ILE:HG21	2:H:49:VAL:HG23	2.01	0.43
1:G:86:ASN:HA	1:G:87:GLN:CB	2.46	0.43
1:A:119:GLU:O	2:B:1:MET:HE2	2.19	0.43
2:K:27:VAL:HG12	2:K:30:PHE:CE1	2.54	0.43
1:D:28:VAL:HG23	1:D:33:PHE:CD1	2.55	0.42
1:J:82:LEU:HD11	1:J:93:HIS:CD2	2.54	0.42
1:J:206:LEU:HD22	1:J:242:GLN:NE2	2.35	0.42
2:H:39:MET:C	2:H:40:LEU:HD12	2.40	0.42
1:D:5:MET:HB2	1:D:168:LEU:HD13	2.02	0.42
1:J:112:GLY:HA3	1:J:160:LEU:HD13	2.01	0.42
1:D:187:ALA:CB	1:D:272:LEU:HD21	2.45	0.42
1:G:199:VAL:N	1:G:249:VAL:O	2.53	0.42
1:G:85:TYR:CE2	1:G:139:ALA:CB	3.03	0.42
2:K:9:VAL:HG12	2:K:23:LEU:HD11	2.02	0.42
1:G:81:LEU:HD23	1:G:81:LEU:O	2.20	0.41
1:A:118:TYR:CD2	1:A:119:GLU:HG2	2.55	0.41
1:G:81:LEU:HD22	1:G:118:TYR:CZ	2.55	0.41
1:G:191:HIS:ND1	1:G:199:VAL:HG13	2.35	0.41
1:D:206:LEU:HD22	1:D:242:GLN:HG3	2.03	0.41
1:G:160:LEU:O	1:G:165:VAL:HG23	2.21	0.41
1:G:194:ARG:HB2	1:G:200:THR:HG23	2.03	0.41
1:D:249:VAL:HG13	1:D:250:PRO:HD2	2.03	0.41
1:A:192:HIS:NE2	2:B:98:ASP:O	2.54	0.40
1:G:187:ALA:CB	1:G:272:LEU:CD1	2.97	0.40
2:H:79:ALA:HB2	2:H:94:TYR:CD2	2.55	0.40
1:J:100:GLY:O	1:J:160:LEU:HD22	2.21	0.40
1:J:231:VAL:HG11	1:J:244:TRP:CD1	2.56	0.40
2:K:35:ILE:HD11	2:K:82:VAL:CG1	2.50	0.40
2:H:46:ILE:HG22	2:H:47:PRO:CA	2.50	0.40



a 1	C		
Continued	trom	previous	paae
	J	<i>P</i> · · · · · · · · · · · · · · · · · · ·	r - g - · · ·

Atom-1 Atom-2		Interatomic distance (Å)	Clash overlap (Å)
1:G:85:TYR:C	1:G:87:GLN:HB2	2.42	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	276/277~(100%)	264 (96%)	12 (4%)	0	100	100
1	D	276/277~(100%)	270 (98%)	6 (2%)	0	100	100
1	G	274/277~(99%)	254 (93%)	16 (6%)	4 (2%)	10	15
1	J	274/277~(99%)	263~(96%)	11 (4%)	0	100	100
2	В	97/99~(98%)	95~(98%)	2 (2%)	0	100	100
2	Е	98/99~(99%)	95~(97%)	3 (3%)	0	100	100
2	Н	97/99~(98%)	94 (97%)	2 (2%)	1 (1%)	15	23
2	K	97/99~(98%)	94 (97%)	3(3%)	0	100	100
3	С	6/9~(67%)	5 (83%)	1 (17%)	0	100	100
3	F	6/9~(67%)	5 (83%)	1 (17%)	0	100	100
3	Ι	6/9~(67%)	5 (83%)	0	1 (17%)	0	0
3	L	6/9 ( $67%$ )	6 (100%)	0	0	100	100
All	All	$151\overline{3}/1540~(98\%)$	1450 (96%)	57 (4%)	6 (0%)	34	48

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	176	ASN
2	Н	47	PRO
1	G	86	ASN



 $Continued \ from \ previous \ page...$ 

Mol	Chain	Res	Type
3	Ι	8	ALA
1	G	87	GLN
1	G	275	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	237/235~(101%)	230~(97%)	7 (3%)	41 59		
1	D	236/235~(100%)	231~(98%)	5(2%)	53 72		
1	G	234/235~(100%)	226~(97%)	8~(3%)	37 53		
1	J	234/235~(100%)	230~(98%)	4 (2%)	60 77		
2	В	93/93~(100%)	91~(98%)	2(2%)	52 70		
2	Ε	94/93~(101%)	92~(98%)	2(2%)	53 72		
2	Η	93/93~(100%)	92~(99%)	1 (1%)	73 85		
2	Κ	93/93~(100%)	91~(98%)	2(2%)	52 70		
3	С	5/5~(100%)	4 (80%)	1 (20%)	1 1		
3	F	5/5~(100%)	5~(100%)	0	100 100		
3	Ι	5/5~(100%)	5~(100%)	0	100 100		
3	L	5/5~(100%)	5 (100%)	0	100 100		
All	All	1334/1332~(100%)	1302 (98%)	32 (2%)	50 67		

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	41	GLU
1	А	99	SER
1	А	108	ARG
1	А	149[A]	GLN
1	А	149[B]	GLN
1	А	231	VAL
1	А	275	GLU



Mol	Chain	Res	Type
2	В	48	LYS
2	В	70	PHE
3	С	9	LEU
1	D	111	ARG
1	D	114	LEU
1	D	196	LYS
1	D	226	GLN
1	D	272	LEU
2	Е	4	THR
2	Е	53	ASP
1	G	35	ARG
1	G	48	ARG
1	G	65	GLN
1	G	72	GLN
1	G	87	GLN
1	G	98	MET
1	G	180	LEU
1	G	226	GLN
2	Н	70	PHE
1	J	39	ASP
1	J	178	THR
1	J	183	ASP
1	J	272	LEU
2	K	70	PHE
2	Κ	87	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	65	GLN
1	А	87	GLN
1	А	97	GLN
2	В	29	GLN
3	С	5	ASN
1	D	54	GLN
1	D	87	GLN
1	D	97	GLN
1	D	226	GLN
1	D	255	GLN
2	Е	8	GLN
2	Е	29	GLN
3	F	5	ASN



	5	1	1 5
$\mathbf{Mol}$	Chain	Res	Type
1	G	72	GLN
1	G	97	GLN
1	G	127	ASN
1	G	191	HIS
1	G	226	GLN
3	Ι	5	ASN
1	J	72	GLN
1	J	97	GLN
1	J	242	GLN
3	L	5	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal Trees		Chain	Dec	T:nl.	Bond lengths			Bond angles							
	туре	Chain	nes	nes	nes	nes	nes	Res		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PRQ	L	7	3	13,14,15	<b>3.03</b>	3 (23%)	12,18,20	1.48	1 (8%)					
3	PRQ	С	7	3	13,14,15	2.92	3 (23%)	12,18,20	1.76	3 (25%)					
3	PRQ	F	7	3	13,14,15	<mark>3.03</mark>	3 (23%)	12,18,20	1.49	3 (25%)					
3	PRQ	Ι	7	3	13,14,15	3.00	3 (23%)	12,18,20	1.61	3 (25%)					

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	$\mathbf{PRQ}$	L	7	3	-	2/9/11/12	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PRQ	С	7	3	-	2/9/11/12	0/1/1/1
3	PRQ	F	7	3	-	2/9/11/12	0/1/1/1
3	PRQ	Ι	7	3	-	2/9/11/12	0/1/1/1

Continued from previous page...

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
3	F	7	PRQ	OAC-NAN	8.18	1.36	1.22
3	L	7	PRQ	OAC-NAN	8.17	1.36	1.22
3	Ι	7	PRQ	OAC-NAN	8.07	1.36	1.22
3	С	7	PRQ	OAC-NAN	7.72	1.35	1.22
3	С	7	PRQ	CAL-NAN	-6.66	1.33	1.45
3	Ι	7	PRQ	CAL-NAN	-6.63	1.33	1.45
3	L	7	PRQ	CAL-NAN	-6.63	1.33	1.45
3	F	7	PRQ	CAL-NAN	-6.59	1.33	1.45
3	F	7	PRQ	CAJ-C	2.44	1.56	1.49
3	L	7	PRQ	CAJ-C	2.33	1.55	1.49
3	Ι	7	PRQ	CAJ-C	2.31	1.55	1.49
3	C	7	PRQ	CAJ-C	2.29	1.55	1.49

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	С	7	PRQ	CAL-CAK-CA	-3.73	120.13	124.58
3	L	7	PRQ	CAL-CAK-CA	-3.42	120.50	124.58
3	F	7	PRQ	CAL-CAK-CA	-3.33	120.60	124.58
3	Ι	7	PRQ	CAL-CAK-CA	-3.15	120.83	124.58
3	С	7	PRQ	CAI-CAL-NAN	2.86	119.53	116.47
3	Ι	7	PRQ	CAI-CAL-NAN	2.61	119.27	116.47
3	Ι	7	PRQ	CAH-CAK-CAL	2.51	119.74	117.05
3	С	7	PRQ	CAH-CAK-CAL	2.44	119.67	117.05
3	F	7	PRQ	CAH-CAK-CAL	2.23	119.44	117.05
3	F	7	PRQ	CAI-CAL-NAN	2.07	118.69	116.47

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	С	7	PRQ	CAK-CAL-NAN-OAC
3	С	7	PRQ	CAI-CAL-NAN-OAC



Mol	Chain	Res	Type	Atoms
3	F	7	PRQ	CAK-CAL-NAN-OAC
3	F	7	PRQ	CAI-CAL-NAN-OAC
3	L	7	PRQ	CAI-CAL-NAN-OAC
3	Ι	7	PRQ	CAI-CAL-NAN-OAC
3	Ι	7	PRQ	CAK-CAL-NAN-OAC
3	L	7	PRQ	CAK-CAL-NAN-OAC

 $Continued \ from \ previous \ page...$ 

There are no ring outliers.

No monomer is involved in short contacts.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

#### 5.7 Other polymers (i)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{\AA}^2)$	Q<0.9
1	А	276/277~(99%)	-0.14	3 (1%) 80 79	46,54,61,64	0
1	D	276/277~(99%)	-0.20	0 100 100	47, 53, 59, 61	0
1	G	276/277~(99%)	0.48	19 (6%) 16 13	49,57,63,64	0
1	J	276/277~(99%)	0.33	12 (4%) 35 31	51,58,63,66	0
2	В	99/99~(100%)	-0.24	0 100 100	48, 52, 57, 62	0
2	Е	99/99~(100%)	-0.11	1 (1%) 82 81	46, 52, 57, 59	0
2	Н	99/99~(100%)	0.59	8 (8%) 12 9	56, 59, 62, 63	0
2	K	99/99~(100%)	0.48	5 (5%) 28 25	52,55,63,64	0
3	С	8/9~(88%)	-0.15	0 100 100	40, 41, 41, 41	0
3	F	8/9~(88%)	0.23	0 100 100	37, 41, 41, 41	0
3	Ι	8/9~(88%)	0.16	0 100 100	66,66,67,67	0
3	L	8/9~(88%)	0.29	0 100 100	68, 68, 70, 70	0
All	All	1532/1540~(99%)	0.13	48 (3%) 49 45	37, 55, 62, 70	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	105	SER	7.9
2	Κ	1	MET	5.8
1	G	105	SER	5.0
1	J	250	PRO	4.1
2	Н	54	MET	3.8
1	G	199	VAL	3.3
1	G	250	PRO	3.2
2	Е	1	MET	3.1
1	G	274	TRP	3.0
1	G	192	HIS	2.9
2	Н	68	THR	2.9



Continued from previous page								
Mol	Chain	Res	Type	RSRZ				
1	J	107	TRP	2.9				
1	J	196	LYS	2.9				
1	G	249	VAL	2.9				
2	Н	1	MET	2.8				
2	Н	69	GLU	2.8				
1	G	106	ASP	2.8				
1	J	239	GLY	2.7				
2	Κ	22	ILE	2.6				
1	G	145	ARG	2.6				
2	Κ	18	GLY	2.6				
1	G	270	LEU	2.6				
1	J	249	VAL	2.6				
1	G	133	TRP	2.5				
1	J	182	THR	2.5				
1	J	257	TYR	2.5				
1	G	1	GLY	2.4				
1	G	154	GLU	2.4				
1	J	104	GLY	2.4				
1	J	200	THR	2.3				
2	Н	55	SER	2.3				
1	А	1	GLY	2.3				
2	Н	20	PRO	2.3				
1	А	105	SER	2.3				
1	J	251	LEU	2.3				
1	G	253	LYS	2.3				
1	G	276	PRO	2.2				
2	Н	2	GLN	2.2				
2	Н	70	PHE	2.2				
1	G	173	LYS	2.1				
1	J	274	TRP	2.1				
1	G	195	SER	2.1				
1	G	225	THR	2.1				
1	G	44	ARG	2.1				
2	K	69	GLU	2.1				
2	K	98	ASP	2.0				
1	G	204	TRP	2.0				
1	А	276[A]	PRO	2.0				

1 L.  $\alpha$ · ·

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	$Q{<}0.9$
3	PRQ	Ι	7	14/15	0.88	0.20	$66,\!67,\!67,\!67$	0
3	PRQ	L	7	14/15	0.91	0.22	68,70,70,70	0
3	PRQ	F	7	14/15	0.93	0.19	37,40,41,41	0
3	PRQ	С	7	14/15	0.95	0.17	41,43,45,46	0

median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

There are no ligands in this entry.

### 6.5 Other polymers (i)

There are no such residues in this entry.

