



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 29, 2024 – 02:05 PM EDT

PDB ID : 8VER  
Title : Structure of YicC endoribonuclease  
Authors : Dahlin, H.R.; Khamrui, S.; Lazarus, M.B.  
Deposited on : 2023-12-20  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3657 (2.80-2.80)
Clashscore	180529	4123 (2.80-2.80)
Ramachandran outliers	177936	4071 (2.80-2.80)
Sidechain outliers	177891	4073 (2.80-2.80)
RSRZ outliers	164620	3659 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	 2% 91% 8%
1	B	289	 0% 89% 10%
1	C	289	 0% 86% 11%
1	D	289	 2% 91% 6%
1	E	289	 0% 91% 8%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	289	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment at the beginning (representing 1% of the chain), a large green segment (representing 89%), a yellow segment (representing 7%), and a small grey segment at the end (representing 4%). A percentage sign (%) is located at the top left of the bar, and a dot (•) is at the top right. The values 89% and 7% are printed below the green and yellow segments respectively.</p>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 13856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoribonuclease YicC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2325	1450	418	447	10	0	0	0
1	B	287	2329	1452	419	448	10	0	0	0
1	C	282	2288	1427	412	439	10	0	0	0
1	D	282	2287	1426	412	439	10	0	0	0
1	E	286	2324	1449	418	447	10	0	1	0
1	F	277	2243	1401	402	430	10	0	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P23839
A	0	SER	-	expression tag	UNP P23839
A	187	THR	ALA	engineered mutation	UNP P23839
B	-1	GLY	-	expression tag	UNP P23839
B	0	SER	-	expression tag	UNP P23839
B	187	THR	ALA	engineered mutation	UNP P23839
C	-1	GLY	-	expression tag	UNP P23839
C	0	SER	-	expression tag	UNP P23839
C	187	THR	ALA	engineered mutation	UNP P23839
D	-1	GLY	-	expression tag	UNP P23839
D	0	SER	-	expression tag	UNP P23839
D	187	THR	ALA	engineered mutation	UNP P23839
E	-1	GLY	-	expression tag	UNP P23839
E	0	SER	-	expression tag	UNP P23839
E	187	THR	ALA	engineered mutation	UNP P23839
F	-1	GLY	-	expression tag	UNP P23839
F	0	SER	-	expression tag	UNP P23839

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	187	THR	ALA	engineered mutation	UNP P23839

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>			<b>ZeroOcc</b>	<b>AltConf</b>
2	D	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Endoribonuclease YicC




- Molecule 1: Endoribonuclease YicC




- Molecule 1: Endoribonuclease YicC



- Molecule 1: Endoribonuclease YicC





- Molecule 1: Endoribonuclease YicC





- Molecule 1: Endoribonuclease YicC





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.92Å 76.54Å 136.34Å 90.00° 100.29° 90.00°	Depositor
Resolution (Å)	38.34 – 2.80 38.34 – 2.80	Depositor EDS
% Data completeness (in resolution range)	65.8 (38.34-2.80) 65.9 (38.34-2.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.226 , 0.267 0.226 , 0.267	Depositor DCC
$R_{free}$ test set	2853 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	88.7	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 49.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	13856	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	0/2351	0.47	0/3168
1	B	0.23	0/2356	0.47	0/3176
1	C	0.23	0/2314	0.47	0/3117
1	D	0.23	0/2313	0.47	0/3116
1	E	0.23	0/2350	0.48	0/3166
1	F	0.23	0/2269	0.47	0/3057
All	All	0.23	0/13953	0.47	0/18800

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2325	0	2359	14	0
1	B	2329	0	2363	18	0
1	C	2288	0	2322	18	0
1	D	2287	0	2320	10	0
1	E	2324	0	2355	13	0
1	F	2243	0	2278	12	0
2	A	30	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	10	0	0	0	0
2	D	15	0	0	0	0
2	F	5	0	0	0	0
All	All	13856	0	13997	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:29:GLN:HG3	1:C:61:LYS:HB2	1.75	0.66
1:E:2:ILE:HG22	1:E:238:VAL:HG21	1.77	0.65
1:A:281:GLU:HG2	1:C:31:TYR:HB3	1.81	0.62
1:B:89:VAL:O	1:B:93:ASN:ND2	2.32	0.62
1:B:171:ARG:NH2	1:B:219:ASP:OD1	2.35	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/289 (98%)	277 (98%)	5 (2%)	0	100	100
1	B	285/289 (99%)	279 (98%)	6 (2%)	0	100	100
1	C	278/289 (96%)	274 (99%)	4 (1%)	0	100	100
1	D	278/289 (96%)	275 (99%)	3 (1%)	0	100	100
1	E	282/289 (98%)	281 (100%)	1 (0%)	0	100	100
1	F	273/289 (94%)	270 (99%)	3 (1%)	0	100	100
All	All	1678/1734 (97%)	1656 (99%)	22 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/252 (100%)	250 (100%)	1 (0%)	89	96
1	B	251/252 (100%)	251 (100%)	0	100	100
1	C	246/252 (98%)	245 (100%)	1 (0%)	89	96
1	D	246/252 (98%)	245 (100%)	1 (0%)	89	96
1	E	250/252 (99%)	250 (100%)	0	100	100
1	F	241/252 (96%)	241 (100%)	0	100	100
All	All	1485/1512 (98%)	1482 (100%)	3 (0%)	92	97

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ARG
1	C	176	GLU
1	D	6	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	93	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	303	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	C	301	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	C	302	-	4,4,4	0.23	0	6,6,6	0.07	0
2	SO4	D	303	-	4,4,4	0.23	0	6,6,6	0.09	0
2	SO4	D	301	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	D	302	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	A	306	-	4,4,4	0.24	0	6,6,6	0.08	0
2	SO4	A	301	-	4,4,4	0.24	0	6,6,6	0.07	0
2	SO4	A	304	-	4,4,4	0.23	0	6,6,6	0.07	0
2	SO4	A	305	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	A	302	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	F	301	-	4,4,4	0.23	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## Ligand SO4 C 302




Bond lengths



Bond angles




Torsions




Rings




## Ligand SO4 D 303




Bond lengths




Bond angles




Torsions




Rings




## Ligand SO4 D 302




Bond lengths



Bond angles




Torsions




Rings


## Ligand SO4 A 306




Bond lengths




Bond angles




Torsions




Rings




## Ligand SO4 A 304




Bond lengths




Bond angles




Torsions




Rings




## Ligand SO4 A 302




Bond lengths



Bond angles




Torsions



Rings





### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/289 (98%)	-0.16	5 (1%) 69 61	38, 71, 125, 185	0
1	B	287/289 (99%)	0.05	2 (0%) 84 79	48, 99, 144, 193	0
1	C	282/289 (97%)	-0.08	3 (1%) 77 71	40, 75, 139, 168	0
1	D	282/289 (97%)	0.24	6 (2%) 63 55	63, 97, 146, 214	0
1	E	286/289 (98%)	0.19	3 (1%) 79 73	59, 104, 172, 218	1 (0%)
1	F	277/289 (95%)	0.07	3 (1%) 77 71	56, 90, 131, 177	0
All	All	1700/1734 (98%)	0.05	22 (1%) 74 67	38, 90, 148, 218	1 (0%)

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	73	SER	3.6
1	E	42	PHE	3.3
1	D	118	ALA	3.2
1	D	64	CYS	3.2
1	A	195	GLN	3.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	306	5/5	0.69	0.14	111,111,115,126	0
2	SO4	D	302	5/5	0.77	0.11	91,91,108,111	0
2	SO4	C	301	5/5	0.79	0.10	99,104,109,110	0
2	SO4	A	301	5/5	0.81	0.11	110,110,110,114	0
2	SO4	A	304	5/5	0.82	0.10	68,76,85,99	0
2	SO4	D	303	5/5	0.82	0.09	73,74,84,96	0
2	SO4	F	301	5/5	0.83	0.06	122,127,136,147	0
2	SO4	D	301	5/5	0.84	0.11	92,93,112,112	0
2	SO4	A	305	5/5	0.85	0.12	90,99,114,119	0
2	SO4	A	302	5/5	0.87	0.13	82,82,104,111	0
2	SO4	C	302	5/5	0.87	0.09	77,78,102,110	0
2	SO4	A	303	5/5	0.94	0.11	54,54,64,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around SO4 A 306:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 D 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 C 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SO4 A 301:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around SO4 A 304:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around SO4 D 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 F 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 D 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SO4 A 305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SO4 A 302:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around SO4 C 302:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SO4 A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.