

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 5, 2023 – 07:17 AM EDT

PDB ID	:	6VFV
Title	:	Crystal structure of human protocadherin 8 EC5-EC6
Authors	:	Harrison, O.J.; Brasch, J.; Shapiro, L.
Deposited on		
Resolution	:	2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\hbox{-}RAY\,DIFFRACTION$ 

The reported resolution of this entry is 2.90 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 3445 atoms, of which 1688 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protocadherin-8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	А	222	Total 3292	C 1033	H 1622	N 306	O 331	0	0	0

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
А	684	HIS	-	expression tag	UNP O95206
А	685	HIS	-	expression tag	UNP O95206
А	686	HIS	-	expression tag	UNP O95206
А	687	HIS	-	expression tag	UNP O95206
А	688	HIS	-	expression tag	UNP O95206
А	689	HIS	-	expression tag	UNP O95206

• Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-6)-beta-D-mannopyranos e-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acet amido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	5	Total 111	C 34	Н 51	N 2	0 24	0	0	0

• Molecule 3 is alpha-D-mannopyranose (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	А	1	Total	С	Η	Ο	0	0
0	11	Ĩ	20	6	9	5	Ŭ	Ŭ

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	4	Total Ca 4 4	0	0

• Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	А	1	Total 10	$\begin{array}{c} \mathrm{C} \\ \mathrm{2} \end{array}$	Н 6	0 2	0	0

• Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	1	Total Cl 1 1	0	0

• Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	А	7	Total O 7 7	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	41.66Å 29.50Å 92.44Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $98.46^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	19.90 - 2.90	Depositor	
% Data completeness	99.5 (19.90-2.90)	Depositor	
(in resolution range)			
R <sub>merge</sub>	0.13	Depositor	
R <sub>sym</sub>	(Not available)	Depositor	
$< I/\sigma(I) > 1$	1.23 (at 2.90 Å)	Xtriage	
Refinement program	PHENIX 1.10.1_2155	Depositor	
$R, R_{free}$	0.225 , $0.264$	Depositor	
Wilson B-factor $(Å^2)$	39.4	Xtriage	
Anisotropy	0.812	Xtriage	
L-test for twinning <sup>2</sup>	$ < L >=0.49, < L^2>=0.32$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3445	wwPDB-VP	
Average B, all atoms $(Å^2)$	52.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.65% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 4 Model quality (i)

## 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bo	ond leng	$\mathbf{ths}$	B	ond ang	les
10101		Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	NAG	В	1	2,1	14,14,15	0.26	0	17,19,21	0.56	0
2	NAG	В	2	2	14,14,15	0.46	0	17,19,21	0.55	0
2	BMA	В	3	2	11,11,12	1.58	3 (27%)	15,15,17	2.08	2 (13%)
2	MAN	В	4	2	11,11,12	0.85	0	15,15,17	1.36	2 (13%)
2	FUC	В	5	2	10,10,11	1.25	2 (20%)	14,14,16	1.53	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	В	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	В	2	2	-	1/6/23/26	0/1/1/1
2	BMA	В	3	2	-	1/2/19/22	0/1/1/1
2	MAN	В	4	2	-	2/2/19/22	0/1/1/1
2	FUC	В	5	2	-	-	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\mathrm{Ideal}(\mathrm{\AA})$
2	В	3	BMA	O5-C5	3.55	1.50	1.43
2	В	5	FUC	O5-C5	2.83	1.49	1.43
2	В	5	FUC	C1-C2	2.31	1.57	1.52
2	В	3	BMA	C4-C5	2.29	1.57	1.53
2	В	3	BMA	O5-C1	2.01	1.46	1.43

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\mathbf{Ideal}(^{o})$
2	В	3	BMA	C1-O5-C5	6.99	121.66	112.19
2	В	4	MAN	C1-O5-C5	4.20	117.88	112.19
2	В	5	FUC	C1-O5-C5	3.09	119.77	112.78
2	В	5	FUC	C1-C2-C3	2.79	113.09	109.67
2	В	5	FUC	O5-C1-C2	2.37	114.43	110.77

There are no chirality outliers.

5 of 6 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
2	В	1	NAG	O5-C5-C6-O6
2	В	1	NAG	C4-C5-C6-O6
2	В	3	BMA	O5-C5-C6-O6
2	В	4	MAN	C4-C5-C6-O6
2	В	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 4.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



	Mol Type	Type Chain		in Res	Link	Bo	ond leng	$\mathbf{ths}$	Bond angles		
		Chain			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
	5	EDO	А	711	-	$3,\!3,\!3$	0.47	0	2,2,2	0.23	0
	3	MAN	А	706	1	$11,\!11,\!12$	0.72	0	$15,\!15,\!17$	0.96	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	А	711	-	-	0/1/1/1	-
3	MAN	А	706	1	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mo	l Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	A	706	MAN	C1-O5-C5	2.30	115.31	112.19
3	А	706	MAN	O2-C2-C3	-2.11	105.92	110.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 4.7 Other polymers (i)

There are no such residues in this entry.

## 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 5 Fit of model and data (i)

## 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

## 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

