



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 01:02 am BST

PDB ID : 1VGF  
Title : volvatoxin A2 (diamond crystal form)  
Authors : Lin, S.-C.; Lo, Y.-C.; Lin, J.-Y.; Liaw, Y.-C.  
Deposited on : 2004-04-24  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 62%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 31%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">3%      62%      31%      . . .</p>
1	B	199	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 64%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 31%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: orange; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 10%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 10px;">3%      64%      31%      . .</p>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called volvatoxin A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	194	Total 1547	C 1004	N 241	O 301	S 1	0	0	0
1	B	194	Total 1547	C 1004	N 241	O 301	S 1	0	0	0

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	14	Total	O	0	0
			14	14		
3	B	20	Total	O	0	0
			20	20		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: volvatoxin A2



- Molecule 1: volvatoxin A2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.31Å 56.31Å 264.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.54 – 2.60 28.54 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.2 (28.54-2.60) 99.2 (28.54-2.60)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.23 (at 2.61Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.214 , 0.268 0.208 , 0.262	Depositor DCC
$R_{free}$ test set	1371 reflections (9.77%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.4	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3156	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/1584	0.61	0/2150
1	B	0.39	0/1584	0.58	0/2150
All	All	0.39	0/3168	0.60	0/4300

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	31	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1547	0	1515	51	0
1	B	1547	0	1515	50	0
2	A	12	0	9	0	0
2	B	16	0	12	0	0
3	A	14	0	0	0	0
3	B	20	0	0	0	0
All	All	3156	0	3051	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:LEU:H	1:A:99:LEU:HD23	1.33	0.91
1:B:1126:VAL:HG23	1:B:1127:LYS:H	1.41	0.84
1:B:1098:LYS:O	1:B:1102:ILE:HG12	1.79	0.82
1:A:6:VAL:HG21	1:A:29:GLY:HA2	1.64	0.78
1:A:70:GLN:HG2	1:A:181:ALA:HB2	1.69	0.75
1:A:98:LYS:N	1:A:98:LYS:HD2	2.05	0.70
1:B:1012:GLN:HB2	1:B:1140:VAL:HG12	1.73	0.70
1:B:1070:GLN:HG2	1:B:1181:ALA:HB2	1.75	0.69
1:B:1126:VAL:HG23	1:B:1127:LYS:N	2.07	0.69
1:A:53:TYR:CE2	1:A:58:LEU:HB3	2.31	0.65
1:B:1039:ALA:H	1:B:1120:GLN:NE2	1.94	0.64
1:A:98:LYS:H	1:A:98:LYS:HD2	1.63	0.63
1:B:1075:VAL:O	1:B:1079:VAL:HG23	1.98	0.63
1:B:1169:TRP:O	1:B:1172:LEU:HD13	1.98	0.63
1:A:6:VAL:HG21	1:A:29:GLY:CA	2.29	0.61
1:B:1160:THR:HB	1:B:1181:ALA:HB3	1.82	0.61
1:A:72:SER:HA	1:A:178:LYS:O	2.01	0.61
1:A:57:ASP:HA	1:A:192:LYS:HG2	1.81	0.60
1:B:1006:VAL:HG13	1:B:1032:LEU:HD22	1.84	0.59
1:B:1068:ILE:N	1:B:1068:ILE:HD12	2.17	0.59
1:B:1161:ALA:HB2	1:B:1180:PHE:CD1	2.36	0.59
1:B:1008:GLN:HE22	1:B:1102:ILE:CG2	2.15	0.59
1:A:160:THR:HB	1:A:181:ALA:HB3	1.84	0.59
1:A:64:ASP:HB2	1:A:90:PHE:CZ	2.38	0.58
1:A:6:VAL:HG12	1:A:32:LEU:HD22	1.86	0.58
1:B:1012:GLN:O	1:B:1141:PRO:HD3	2.04	0.57
1:B:1070:GLN:HB3	1:B:1179:ASN:ND2	2.19	0.57
1:A:13:LEU:HD23	1:A:141:PRO:HD3	1.86	0.56

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:TYR:CZ	1:A:55:GLY:HA3	2.40	0.56
1:A:122:ASN:ND2	1:A:123:THR:H	2.02	0.56
1:B:1156:THR:OG1	1:B:1185:ASP:HB2	2.06	0.56
1:A:161:ALA:HB2	1:A:180:PHE:CD1	2.44	0.53
1:A:122:ASN:HD22	1:A:123:THR:H	1.56	0.53
1:B:1146:ALA:HB1	1:B:1149:TYR:CD2	2.44	0.53
1:B:1056:GLU:N	1:B:1056:GLU:OE1	2.42	0.53
1:A:149:TYR:CE1	1:A:192:LYS:HB3	2.43	0.53
1:A:74:GLU:HG2	1:A:175:SER:O	2.09	0.53
1:B:1079:VAL:HG11	1:B:1104:LEU:HB2	1.92	0.52
1:B:1149:TYR:CE1	1:B:1192:LYS:HB2	2.45	0.52
1:B:1146:ALA:HB1	1:B:1149:TYR:HD2	1.75	0.51
1:B:1024:VAL:HG12	1:B:1137:LEU:HD11	1.91	0.51
1:A:146:ALA:N	1:A:147:PRO:HD3	2.26	0.50
1:A:69:ASN:O	1:A:71:GLN:HG3	2.11	0.50
1:A:68:ILE:HD12	1:A:68:ILE:N	2.26	0.50
1:B:1126:VAL:CG2	1:B:1127:LYS:H	2.18	0.49
1:A:39:ALA:H	1:A:120:GLN:NE2	2.10	0.49
1:B:1087:HIS:HA	1:B:1096:LEU:HD23	1.94	0.49
1:A:30:LYS:HG3	1:A:31:TYR:CE2	2.48	0.49
1:B:1104:LEU:O	1:B:1108:THR:HG23	2.12	0.49
1:B:1070:GLN:HB3	1:B:1179:ASN:HD21	1.78	0.48
1:A:121:PHE:CG	1:A:122:ASN:N	2.81	0.48
1:B:1070:GLN:CG	1:B:1181:ALA:HB2	2.41	0.48
1:A:13:LEU:HD11	1:A:139:SER:HB3	1.96	0.48
1:A:172:LEU:CD2	1:A:176:THR:HG21	2.44	0.47
1:A:137:LEU:HD12	1:A:138:PHE:N	2.29	0.47
1:B:1117:GLY:O	1:B:1135:ARG:NH2	2.44	0.47
1:A:104:LEU:O	1:A:108:THR:HG23	2.14	0.47
1:A:172:LEU:HD23	1:A:176:THR:HG21	1.97	0.47
1:A:35:GLU:O	1:A:36:GLN:HB2	2.14	0.47
1:A:137:LEU:HD12	1:A:137:LEU:C	2.35	0.46
1:B:1062:LYS:HE2	1:B:1064:ASP:OD1	2.15	0.46
1:A:124:ASN:C	1:A:124:ASN:HD22	2.18	0.46
1:B:1161:ALA:HB2	1:B:1180:PHE:CE1	2.50	0.46
1:A:162:ASP:OD1	1:A:179:ASN:HB3	2.17	0.45
1:A:187:LEU:HD23	1:A:187:LEU:C	2.36	0.45
1:A:178:LYS:HB2	1:A:180:PHE:CZ	2.52	0.45
1:B:1071:GLN:NE2	1:B:1081:LYS:NZ	2.64	0.45
1:B:1008:GLN:HE22	1:B:1102:ILE:HG21	1.82	0.45
1:A:75:VAL:O	1:A:79:VAL:HG23	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1178:LYS:HB2	1:B:1180:PHE:CE2	2.52	0.44
1:A:38:LYS:HB2	1:A:38:LYS:NZ	2.32	0.44
1:B:1042:ASP:OD1	1:B:1044:PRO:HD2	2.18	0.44
1:B:1126:VAL:O	1:B:1128:LYS:HG2	2.18	0.44
1:A:178:LYS:HB2	1:A:180:PHE:CE2	2.53	0.43
1:B:1148:SER:HB2	1:B:1193:LYS:HB3	2.00	0.43
1:A:50:ILE:HD13	1:A:189:LEU:HD21	1.98	0.43
1:A:156:THR:OG1	1:A:185:ASP:HB2	2.18	0.43
1:B:1162:ASP:N	1:B:1162:ASP:OD1	2.49	0.43
1:B:1086:LEU:HD23	1:B:1099:LEU:HD13	2.01	0.43
1:A:187:LEU:HD23	1:A:188:GLU:N	2.34	0.42
1:B:1126:VAL:HG23	1:B:1127:LYS:HG3	2.01	0.42
1:B:1126:VAL:CG2	1:B:1127:LYS:N	2.78	0.42
1:B:1177:LYS:HZ2	1:B:1177:LYS:HB2	1.84	0.42
1:A:177:LYS:HB2	1:A:177:LYS:HE3	1.92	0.42
1:B:1169:TRP:C	1:B:1172:LEU:HD13	2.39	0.42
1:A:11:ASP:OD1	1:A:139:SER:HB2	2.19	0.42
1:B:1121:PHE:CG	1:B:1122:ASN:N	2.88	0.42
1:A:139:SER:HA	1:A:151:TYR:O	2.20	0.42
1:B:1086:LEU:CD2	1:B:1099:LEU:HD13	2.50	0.42
1:B:1008:GLN:NE2	1:B:1102:ILE:HG21	2.35	0.42
1:A:148:SER:O	1:A:192:LYS:HA	2.21	0.41
1:B:1066:SER:HB2	1:B:1085:PHE:CE1	2.55	0.41
1:A:138:PHE:O	1:A:152:SER:HA	2.20	0.41
1:A:96:LEU:O	1:A:98:LYS:N	2.53	0.41
1:B:1062:LYS:HG2	1:B:1064:ASP:OD1	2.20	0.41
1:A:38:LYS:CB	1:A:38:LYS:NZ	2.84	0.41
1:A:23:GLN:HG3	1:A:195:PHE:O	2.21	0.41
1:A:99:LEU:H	1:A:99:LEU:CD2	2.11	0.41
1:B:1053:TYR:CE2	1:B:1058:LEU:HB3	2.56	0.41
1:B:1014:PRO:HD2	1:B:1017:LEU:HD12	2.02	0.40
1:B:1039:ALA:O	1:B:1120:GLN:HG2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/199 (96%)	177 (92%)	13 (7%)	2 (1%)	15	32
1	B	192/199 (96%)	179 (93%)	9 (5%)	4 (2%)	7	13
All	All	384/398 (96%)	356 (93%)	22 (6%)	6 (2%)	9	19

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	97	SER
1	A	98	LYS
1	B	1114	SER
1	B	1127	LYS
1	B	1015	GLU
1	B	1125	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/175 (98%)	160 (94%)	11 (6%)	17	35
1	B	171/175 (98%)	164 (96%)	7 (4%)	30	56
All	All	342/350 (98%)	324 (95%)	18 (5%)	22	45

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	ASP
1	A	64	ASP
1	A	73	GLN
1	A	97	SER
1	A	98	LYS
1	A	99	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	122	ASN
1	A	124	ASN
1	A	137	LEU
1	A	170	TRP
1	A	191	VAL
1	B	1025	LEU
1	B	1037	ASP
1	B	1056	GLU
1	B	1069	ASN
1	B	1096	LEU
1	B	1099	LEU
1	B	1170	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	GLN
1	A	71	GLN
1	A	73	GLN
1	A	120	GLN
1	A	122	ASN
1	A	124	ASN
1	A	179	ASN
1	A	183	GLN
1	B	1008	GLN
1	B	1052	ASN
1	B	1065	GLN
1	B	1069	ASN
1	B	1071	GLN
1	B	1073	GLN
1	B	1120	GLN
1	B	1179	ASN
1	B	1183	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACT	B	3001	-	1,3,3	2.95	1 (100%)	0,3,3	0.00	-
2	ACT	B	3005	-	1,3,3	2.79	1 (100%)	0,3,3	0.00	-
2	ACT	B	3002	-	1,3,3	3.16	1 (100%)	0,3,3	0.00	-
2	ACT	A	3006	-	1,3,3	3.02	1 (100%)	0,3,3	0.00	-
2	ACT	A	3003	-	1,3,3	2.94	1 (100%)	0,3,3	0.00	-
2	ACT	B	3007	-	1,3,3	2.89	1 (100%)	0,3,3	0.00	-
2	ACT	A	3004	-	1,3,3	3.16	1 (100%)	0,3,3	0.00	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	3004	ACT	CH3-C	3.16	1.52	1.48
2	B	3002	ACT	CH3-C	3.16	1.52	1.48
2	A	3006	ACT	CH3-C	3.02	1.52	1.48
2	B	3001	ACT	CH3-C	2.95	1.52	1.48
2	A	3003	ACT	CH3-C	2.94	1.52	1.48
2	B	3007	ACT	CH3-C	2.89	1.52	1.48
2	B	3005	ACT	CH3-C	2.79	1.52	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	194/199 (97%)	0.01	6 (3%) 49 42	24, 49, 73, 84	0
1	B	194/199 (97%)	0.03	5 (2%) 56 50	28, 49, 70, 80	0
All	All	388/398 (97%)	0.02	11 (2%) 53 46	24, 49, 71, 84	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1126	VAL	4.1
1	A	126	VAL	3.5
1	B	1125	ASN	2.9
1	A	125	ASN	2.9
1	B	1129	ASN	2.5
1	A	198	PRO	2.4
1	A	54	THR	2.4
1	A	177	LYS	2.4
1	B	1056	GLU	2.3
1	A	70	GLN	2.1
1	B	1128	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ACT	B	3001	4/4	0.85	0.16	61,62,63,64	0
2	ACT	B	3002	4/4	0.85	0.21	71,71,71,72	0
2	ACT	A	3004	4/4	0.89	0.14	70,73,73,74	0
2	ACT	A	3003	4/4	0.90	0.29	80,80,80,80	0
2	ACT	B	3007	4/4	0.93	0.21	63,64,65,65	0
2	ACT	B	3005	4/4	0.93	0.12	51,54,54,56	0
2	ACT	A	3006	4/4	0.97	0.21	42,42,43,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.