



# Full wwPDB X-ray Structure Validation Report i

May 25, 2020 – 01:55 pm BST

PDB ID : 3VGB  
Title : Crystal structure of glycosyltrehalose trehalohydrolase (GTHase) from Sulfolobus solfataricus KM1  
Authors : Okazaki, N.; Tamada, T.; Feese, M.D.; Kato, M.; Miura, Y.; Komeda, T.; Kobayashi, K.; Kondo, K.; Kuroki, R.  
Deposited on : 2011-08-09  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.1.3  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	558	<p>2% 75% 19% ..</p>

## 2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 4720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malto-oligosyltrehalose trehalohydrolase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	550	4512	2907	741	855	9	0	0	0

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: C<sub>6</sub>H<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	13	6	7	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	177	Total O 177 177	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Malto-oligosyltrehalose trehalohydrolase



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.27Å    78.27Å    282.10Å 90.00°    90.00°    120.00°	Depositor
Resolution (Å)	67.78 – 2.65 67.78 – 2.65	Depositor EDS
% Data completeness (in resolution range)	97.4 (67.78-2.65) 97.4 (67.78-2.65)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.64 (at 2.65Å)	Xtriage
Refinement program	REFMAC	Depositor
$R$ , $R_{free}$	0.202 , 0.243 0.206 , 0.245	Depositor DCC
$R_{free}$ test set	1486 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.5	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.057 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	4720	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/4628	0.55	0/6253

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4512	0	4369	70	0
2	A	13	0	5	0	0
3	A	18	0	24	3	0
4	A	177	0	0	7	0
All	All	4720	0	4398	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:ILE:HD11	1:A:413:ILE:HD12	1.48	0.95
1:A:354:ARG:HG3	1:A:354:ARG:HH11	1.49	0.76
1:A:264:HIS:HD2	1:A:266:LEU:H	1.36	0.73
1:A:105:TYR:HB2	1:A:133:ILE:HG12	1.71	0.71
1:A:321:GLU:HB3	1:A:323:GLN:NE2	2.06	0.71
1:A:104:ILE:HB	1:A:412:MET:HG3	1.73	0.71
1:A:311:HIS:HD2	1:A:312:HIS:ND1	1.89	0.70
1:A:92:ASN:O	1:A:278:ARG:NH2	2.24	0.69
1:A:143:GLN:HE22	1:A:162:ASN:H	1.42	0.67
1:A:387:ARG:HD2	1:A:423:ASN:O	1.95	0.67
1:A:395:GLU:OE1	1:A:535:ASN:ND2	2.30	0.64
1:A:535:ASN:HB3	1:A:537:SER:H	1.63	0.62
1:A:174:VAL:HB	1:A:184:VAL:HG11	1.82	0.61
1:A:252:ASP:OD1	3:A:1103:GOL:H31	2.00	0.61
1:A:387:ARG:NH2	1:A:415:MET:O	2.34	0.61
1:A:392:VAL:HB	1:A:396:SER:HB2	1.83	0.61
1:A:141:ILE:HD12	1:A:160:VAL:HG13	1.85	0.58
1:A:72:ARG:HG2	4:A:2089:HOH:O	2.05	0.57
1:A:124:LYS:HE2	3:A:1101:GOL:H32	1.87	0.56
1:A:427:PHE:HB3	1:A:450:GLN:HE22	1.72	0.55
1:A:444:ARG:NH1	1:A:450:GLN:HE21	2.05	0.55
1:A:264:HIS:CD2	1:A:266:LEU:H	2.21	0.55
1:A:453:ASP:HB3	1:A:456:ASP:HB2	1.88	0.55
1:A:437:ILE:HG23	1:A:455:GLN:HG3	1.90	0.54
1:A:226:SER:HA	1:A:229:VAL:HG13	1.88	0.54
1:A:354:ARG:HH11	1:A:354:ARG:CG	2.20	0.54
1:A:357:HIS:HE1	4:A:2026:HOH:O	1.91	0.53
1:A:427:PHE:HB2	1:A:444:ARG:NH2	2.25	0.51
1:A:53:ARG:HB3	1:A:84:ILE:O	2.11	0.51
1:A:480:LEU:HD13	1:A:532:LEU:HD23	1.93	0.51
1:A:321:GLU:HB3	1:A:323:GLN:HE21	1.76	0.50
1:A:100:GLU:HG2	4:A:2004:HOH:O	2.12	0.50
1:A:531:LEU:HD21	1:A:557:LYS:HG3	1.94	0.50
1:A:155:VAL:CG2	1:A:198:ASN:HB2	2.42	0.49
1:A:415:MET:O	1:A:415:MET:HG3	2.13	0.49
1:A:16:TRP:HZ2	1:A:225:GLU:HG2	1.79	0.48
1:A:494:ARG:HH12	1:A:509:ARG:HD3	1.79	0.48
1:A:71:SER:CB	1:A:74:GLN:HE22	2.26	0.47
1:A:308:ASP:OD1	1:A:372:TYR:OH	2.27	0.47
1:A:24:LYS:HB2	1:A:59:VAL:HG13	1.95	0.47
1:A:157:LEU:HD11	4:A:2071:HOH:O	2.14	0.47
1:A:333:ASP:HA	1:A:336:VAL:HG13	1.95	0.47

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:332:LEU:HD21	1:A:504:LEU:HD23	1.98	0.46
1:A:444:ARG:HD2	1:A:444:ARG:HA	1.62	0.46
1:A:487:LEU:HG	1:A:511:TYR:HB3	1.96	0.46
1:A:52:VAL:HA	1:A:53:ARG:HA	1.66	0.46
1:A:141:ILE:HG12	1:A:186:LEU:HD11	1.98	0.45
1:A:409:TYR:O	1:A:484:ARG:NH2	2.49	0.45
1:A:402:ALA:HA	1:A:480:LEU:HD21	1.99	0.44
1:A:273:VAL:HG21	1:A:280:VAL:HG22	1.99	0.44
1:A:493:ARG:HD2	4:A:2012:HOH:O	2.17	0.44
1:A:10:GLU:HB2	1:A:48:ASN:HA	1.99	0.44
1:A:268:GLU:O	1:A:272:VAL:HG23	2.18	0.44
1:A:124:LYS:CE	3:A:1101:GOL:H32	2.48	0.43
1:A:89:LYS:HG3	1:A:243:GLU:OE1	2.19	0.43
1:A:194:GLY:HA2	1:A:195:PRO:HD3	1.86	0.43
1:A:295:LYS:HE3	4:A:2157:HOH:O	2.19	0.43
1:A:133:ILE:CD1	1:A:413:ILE:HD12	2.35	0.42
1:A:20:GLN:OE1	1:A:60:LEU:HB3	2.18	0.42
1:A:354:ARG:NH1	1:A:354:ARG:CG	2.82	0.42
1:A:89:LYS:HG2	1:A:242:LYS:HB3	2.00	0.42
1:A:544:GLU:HG2	1:A:544:GLU:H	1.37	0.42
1:A:297:LYS:O	1:A:298:CYS:HB2	2.20	0.41
1:A:122:ILE:HA	1:A:125:LEU:HD22	2.02	0.41
1:A:210:LYS:HD3	4:A:2137:HOH:O	2.21	0.41
1:A:155:VAL:HG21	1:A:198:ASN:HB2	2.03	0.40
1:A:69:PRO:HG2	1:A:203:LEU:HD11	2.04	0.40
1:A:435:LYS:HD2	1:A:435:LYS:HA	1.90	0.40
1:A:10:GLU:CB	1:A:48:ASN:HA	2.51	0.40
1:A:138:ILE:HD13	1:A:184:VAL:HG23	2.04	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	548/558 (98%)	524 (96%)	22 (4%)	2 (0%)	34   48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	LYS
1	A	62	ASP

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	487/494 (99%)	433 (89%)	54 (11%)	6   8

All (54) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	GLU
1	A	20	GLN
1	A	25	LEU
1	A	45	ILE
1	A	53	ARG
1	A	65	GLU
1	A	72	ARG
1	A	74	GLN
1	A	87	GLU
1	A	93	ASN
1	A	94	GLU
1	A	98	LYS
1	A	121	VAL
1	A	125	LEU
1	A	128	LEU
1	A	155	VAL
1	A	174	VAL
1	A	179	LYS
1	A	184	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	193	VAL
1	A	215	TRP
1	A	217	LEU
1	A	225	GLU
1	A	229	VAL
1	A	237	VAL
1	A	280	VAL
1	A	290	ARG
1	A	292	VAL
1	A	323	GLN
1	A	332	LEU
1	A	336	VAL
1	A	339	TYR
1	A	340	LYS
1	A	361	VAL
1	A	370	VAL
1	A	435	LYS
1	A	438	GLN
1	A	444	ARG
1	A	445	LYS
1	A	446	LYS
1	A	447	GLU
1	A	456	ASP
1	A	458	SER
1	A	483	MET
1	A	487	LEU
1	A	497	VAL
1	A	501	GLU
1	A	509	ARG
1	A	518	SER
1	A	525	LYS
1	A	531	LEU
1	A	544	GLU
1	A	553	PHE
1	A	555	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	GLN
1	A	93	ASN
1	A	143	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	245	ASN
1	A	264	HIS
1	A	301	ASN
1	A	311	HIS
1	A	323	GLN
1	A	357	HIS
1	A	450	GLN
1	A	499	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FLC	A	1001	-	3,12,12	0.70	0	3,17,17	0.60	0
3	GOL	A	1103	-	5,5,5	0.36	0	5,5,5	0.38	0
3	GOL	A	1101	-	5,5,5	0.36	0	5,5,5	0.29	0
3	GOL	A	1102	-	5,5,5	0.36	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	A	1001	-	-	0/6/16/16	-
3	GOL	A	1103	-	-	0/4/4/4	-
3	GOL	A	1101	-	-	1/4/4/4	-
3	GOL	A	1102	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1102	GOL	O2-C2-C3-O3
3	A	1101	GOL	C1-C2-C3-O3
3	A	1102	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1103	GOL	1	0
3	A	1101	GOL	2	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	550/558 (98%)	0.21	9 (1%) 72 69	24, 40, 78, 101	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	GLY	4.7
1	A	211	TYR	3.8
1	A	212	LYS	2.7
1	A	46	THR	2.6
1	A	217	LEU	2.6
1	A	9	ASN	2.4
1	A	33	TYR	2.2
1	A	446	LYS	2.1
1	A	52	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates i

There are no carbohydrates in this entry.

### 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	1101	6/6	0.82	0.33	79,80,81,81	0
3	GOL	A	1103	6/6	0.90	0.29	58,58,59,59	0
3	GOL	A	1102	6/6	0.94	0.38	65,66,66,67	0
2	FLC	A	1001	13/13	0.96	0.14	49,50,50,50	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.