



# wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 18, 2023 – 11:40 AM EST

PDB ID : 1VHA  
Title : Crystal structure of 2C-methyl-D-erythritol 2,4-cyclodiphosphate synthase  
Authors : Structural GenomiX  
Deposited on : 2003-12-01  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	 5% 67% 20% • 10%
1	B	170	 5% 66% 22% • 11%
1	C	170	 5% 71% 18% • 10%
1	D	170	 5% 64% 24% • 11%
1	E	170	 5% 65% 22% • 11%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	170	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACY	A	170	-	-	X	-
3	ACY	E	169	-	-	-	X
4	POP	B	170[C]	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	153	1160	730	209	217	2	2	0	1	0
1	B	152	1148	723	207	214	2	2	0	0	0
1	C	153	1157	729	209	215	2	2	0	0	0
1	D	152	1151	724	207	216	2	2	0	1	0
1	E	152	1148	723	207	214	2	2	0	0	0
1	F	152	1151	724	207	216	2	2	0	1	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	cloning artifact	UNP P44815
A	0	SER	-	cloning artifact	UNP P44815
A	1	LEU	-	cloning artifact	UNP P44815
A	67	MSE	MET	modified residue	UNP P44815
A	106	MSE	MET	modified residue	UNP P44815
A	113	MSE	MET	modified residue	UNP P44815
A	159	GLU	-	cloning artifact	UNP P44815
A	160	GLY	-	cloning artifact	UNP P44815
A	161	GLY	-	cloning artifact	UNP P44815
A	162	SER	-	cloning artifact	UNP P44815
A	163	HIS	-	cloning artifact	UNP P44815
A	164	HIS	-	cloning artifact	UNP P44815
A	165	HIS	-	cloning artifact	UNP P44815
A	166	HIS	-	cloning artifact	UNP P44815
A	167	HIS	-	cloning artifact	UNP P44815
A	168	HIS	-	cloning artifact	UNP P44815
B	-1	MSE	-	cloning artifact	UNP P44815

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	SER	-	cloning artifact	UNP P44815
B	1	LEU	-	cloning artifact	UNP P44815
B	67	MSE	MET	modified residue	UNP P44815
B	106	MSE	MET	modified residue	UNP P44815
B	113	MSE	MET	modified residue	UNP P44815
B	159	GLU	-	cloning artifact	UNP P44815
B	160	GLY	-	cloning artifact	UNP P44815
B	161	GLY	-	cloning artifact	UNP P44815
B	162	SER	-	cloning artifact	UNP P44815
B	163	HIS	-	cloning artifact	UNP P44815
B	164	HIS	-	cloning artifact	UNP P44815
B	165	HIS	-	cloning artifact	UNP P44815
B	166	HIS	-	cloning artifact	UNP P44815
B	167	HIS	-	cloning artifact	UNP P44815
B	168	HIS	-	cloning artifact	UNP P44815
C	-1	MSE	-	cloning artifact	UNP P44815
C	0	SER	-	cloning artifact	UNP P44815
C	1	LEU	-	cloning artifact	UNP P44815
C	67	MSE	MET	modified residue	UNP P44815
C	106	MSE	MET	modified residue	UNP P44815
C	113	MSE	MET	modified residue	UNP P44815
C	159	GLU	-	cloning artifact	UNP P44815
C	160	GLY	-	cloning artifact	UNP P44815
C	161	GLY	-	cloning artifact	UNP P44815
C	162	SER	-	cloning artifact	UNP P44815
C	163	HIS	-	cloning artifact	UNP P44815
C	164	HIS	-	cloning artifact	UNP P44815
C	165	HIS	-	cloning artifact	UNP P44815
C	166	HIS	-	cloning artifact	UNP P44815
C	167	HIS	-	cloning artifact	UNP P44815
C	168	HIS	-	cloning artifact	UNP P44815
D	-1	MSE	-	cloning artifact	UNP P44815
D	0	SER	-	cloning artifact	UNP P44815
D	1	LEU	-	cloning artifact	UNP P44815
D	67	MSE	MET	modified residue	UNP P44815
D	106	MSE	MET	modified residue	UNP P44815
D	113	MSE	MET	modified residue	UNP P44815
D	159	GLU	-	cloning artifact	UNP P44815
D	160	GLY	-	cloning artifact	UNP P44815
D	161	GLY	-	cloning artifact	UNP P44815
D	162	SER	-	cloning artifact	UNP P44815
D	163	HIS	-	cloning artifact	UNP P44815

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	164	HIS	-	cloning artifact	UNP P44815
D	165	HIS	-	cloning artifact	UNP P44815
D	166	HIS	-	cloning artifact	UNP P44815
D	167	HIS	-	cloning artifact	UNP P44815
D	168	HIS	-	cloning artifact	UNP P44815
E	-1	MSE	-	cloning artifact	UNP P44815
E	0	SER	-	cloning artifact	UNP P44815
E	1	LEU	-	cloning artifact	UNP P44815
E	67	MSE	MET	modified residue	UNP P44815
E	106	MSE	MET	modified residue	UNP P44815
E	113	MSE	MET	modified residue	UNP P44815
E	159	GLU	-	cloning artifact	UNP P44815
E	160	GLY	-	cloning artifact	UNP P44815
E	161	GLY	-	cloning artifact	UNP P44815
E	162	SER	-	cloning artifact	UNP P44815
E	163	HIS	-	cloning artifact	UNP P44815
E	164	HIS	-	cloning artifact	UNP P44815
E	165	HIS	-	cloning artifact	UNP P44815
E	166	HIS	-	cloning artifact	UNP P44815
E	167	HIS	-	cloning artifact	UNP P44815
E	168	HIS	-	cloning artifact	UNP P44815
F	-1	MSE	-	cloning artifact	UNP P44815
F	0	SER	-	cloning artifact	UNP P44815
F	1	LEU	-	cloning artifact	UNP P44815
F	67	MSE	MET	modified residue	UNP P44815
F	106	MSE	MET	modified residue	UNP P44815
F	113	MSE	MET	modified residue	UNP P44815
F	159	GLU	-	cloning artifact	UNP P44815
F	160	GLY	-	cloning artifact	UNP P44815
F	161	GLY	-	cloning artifact	UNP P44815
F	162	SER	-	cloning artifact	UNP P44815
F	163	HIS	-	cloning artifact	UNP P44815
F	164	HIS	-	cloning artifact	UNP P44815
F	165	HIS	-	cloning artifact	UNP P44815
F	166	HIS	-	cloning artifact	UNP P44815
F	167	HIS	-	cloning artifact	UNP P44815
F	168	HIS	-	cloning artifact	UNP P44815


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mn 1 1	0	0
2	B	1	Total Mn 1 1	0	0
2	C	1	Total Mn 1 1	0	0
2	D	1	Total Mn 1 1	0	0
2	F	1	Total Mn 1 1	0	0

- Molecule 3 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	E	1	Total C O 4 2 2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is PYROPHOSPHATE 2- (three-letter code: POP) (formula:  $\text{H}_2\text{O}_7\text{P}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	P	0	1
			27	21	6		
4	D	1	Total	O	P	0	1
			27	21	6		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	42	Total	O	0	0
			42	42		
5	B	39	Total	O	0	0
			39	39		
5	C	46	Total	O	0	0
			46	46		
5	D	33	Total	O	0	0
			33	33		
5	E	28	Total	O	0	0
			28	28		
5	F	38	Total	O	0	0
			38	38		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase



- Molecule 1: 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase




- Molecule 1: 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase




- Molecule 1: 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase






● Molecule 1: 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase



● Molecule 1: 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.72Å 104.72Å 195.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.55 – 2.35 45.55 – 2.35	Depositor EDS
% Data completeness (in resolution range)	(Not available) (45.55-2.35) 100.0 (45.55-2.35)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.44 (at 2.34Å)	Xtrriage
Refinement program	REFMAC 4.0	Depositor
R, $R_{free}$	0.244 , 0.288 0.213 , 0.258	Depositor DCC
$R_{free}$ test set	2211 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.7	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7224	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: POP, MN, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.36	0/1180	0.57	0/1589
1	B	0.36	0/1163	0.58	0/1567
1	C	0.36	0/1172	0.58	1/1578 (0.1%)
1	D	0.33	0/1171	0.55	0/1578
1	E	0.32	0/1163	0.55	0/1567
1	F	0.38	0/1171	0.58	0/1578
All	All	0.35	0/7020	0.57	1/9457 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	93	GLY	N-CA-C	-5.54	99.24	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1160	0	1167	45	0
1	B	1148	0	1154	37	0
1	C	1157	0	1167	35	0
1	D	1151	0	1154	31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1148	0	1154	33	0
1	F	1151	0	1154	38	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	F	1	0	0	0	0
3	A	4	0	3	2	0
3	B	4	0	3	1	0
3	C	4	0	3	1	0
3	D	4	0	3	1	0
3	E	4	0	3	0	0
3	F	4	0	3	1	0
4	B	27	0	0	7	0
4	D	27	0	0	5	0
5	A	42	0	0	0	0
5	B	39	0	0	0	0
5	C	46	0	0	0	0
5	D	33	0	0	0	0
5	E	28	0	0	1	0
5	F	38	0	0	0	0
All	All	7224	0	6968	203	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 203 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:2:ILE:HG23	1:F:154:LEU:HD11	1.43	0.99
1:A:70:LYS:HG2	1:A:71:ASN:H	1.31	0.92
1:A:40:VAL:HG13	1:A:149:CYS:HB2	1.50	0.92
1:A:15:GLU:HG2	1:A:30:THR:HB	1.52	0.91
1:B:107:ARG:HH21	1:B:107:ARG:HB2	1.34	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/170 (88%)	142 (95%)	7 (5%)	1 (1%)	22	23
1	B	148/170 (87%)	143 (97%)	4 (3%)	1 (1%)	22	23
1	C	149/170 (88%)	143 (96%)	4 (3%)	2 (1%)	12	10
1	D	149/170 (88%)	142 (95%)	5 (3%)	2 (1%)	12	10
1	E	148/170 (87%)	141 (95%)	6 (4%)	1 (1%)	22	23
1	F	149/170 (88%)	141 (95%)	6 (4%)	2 (1%)	12	10
All	All	893/1020 (88%)	852 (95%)	32 (4%)	9 (1%)	15	15

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	104	PRO
1	C	104	PRO
1	C	105	LYS
1	D	104	PRO
1	E	104	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/129 (92%)	114 (97%)	4 (3%)	37	46
1	B	116/129 (90%)	111 (96%)	5 (4%)	29	35
1	C	117/129 (91%)	116 (99%)	1 (1%)	78	87

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	117/129 (91%)	113 (97%)	4 (3%)	37	46
1	E	116/129 (90%)	113 (97%)	3 (3%)	46	56
1	F	117/129 (91%)	115 (98%)	2 (2%)	60	72
All	All	701/774 (91%)	682 (97%)	19 (3%)	44	55

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	3	ARG
1	F	35	HIS
1	F	131	LYS
1	E	134	THR
1	B	119	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	35	HIS
1	D	71	ASN
1	F	35	HIS
1	D	122	GLN
1	E	122	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 17 ligands modelled in this entry, 5 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	ACY	A	170	4	3,3,3	0.90	0	3,3,3	1.72	1 (33%)
3	ACY	E	169	4	3,3,3	0.93	0	3,3,3	1.76	1 (33%)
3	ACY	F	170	4	3,3,3	0.93	0	3,3,3	1.70	1 (33%)
4	POP	D	170[B]	3	6,8,8	1.16	0	13,13,13	0.81	0
3	ACY	C	170	4	3,3,3	1.03	0	3,3,3	1.75	1 (33%)
4	POP	B	170[C]	3	6,8,8	1.15	0	13,13,13	0.79	0
4	POP	B	170[B]	3	6,8,8	1.14	0	13,13,13	0.84	0
4	POP	D	170[A]	3,1	6,8,8	1.14	0	13,13,13	0.81	0
4	POP	D	170[C]	3	6,8,8	1.15	0	13,13,13	0.82	0
4	POP	B	170[A]	3	6,8,8	1.10	0	13,13,13	0.84	0
3	ACY	B	171	4	3,3,3	0.92	0	3,3,3	1.73	1 (33%)
3	ACY	D	171	4	3,3,3	0.88	0	3,3,3	1.71	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	POP	D	170[B]	3	-	3/6/6/6	-
4	POP	B	170[B]	3	-	3/6/6/6	-
4	POP	D	170[A]	3,1	-	2/6/6/6	-
4	POP	D	170[C]	3	-	3/6/6/6	-
4	POP	B	170[A]	3	-	1/6/6/6	-
4	POP	B	170[C]	3	-	1/6/6/6	-

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	169	ACY	O-C-CH3	-2.46	112.76	122.33
3	C	170	ACY	O-C-CH3	-2.44	112.84	122.33
3	B	171	ACY	O-C-CH3	-2.41	112.93	122.33
3	A	170	ACY	O-C-CH3	-2.39	113.02	122.33
3	D	171	ACY	O-C-CH3	-2.39	113.04	122.33

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	170[C]	POP	P1-O-P2-O5
4	B	170[B]	POP	P1-O-P2-O4
4	D	170[B]	POP	P1-O-P2-O4
4	D	170[C]	POP	P1-O-P2-O4
4	B	170[A]	POP	P1-O-P2-O5

There are no ring outliers.

10 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	170	ACY	2	0
3	F	170	ACY	1	0
4	D	170[B]	POP	2	0
3	C	170	ACY	1	0
4	B	170[C]	POP	4	0
4	B	170[B]	POP	1	0
4	D	170[A]	POP	3	0
4	B	170[A]	POP	2	0
3	B	171	ACY	1	0
3	D	171	ACY	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	151/170 (88%)	0.20	8 (5%) 26 38	21, 38, 64, 74	0
1	B	150/170 (88%)	0.29	9 (6%) 21 32	22, 33, 63, 95	0
1	C	151/170 (88%)	0.24	8 (5%) 26 38	18, 35, 61, 69	0
1	D	150/170 (88%)	0.19	9 (6%) 21 32	25, 44, 68, 88	0
1	E	150/170 (88%)	0.45	8 (5%) 26 38	25, 46, 72, 91	0
1	F	150/170 (88%)	0.08	6 (4%) 38 51	24, 37, 68, 95	0
All	All	902/1020 (88%)	0.24	48 (5%) 26 38	18, 38, 67, 95	0

The worst 5 of 48 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	62	PHE	9.4
1	B	63	PRO	7.8
1	F	63	PRO	6.7
1	B	71	ASN	6.0
1	E	62	PHE	5.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	ACY	E	169	4/4	0.64	0.44	35,37,39,40	4
3	ACY	D	171	4/4	0.73	0.32	32,32,33,35	4
3	ACY	F	170	4/4	0.73	0.30	31,32,32,33	4
3	ACY	A	170	4/4	0.75	0.35	34,34,36,37	4
3	ACY	B	171	4/4	0.81	0.27	25,27,28,28	4
2	MN	F	169	1/1	0.86	0.13	55,55,55,55	1
2	MN	D	169	1/1	0.88	0.16	62,62,62,62	1
2	MN	B	169	1/1	0.89	0.22	54,54,54,54	1
3	ACY	C	170	4/4	0.91	0.29	20,20,21,23	4
2	MN	A	169	1/1	0.92	0.20	52,52,52,52	1
2	MN	C	169	1/1	0.94	0.38	64,64,64,64	1
4	POP	B	170[A]	9/9	0.98	0.13	13,15,18,18	9
4	POP	B	170[B]	9/9	0.98	0.13	8,10,13,15	9
4	POP	B	170[C]	9/9	0.98	0.13	39,40,41,41	9
4	POP	D	170[A]	9/9	0.98	0.12	16,17,22,23	9
4	POP	D	170[B]	9/9	0.98	0.12	27,29,33,33	9
4	POP	D	170[C]	9/9	0.98	0.12	26,27,32,32	9

## 6.5 Other polymers [i](#)

There are no such residues in this entry.