

# wwPDB X-ray Structure Validation Summary Report (i)

#### Oct 5, 2023 – 01:29 AM EDT

PDB ID	:	6VHI
Title	:	Crystal structure of the human ILRUN Fw domain
Authors	:	Caputo, A.T.; Adams, T.E.
Deposited on		
Resolution	:	2.46  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

:	FAILED
:	1.13
:	FAILED
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	Engh & Huber $(2001)$
:	Parkinson et al. (1996)
:	2.35.1
	: : : :

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\hbox{-}RAY\,DIFFRACTION$ 

The reported resolution of this entry is 2.46 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



# 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1643 atoms, of which 803 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein ILRUN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	А	108	Total 1637	C 530	Н 803	N 135	O 160	S 9	818	0	0

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	6	Total O 6 6	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



# 3 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	135.93Å $40.98$ Å $23.29$ Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $91.43^{\circ}$ $90.00^{\circ}$	Depositor	
Resolution (Å)	67.94 - 2.46	Depositor	
% Data completeness	99.9 (67.94-2.46)	Depositor	
(in resolution range)		-	
$R_{merge}$	0.12	Depositor	
R <sub>sym</sub>	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$2.52 (at 2.45 \text{\AA})$	Xtriage	
Refinement program	BUSTER 2.10.3	Depositor	
$R, R_{free}$	0.192 , $0.222$	Depositor	
Wilson B-factor $(Å^2)$	35.9	Xtriage	
Anisotropy	1.216	Xtriage	
L-test for twinning <sup>2</sup>	$<  L  > = 0.48, < L^2 > = 0.32$	Xtriage	
Estimated twinning fraction	0.047 for -h,-k,l	Xtriage	
Total number of atoms	1643	wwPDB-VP	
Average B, all atoms $(Å^2)$	57.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.17% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 4 Model quality (i)

## 4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles (i)

#### 4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

#### 4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry (i)

There are no ligands in this entry.

#### 4.7 Other polymers (i)

There are no such residues in this entry.



# 4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 5 Fit of model and data (i)

# 5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

## 5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

## 5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

