



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2024 – 04:33 AM EDT

PDB ID : 6VJI  
Title : Structure of mammalian NEIL2 from Monodelphis domestica  
Authors : Eckenroth, B.E.; Doubleie, S.  
Deposited on : 2020-01-16  
Resolution : 2.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	 10% 64% 10% 26%
1	B	344	 13% 60% 13% 26%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nei like DNA glycosylase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	255	1912	1248	315	338	7	4	0	1	0
1	B	253	1895	1240	314	331	6	4	0	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	337	LEU	-	expression tag	UNP F7AMK3
A	338	GLU	-	expression tag	UNP F7AMK3
A	339	HIS	-	expression tag	UNP F7AMK3
A	340	HIS	-	expression tag	UNP F7AMK3
A	341	HIS	-	expression tag	UNP F7AMK3
A	342	HIS	-	expression tag	UNP F7AMK3
A	343	HIS	-	expression tag	UNP F7AMK3
A	344	HIS	-	expression tag	UNP F7AMK3
B	337	LEU	-	expression tag	UNP F7AMK3
B	338	GLU	-	expression tag	UNP F7AMK3
B	339	HIS	-	expression tag	UNP F7AMK3
B	340	HIS	-	expression tag	UNP F7AMK3
B	341	HIS	-	expression tag	UNP F7AMK3
B	342	HIS	-	expression tag	UNP F7AMK3
B	343	HIS	-	expression tag	UNP F7AMK3
B	344	HIS	-	expression tag	UNP F7AMK3

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	5	Total O 5 5	0	0
3	B	1	Total O 1 1	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nei like DNA glycosylase 2



- Molecule 1: Nei like DNA glycosylase 2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.89Å 67.89Å 149.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.96 – 2.54 37.96 – 2.54	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.96-2.54) 99.7 (37.96-2.54)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.66 (at 2.54Å)	Xtrriage
Refinement program	PHENIX 1.14-3260	Depositor
R, $R_{free}$	0.250 , 0.275 0.253 , 0.276	Depositor DCC
$R_{free}$ test set	2473 reflections (9.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	77.0	Xtrriage
Anisotropy	0.370	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 81.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.032 for -h,-k,l 0.054 for h,-h-k,-l 0.378 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3815	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	98.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/1966	0.43	0/2682
1	B	0.24	0/1949	0.44	0/2660
All	All	0.24	0/3915	0.43	0/5342

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1912	0	1797	19	0
1	B	1895	0	1784	25	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	1	0	0	0	0
All	All	3815	0	3581	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:VAL:HG21	1:A:246:ILE:HD11	1.74	0.70
1:A:246:ILE:O	1:A:322:GLN:NE2	2.25	0.67
1:A:133:LEU:HD11	1:B:70:LEU:HD11	1.81	0.63
1:B:237:LYS:HZ3	1:B:315:THR:HG22	1.64	0.62
1:B:246:ILE:O	1:B:322:GLN:NE2	2.34	0.61
1:A:294:CYS:SG	1:A:298:HIS:N	2.74	0.60
1:B:210:LEU:HD13	1:B:264:LEU:HD22	1.88	0.56
1:A:238:ASN:ND2	1:A:316:TRP:H	2.03	0.56
1:A:136:CYS:HB2	1:A:188:CYS:SG	2.48	0.53
1:B:196:THR:HB	1:B:205:ASP:HB2	1.90	0.53
1:A:70:LEU:HD11	1:B:133:LEU:HD11	1.91	0.52
1:B:246:ILE:HD11	1:B:259:ASN:HB3	1.89	0.52
1:B:294:CYS:SG	1:B:298:HIS:N	2.81	0.51
1:B:136:CYS:HB2	1:B:188:CYS:SG	2.52	0.49
1:A:294:CYS:SG	1:A:297:GLY:N	2.85	0.49
1:A:238:ASN:ND2	1:A:316:TRP:HE3	2.11	0.49
1:B:168:ARG:HG3	1:B:181:TYR:O	2.13	0.47
1:A:58:LEU:HB3	1:A:66:GLU:HG3	1.97	0.47
1:A:196:THR:HB	1:A:205:ASP:HB2	1.97	0.46
1:B:235:ILE:HG12	1:B:289:TYR:CZ	2.51	0.45
1:A:206:ARG:NH1	1:A:265:ASP:OD1	2.42	0.45
1:B:5:PRO:HG3	1:B:140:GLY:O	2.17	0.45
1:B:250:SER:OG	1:B:324:LYS:O	2.32	0.45
1:A:10:PHE:CD1	1:A:145:VAL:HG12	2.52	0.45
1:B:28:ASN:OD1	1:B:168:ARG:HA	2.18	0.44
1:B:41:ARG:O	1:B:56:PHE:HA	2.17	0.44
1:B:237:LYS:NZ	1:B:315:THR:HG22	2.31	0.44
1:A:37:LEU:HA	1:A:40:LEU:HD12	2.00	0.43
1:B:276:GLN:NE2	1:B:280:GLU:OE2	2.51	0.43
1:B:294:CYS:SG	1:B:297:GLY:N	2.90	0.43
1:A:28:ASN:ND2	1:A:168:ARG:HA	2.33	0.43
1:B:247:HIS:HB2	1:B:323:PRO:O	2.18	0.43
1:A:242:TYR:CD1	1:A:288:ILE:HD12	2.53	0.43
1:B:180:PHE:HB3	1:B:183:CYS:SG	2.59	0.43
1:B:213:LEU:HD21	1:B:263:LEU:HD23	2.01	0.42
1:A:41:ARG:O	1:A:56:PHE:HA	2.20	0.42
1:B:37:LEU:HA	1:B:40:LEU:HD12	2.02	0.42
1:B:303:ASP:OD1	1:B:304:SER:N	2.50	0.41
1:B:10:PHE:CD1	1:B:145:VAL:HG12	2.55	0.41
1:A:28:ASN:HD21	1:A:168:ARG:HA	1.86	0.41
1:B:290:GLN:HG2	1:B:316:TRP:CZ2	2.56	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:ALA:O	1:A:326:GLU:C	2.60	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/344 (72%)	242 (98%)	6 (2%)	0	100	100
1	B	246/344 (72%)	240 (98%)	6 (2%)	0	100	100
All	All	494/688 (72%)	482 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	195/300 (65%)	188 (96%)	7 (4%)	35	47
1	B	192/300 (64%)	188 (98%)	4 (2%)	53	68
All	All	387/600 (64%)	376 (97%)	11 (3%)	42	58

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	68	PHE
1	A	146	ARG
1	A	246	ILE
1	A	284	LEU
1	A	311	PHE
1	A	312	GLN
1	B	68	PHE
1	B	210	LEU
1	B	284	LEU
1	B	311	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	ASN
1	A	276	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	252/344 (73%)	0.97	33 (13%) <b>3</b> <b>4</b>	57, 98, 138, 158	0
1	B	250/344 (72%)	1.06	45 (18%) <b>1</b> <b>1</b>	59, 98, 137, 168	2 (0%)
All	All	502/688 (72%)	1.02	78 (15%) <b>2</b> <b>2</b>	57, 98, 138, 168	2 (0%)

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	141	LEU	6.2
1	A	212	ALA	5.4
1	A	252	GLY	5.2
1	B	69	LEU	4.7
1	A	283	PRO	4.3
1	B	325	ALA	4.3
1	B	70	LEU	4.2
1	B	131	SER	4.1
1	B	263	LEU	3.9
1	B	320	HIS	3.7
1	A	228	TYR	3.7
1	A	314	LEU	3.7
1	B	142	PHE	3.6
1	A	305	PHE	3.6
1	A	139	PHE	3.6
1	B	132	GLY	3.6
1	A	142	PHE	3.5
1	A	209	ALA	3.5
1	A	249	LEU	3.4
1	A	140	GLY	3.4
1	B	64	LEU	3.3
1	B	236	ILE	3.3
1	B	288	ILE	3.2
1	B	67	SER	3.2

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	68	PHE	3.2
1	B	229	PHE	3.1
1	B	219	VAL	3.1
1	A	254	CYS	3.1
1	A	70	LEU	3.1
1	A	288	ILE	3.0
1	B	71	PRO	3.0
1	A	135	LEU	2.9
1	B	242	TYR	2.9
1	B	249	LEU	2.8
1	B	182	ASN	2.8
1	A	255	LEU	2.8
1	B	245	ARG	2.8
1	A	287	LEU	2.7
1	A	311	PHE	2.7
1	A	284	LEU	2.7
1	B	47	VAL	2.6
1	B	18	VAL	2.6
1	B	250	SER	2.5
1	A	64	LEU	2.5
1	B	240	VAL	2.5
1	B	258	LEU	2.5
1	B	213	LEU	2.4
1	B	228	TYR	2.4
1	B	246	ILE	2.4
1	A	263	LEU	2.4
1	B	62	LEU	2.4
1	A	236	ILE	2.4
1	B	32	ILE	2.4
1	B	255	LEU	2.4
1	B	204	PHE	2.4
1	B	313	ARG	2.4
1	A	200	LEU	2.4
1	B	199	ILE	2.3
1	A	47	VAL	2.3
1	B	306	GLY	2.3
1	A	185	ILE	2.3
1	A	240	VAL	2.2
1	A	260	LEU	2.2
1	B	10	PHE	2.2
1	B	65	PRO	2.2
1	B	264	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	133	LEU	2.2
1	A	264	LEU	2.2
1	B	193	VAL	2.2
1	B	15	ALA	2.2
1	A	224	LEU	2.1
1	B	232	LEU	2.1
1	B	31	LYS	2.1
1	A	52	LEU	2.1
1	A	307	PRO	2.0
1	A	134	TRP	2.0
1	B	137	PHE	2.0
1	B	181	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	B	401	1/1	0.83	0.19	119,119,119,119	0
2	ZN	A	401	1/1	0.91	0.21	123,123,123,123	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.