

wwPDB X-ray Structure Validation Summary Report (i)

Sep 22, 2020 – 06:21 PM EDT

PDB ID	:	6VLT
Title	:	Crystal Structure of Human P450 2C9*2 Genetic Variant in Complex with
		Losartan
Authors	:	Shah, M.B.
Deposited on	:	2020-01-25
Resolution	:	3.12 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.14.6
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.6

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution
Wiethic	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	1292 (3.14-3.10)
Clashscore	141614	1389 (3.14-3.10)
Ramachandran outliers	138981	1337 (3.14-3.10)
Sidechain outliers	138945	1337 (3.14-3.10)
RSRZ outliers	127900	1260 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	476	70%	25%	• •
1	В	476	75%	19%	•••
1	С	476	77%	16%	•••
1	D	476	77%	17%	••
1	Е	476	79%	16%	••



Mol	Chain	Length	Quality of chain		
1	F	476	% 72%	22%	•••
1	G	476	% 74%	21%	
1	Н	476	73%	22%	•••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CM5	Е	501	-	-	-	Х
4	CM5	Е	504	-	-	-	Х
4	CM5	F	501	-	-	-	Х
4	CM5	G	501	-	-	-	Х
4	CM5	Н	501	-	-	-	Х



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 29782 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	461	Total	С	Ν	0	\mathbf{S}	0	0	0
1	A	401	3615	2337	599	656	23	0	0	0
1	В	461	Total	С	Ν	0	S	0	0	0
1	D	401	3639	2348	605	662	24	0	0	0
1	С	460	Total	С	Ν	0	S	0	0	0
1		400	3619	2340	601	655	23	0	0	0
1	Л	461	Total	С	Ν	0	S	0	0	0
1	D	401	3586	2316	594	652	24	0	0	0
1	F	460	Total	С	Ν	0	S	0	0	0
1	Ľ	400	3575	2310	588	654	23	0	0	0
1	Б	460	Total	С	Ν	0	S	0	0	0
1	Г	400	3583	2318	592	650	23	0	0	0
1	С	462	Total	С	Ν	0	S	0	0	0
1	G	402	3580	2315	588	653	24	0	0	0
1	ц	450	Total	С	Ν	0	S	0	0	0
	п	409	3571	2311	586	650	24	0	0	U

• Molecule 1 is a protein called Cytochrome P450 2C9.

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	19	MET	-	initiating methionine	UNP P11712
А	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
А	21	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
А	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
А	23	THR	-	SEE SEQUENCE DETAILS	UNP P11712
А	144	CYS	ARG	engineered mutation	UNP P11712
А	490	ILE	VAL	engineered mutation	UNP P11712
А	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
А	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
А	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
А	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
В	19	MET	-	initiating methionine	UNP P11712
В	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712



Chain	Residue	Modelled	Actual	Comment	Reference
В	21	LYS	_	SEE SEQUENCE DETAILS	UNP P11712
В	22	LYS	_	SEE SEQUENCE DETAILS	UNP P11712
В	23	THR	_	SEE SEQUENCE DETAILS	UNP P11712
В	144	CYS	ARG	engineered mutation	UNP P11712
В	490	ILE	VAL	engineered mutation	UNP P11712
В	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
В	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
В	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
В	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
С	19	MET	-	initiating methionine	UNP P11712
С	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
С	21	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
С	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
С	23	THR	-	SEE SEQUENCE DETAILS	UNP P11712
С	144	CYS	ARG	engineered mutation	UNP P11712
С	490	ILE	VAL	engineered mutation	UNP P11712
С	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
С	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
С	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
С	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
D	19	MET	-	initiating methionine	UNP P11712
D	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
D	21	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
D	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
D	23	THR	-	SEE SEQUENCE DETAILS	UNP P11712
D	144	CYS	ARG	engineered mutation	UNP P11712
D	490	ILE	VAL	engineered mutation	UNP P11712
D	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
D	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
D	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
D	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
E	19	MET	-	initiating methionine	UNP P11712
E	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
E	21	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
E	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
E	23	THR	-	SEE SEQUENCE DETAILS	UNP P11712
E	144	CYS	ARG	engineered mutation	UNP P11712
E	490	ILE	VAL	engineered mutation	UNP P11712
E	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
E	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
E	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
E	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712



	6	V	Ι	Л	
--	---	---	---	---	--

Chain	Residue	Modelled	Actual	Comment	Reference
F	19	MET	-	initiating methionine	UNP P11712
F	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
F	21	LYS	_	SEE SEQUENCE DETAILS	UNP P11712
F	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
F	23	THR	_	SEE SEQUENCE DETAILS	UNP P11712
F	144	CYS	ARG	engineered mutation	UNP P11712
F	490	ILE	VAL	engineered mutation	UNP P11712
F	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
F	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
F	493	HIS	_	SEE SEQUENCE DETAILS	UNP P11712
F	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
G	19	MET	_	initiating methionine	UNP P11712
G	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
G	21	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
G	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
G	23	THR	_	SEE SEQUENCE DETAILS	UNP P11712
G	144	CYS	ARG	engineered mutation	UNP P11712
G	490	ILE	VAL	engineered mutation	UNP P11712
G	491	HIS	_	SEE SEQUENCE DETAILS	UNP P11712
G	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
G	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
G	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
Н	19	MET	-	initiating methionine	UNP P11712
Н	20	ALA	-	SEE SEQUENCE DETAILS	UNP P11712
Н	21	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
Н	22	LYS	-	SEE SEQUENCE DETAILS	UNP P11712
Н	23	THR	-	SEE SEQUENCE DETAILS	UNP P11712
Н	144	CYS	ARG	engineered mutation	UNP P11712
Н	490	ILE	VAL	engineered mutation	UNP P11712
Н	491	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
Н	492	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
Н	493	HIS	-	SEE SEQUENCE DETAILS	UNP P11712
Н	494	HIS	-	SEE SEQUENCE DETAILS	UNP P11712

• Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf		
0	Λ	1	Total	С	Fe	Ν	Ο	0	0		
	A	1	43	34	1	4	4	0	0		
0	В	1	Total	С	Fe	Ν	0	0	0		
	D	1	43	34	1	4	4	0	0		
0	C	1	Total	С	Fe	Ν	0	0	0		
	U	1	43	34	1	4	4	0	0		
0	л	1	Total	С	Fe	Ν	0	0	0		
	D	D			43	34	1	4	4	0	0
9	F	1	Total	С	Fe	Ν	0	0	0		
	Ľ	1	43	34	1	4	4	0	0		
9	F	1	Total	С	Fe	Ν	0	0	0		
	Ľ	1	43	34	1	4	4	0	0		
9	С	1	Total	С	Fe	Ν	0	0	0		
	G		43	34	1	4	4		0		
9	Ц	1	Total	С	Fe	Ν	Ο	0	0		
	11	L	43	34	1	4	4		U		

• Molecule 3 is [2-butyl-5-chloranyl-3-[[4-[2-(2H-1,2,3,4-tetrazol-5-yl)phenyl]phenyl]methyl]im idazol-4-yl]methanol (three-letter code: LSN) (formula: $C_{22}H_{23}ClN_6O$) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
9	٨	1	Total	С	Cl	Ν	0	0	0	
5	A	1	30	22	1	6	1	0	0	
9	D	1	Total	С	Cl	Ν	Ο	0	0	
5	D	1	30	22	1	6	1	0	0	
9	С	1	Total	С	Cl	Ν	Ο	0	0	
5	C	1	30	22	1	6	1	0	0	
2	Л	1	Total	С	Cl	Ν	0	0	0	
0	D	1	30	22	1	6	1	0	U	
9	Б	1	Total	С	Cl	Ν	Ο	0	0	
3	E	1	30	22	1	6	1	0	0	
2	Б	1	Total	С	Cl	Ν	Ο	0	0	
0	Г	1	30	22	1	6	1	0	0	
2	С	1	Total	С	Cl	Ν	Ο	0	0	
3	G		30	22	1	6	1	0		
2	ц	1	Total	С	Cl	Ν	Ο	0	0	
່ ^ວ	11		30	22	1	6	1	0		

• Molecule 4 is 5-CYCLOHEXYL-1-PENTYL-BETA-D-MALTOSIDE (three-letter code: CM5) (formula: $C_{23}H_{42}O_{11}$).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	F	1	Total C	0	0	0	
4	Ľ	1	34 23	11	0	0	
4	F	1	Total C	0	0	0	
4	Ľ	1	34 23	11	0	0	
4	Б	1	Total C	0	0	0	
4	Г	1	34 23	11	0	0	
4	Б	1	Total C	0	0	0	
4	Г	1	34 23	11	0	0	
4	С	1	Total C	0	0	0	
4	G	1	34 23	11	0	0	
4	С	1	Total C	0	0	0	
4	G	1	34 23	11	0	0	
4	Ц	1	Total C	0	0	0	
4		L	34 23	11	U	U	
4	ц	1	Total C	0	0	0	
4	11	L	34 23	11	U	U	





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	Е	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
5	F	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
5	G	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	12	Total O 12 12	0	0
6	В	19	Total O 19 19	0	0
6	С	21	TotalO2121	0	0
6	D	8	Total O 8 8	0	0
6	Е	22	TotalO2222	0	0
6	F	20	TotalO2020	0	0
6	G	16	Total O 16 16	0	0
6	Н	25	TotalO2525	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Cytochrome P450 2C9







• Molecule 1: Cytochrome P450 2C9





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	238.10Å 238.10Å 109.85Å	Deneriten
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
$\mathbf{P}_{\text{oscolution}}(\hat{\mathbf{A}})$	50.01 - 3.12	Depositor
Resolution (A)	39.61 - 3.12	EDS
% Data completeness	99.2 (50.01-3.12)	Depositor
(in resolution range)	99.3 (39.61 - 3.12)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.23 (at 3.12 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
B B.	0.156 , 0.221	Depositor
It, It _{free}	0.181 , 0.221	DCC
R_{free} test set	6047 reflections $(4.93%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	80.9	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.34 , 60.2	EDS
L-test for twinning ²	$< L > = 0.49, < L^2 > = 0.32$	Xtriage
	0.021 for -h,-k,l	
Estimated twinning fraction	0.457 for h,-h-k,-l	Xtriage
	0.022 for -k,-h,-l	
F_o, F_c correlation	0.95	EDS
Total number of atoms	29782	wwPDB-VP
Average B, all atoms $(Å^2)$	80.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.10% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LSN, HEM, CM5, SO4 $\,$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain Bond lengths		Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.62	0/3704	0.89	0/5028
1	В	0.63	0/3728	0.86	0/5057
1	С	0.63	0/3709	0.87	0/5034
1	D	0.62	0/3674	0.86	0/4995
1	Е	0.63	0/3663	0.86	0/4981
1	F	0.62	0/3672	0.84	0/4991
1	G	0.62	0/3668	0.83	0/4988
1	H	0.63	0/3659	0.86	0/4973
All	All	0.62	0/29477	0.86	0/40047

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3615	0	3577	123	0
1	В	3639	0	3613	81	0
1	С	3619	0	3588	92	0
1	D	3586	0	3514	79	0
1	Е	3575	0	3498	74	0



6VLT	
------	--

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	3583	0	3519	93	0
1	G	3580	0	3495	103	0
1	Н	3571	0	3502	92	0
2	А	43	0	30	12	0
2	В	43	0	30	10	0
2	С	43	0	30	14	0
2	D	43	0	30	11	0
2	Е	43	0	30	12	0
2	F	43	0	30	7	0
2	G	43	0	30	8	0
2	Н	43	0	30	5	0
3	А	30	0	0	5	0
3	В	30	0	0	1	0
3	С	30	0	0	0	0
3	D	30	0	0	5	0
3	Е	30	0	0	2	0
3	F	30	0	0	2	0
3	G	30	0	0	2	0
3	Н	30	0	0	2	0
4	Е	68	0	84	15	0
4	F	68	0	84	7	0
4	G	68	0	84	9	0
4	Н	68	0	84	16	0
5	Е	5	0	0	0	0
5	F	5	0	0	0	0
5	G	5	0	0	0	0
6	А	12	0	0	1	0
6	В	19	0	0	0	0
6	С	21	0	0	1	0
6	D	8	0	0	0	0
6	Е	22	0	0	0	0
6	F	20	0	0	1	0
6	G	16	0	0	1	0
6	Н	25	0	0	2	0
All	All	29782	0	28882	811	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 811 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:69:PHE:CD2	1:G:74:ILE:HD12	1.53	1.42
1:F:457:ASN:OD1	1:F:490:ILE:HD12	1.29	1.27
1:C:163:PRO:CB	1:C:461:LEU:HD11	1.66	1.24
1:D:277:ASN:O	1:D:279:PRO:HD2	1.39	1.19
1:E:461:LEU:HD12	1:E:461:LEU:O	1.42	1.18

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	P	\mathbf{erc}	entiles
1	А	457/476~(96%)	386 (84%)	62 (14%)	9 (2%)		7	30
1	В	457/476~(96%)	396 (87%)	52 (11%)	9 (2%)		7	30
1	С	456/476~(96%)	409 (90%)	40 (9%)	7 (2%)		10	38
1	D	457/476~(96%)	403 (88%)	42 (9%)	12 (3%)		5	25
1	Е	456/476~(96%)	397 (87%)	44 (10%)	15 (3%)		4	20
1	F	456/476~(96%)	391 (86%)	53 (12%)	12 (3%)		5	25
1	G	458/476~(96%)	398 (87%)	50 (11%)	10 (2%)		6	28
1	Н	455/476~(96%)	396 (87%)	46 (10%)	13 (3%)		4	23
All	All	3652/3808~(96%)	3176 (87%)	389 (11%)	87 (2%)		6	26

5 of 87 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	278	GLN
1	А	280	SER
1	В	255	MET
1	В	278	GLN
1	С	278	GLN



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	400/429~(93%)	383~(96%)	17~(4%)	29 61
1	В	406/429~(95%)	386~(95%)	20~(5%)	25 57
1	С	402/429~(94%)	385~(96%)	17 (4%)	30 62
1	D	393/429~(92%)	378~(96%)	15 (4%)	33 65
1	Ε	392/429~(91%)	387~(99%)	5 (1%)	69 86
1	F	393/429~(92%)	385~(98%)	8 (2%)	55 79
1	G	390/429~(91%)	378~(97%)	12 (3%)	40 69
1	Н	391/429~(91%)	383 (98%)	8 (2%)	55 79
All	All	3167/3432~(92%)	3065 (97%)	102 (3%)	39 69

 $5~{\rm of}~102$ residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	С	250	GLU
1	D	134	PHE
1	Н	34	THR
1	С	254	SER
1	С	322	LYS

Some side chains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such side chains are listed below:

Mol	Chain	Res	Type
1	F	231	ASN
1	Н	41	ASN
1	G	252	GLN
1	А	418	ASN
1	G	41	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tiple	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
	Type	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	F	502	1	27,50,50	1.91	5 (18%)	17,82,82	1.40	4 (23%)
2	HEM	А	501	1	27,50,50	1.89	6 (22%)	17,82,82	1.49	2 (11%)
4	CM5	F	504	-	36,36,36	0.14	0	49,49,49	0.21	0
4	CM5	Е	504	-	36,36,36	0.14	0	49,49,49	0.20	0
3	LSN	F	503	-	28,33,33	0.62	0	34,45,45	1.33	3 (8%)
3	LSN	D	502	-	28,33,33	0.62	0	34,45,45	1.32	3 (8%)
2	HEM	Н	502	1	27,50,50	1.19	2 (7%)	17,82,82	1.28	2 (11%)
4	CM5	F	501	-	36,36,36	0.14	0	49,49,49	0.21	0
4	CM5	Н	504	-	36,36,36	0.13	0	49,49,49	0.20	0
5	SO4	Е	505	-	4,4,4	0.43	0	6,6,6	0.04	0
4	CM5	G	501	-	36,36,36	0.14	0	49,49,49	0.23	0
2	HEM	G	502	1	27,50,50	1.84	6 (22%)	17,82,82	1.06	1 (5%)
3	LSN	А	502	-	28,33,33	0.61	0	$34,\!45,\!45$	1.32	3 (8%)
2	HEM	В	501	1	27,50,50	1.77	4 (14%)	17,82,82	1.08	1 (5%)
4	CM5	Н	501	-	36,36,36	0.14	0	49,49,49	0.22	0
5	SO4	G	505	-	4,4,4	0.41	0	6,6,6	0.06	0
5	SO4	F	505	-	4,4,4	0.45	0	6,6,6	0.06	0
2	HEM	С	501	1	27,50,50	1.88	5 (18%)	17,82,82	1.24	1 (5%)
4	CM5	G	504	-	36,36,36	0.13	0	49,49,49	0.20	0
2	HEM	D	501	1	27,50,50	1.89	7 (25%)	17,82,82	1.58	2 (11%)



Mal	Mol Type Chain		Dec	Tink	Bo	ond leng	$_{\rm ths}$	B	ond ang	les	
WIOI	туре	Unam	rtes	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	LSN	В	502	-	28,33,33	0.61	0	34,45,45	1.32	3 (8%)	
2	HEM	Е	502	1	$27,\!50,\!50$	1.83	8 (29%)	17,82,82	1.40	1 (5%)	
3	LSN	Н	503	-	28,33,33	0.61	0	34,45,45	1.34	3 (8%)	
3	LSN	Е	503	-	28,33,33	0.61	0	34,45,45	1.33	3 (8%)	
3	LSN	G	503	-	28,33,33	0.61	0	34,45,45	1.33	3 (8%)	
3	LSN	С	502	-	28,33,33	0.61	0	34,45,45	1.32	3 (8%)	
4	CM5	Е	501	-	36,36,36	0.14	0	49,49,49	0.22	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	F	502	1	-	2/6/54/54	-
2	HEM	А	501	1	-	4/6/54/54	-
4	CM5	F	504	-	-	10/17/65/65	0/3/3/3
4	CM5	Е	504	-	-	13/17/65/65	0/3/3/3
3	LSN	F	503	-	-	5/16/18/18	0/4/4/4
3	LSN	D	502	-	-	4/16/18/18	0/4/4/4
2	HEM	Н	502	1	-	0/6/54/54	-
4	CM5	F	501	-	-	12/17/65/65	0/3/3/3
4	CM5	Н	504	-	-	11/17/65/65	0/3/3/3
4	CM5	G	501	-	-	11/17/65/65	0/3/3/3
2	HEM	G	502	1	-	2/6/54/54	-
3	LSN	А	502	-	-	6/16/18/18	0/4/4/4
2	HEM	В	501	1	-	0/6/54/54	-
4	CM5	Н	501	-	-	8/17/65/65	0/3/3/3
2	HEM	С	501	1	-	2/6/54/54	-
4	CM5	G	504	-	-	9/17/65/65	0/3/3/3
2	HEM	D	501	1	-	0/6/54/54	-
3	LSN	В	502	-	-	5/16/18/18	0/4/4/4
2	HEM	Е	502	1	-	1/6/54/54	-
3	LSN	Н	503	-	-	6/16/18/18	0/4/4/4
3	LSN	Е	503	-	-	6/16/18/18	0/4/4/4
3	LSN	G	503	-	-	5/16/18/18	0/4/4/4
3	LSN	С	502	-	-	7/16/18/18	0/4/4/4



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CM5	Е	501	-	-	10/17/65/65	0/3/3/3

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	С	501	HEM	C3B-C2B	-5.43	1.32	1.40
2	F	502	HEM	C3B-C2B	-5.33	1.33	1.40
2	G	502	HEM	C3B-C2B	-5.00	1.33	1.40
2	В	501	HEM	C3B-C2B	-4.97	1.33	1.40
2	Е	502	HEM	C3B-C2B	-4.97	1.33	1.40

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	Е	502	HEM	CAA-CBA-CGA	-4.46	105.18	112.67
3	G	503	LSN	N4-N3-N2	4.29	112.34	109.53
2	D	501	HEM	CAA-CBA-CGA	-4.29	105.47	112.67
3	D	502	LSN	N4-N3-N2	4.29	112.34	109.53
3	Е	503	LSN	N4-N3-N2	4.29	112.34	109.53

There are no chirality outliers.

5 of 139 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	502	HEM	C2D-C3D-CAD-CBD
2	F	502	HEM	C4D-C3D-CAD-CBD
2	А	501	HEM	C1A-C2A-CAA-CBA
2	А	501	HEM	C3A-C2A-CAA-CBA
4	F	504	CM5	C2-C1-O12-C13

There are no ring outliers.

23 monomers are involved in 145 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	502	HEM	7	0
2	А	501	HEM	12	0
4	F	504	CM5	3	0
4	Е	504	CM5	5	0
3	F	503	LSN	2	0
3	D	502	LSN	5	0
2	Н	502	HEM	5	0



G	٦7	Т	Π	
υ	V	Т		L

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	501	CM5	4	0
4	Н	504	CM5	6	0
4	G	501	CM5	5	0
2	G	502	HEM	8	0
3	А	502	LSN	5	0
2	В	501	HEM	10	0
4	Н	501	CM5	10	0
2	С	501	HEM	14	0
4	G	504	CM5	4	0
2	D	501	HEM	11	0
3	В	502	LSN	1	0
2	Е	502	HEM	12	0
3	Н	503	LSN	2	0
3	Е	503	LSN	2	0
3	G	503	LSN	2	0
4	Е	501	CM5	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









































































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	461/476~(96%)	0.16	0 100 100	26, 73, 107, 152	0
1	В	461/476~(96%)	0.13	1 (0%) 95 91	26, 74, 105, 131	0
1	С	460/476~(96%)	0.14	1 (0%) 95 91	26, 75, 103, 136	0
1	D	461/476~(96%)	0.15	0 100 100	26, 73, 107, 148	0
1	Ε	460/476~(96%)	0.17	0 100 100	26, 81, 116, 158	0
1	F	460/476~(96%)	0.20	3 (0%) 87 77	26, 81, 118, 184	0
1	G	462/476~(97%)	0.23	4 (0%) 84 71	26, 83, 117, 182	0
1	Η	459/476~(96%)	0.17	0 100 100	26, 80, 115, 173	0
All	All	3684/3808~(96%)	0.17	9 (0%) 95 91	26, 77, 113, 184	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	240	MET	2.8
1	G	244	ILE	2.8
1	G	134	PHE	2.6
1	F	244	ILE	2.6
1	G	136	MET	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
5	SO4	G	505	5/5	0.62	0.19	169,180,184,187	0
5	SO4	F	505	5/5	0.69	0.19	174,177,187,193	0
4	CM5	G	501	34/34	0.72	0.43	77,138,158,165	0
4	CM5	Н	501	34/34	0.73	0.41	78,128,150,157	0
4	CM5	Е	501	34/34	0.76	0.48	83,130,155,167	0
4	CM5	Е	504	34/34	0.77	0.46	90,121,138,147	0
4	CM5	F	501	34/34	0.78	0.44	78,134,161,164	0
5	SO4	Е	505	5/5	0.80	0.13	163,167,183,183	0
4	CM5	Н	504	34/34	0.83	0.36	80,121,150,161	0
4	CM5	F	504	34/34	0.86	0.38	79,120,148,153	0
4	CM5	G	504	34/34	0.88	0.32	88,115,124,130	0
3	LSN	С	502	30/30	0.92	0.51	69,88,117,122	0
3	LSN	F	503	30/30	0.94	0.43	73,88,101,111	0
3	LSN	В	502	30/30	0.94	0.47	69,93,109,111	0
3	LSN	D	502	30/30	0.94	0.42	71,88,109,111	0
3	LSN	А	502	30/30	0.94	0.40	65,86,112,116	0
3	LSN	Н	503	30/30	0.95	0.43	70,79,108,111	0
3	LSN	G	503	30/30	0.95	0.42	77,94,110,114	0
3	LSN	Е	503	30/30	0.96	0.42	67,76,112,118	0
2	HEM	D	501	43/43	0.99	0.29	51,55,61,69	0
2	HEM	В	501	43/43	0.99	0.27	42,49,60,67	0
2	HEM	Е	502	43/43	0.99	0.26	42,48,62,67	0
2	HEM	Н	502	43/43	0.99	0.29	45,52,63,73	0
2	HEM	F	502	43/43	0.99	0.27	42,55,66,80	0
2	HEM	G	502	43/43	0.99	0.26	45,54,63,73	0
2	HEM	С	501	43/43	0.99	0.27	40,48,59,67	0
2	HEM	A	501	43/43	0.99	0.27	47,52,62,76	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

































































6.5 Other polymers (i)

There are no such residues in this entry.

