



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 07:29 pm BST

PDB ID : 3VND
Title : Crystal structure of tryptophan synthase alpha-subunit from the psychrophile *Shewanella frigidimarina* K14-2
Authors : Mitsuya, D.; Tanaka, S.; Matsumura, H.; Takano, K.; Urano, N.; Ishida, M.
Deposited on : 2012-01-12
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 69% 22% 6% .
1	B	267	 63% 28% 5% . .
1	C	267	 67% 24% 5% .
1	D	267	 64% 26% 6% .
1	E	267	 66% 25% 5% . .
1	F	267	 63% 27% 6% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	267	
1	H	267	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	H	302	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan synthase alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	1907	1220	313	369	5	0	0	0
1	B	259	1907	1220	313	369	5	0	0	0
1	C	259	1907	1220	313	369	5	0	0	0
1	D	259	1907	1220	313	369	5	0	0	0
1	E	259	1907	1220	313	369	5	0	0	0
1	F	259	1907	1220	313	369	5	0	0	0
1	G	259	1907	1220	313	369	5	0	0	0
1	H	259	1907	1220	313	369	5	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	E	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	G	1	Total	O	S	0	0
			5	4	1		
2	G	1	Total	O	S	0	0
			5	4	1		
2	H	1	Total	O	S	0	0
			5	4	1		
2	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is 3,6,9,12,15,18,21-HEPTAOXATRICOSANE-1,23-DIOL (three-letter code: PE8) (formula: C₁₆H₃₄O₉).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			25	16	9		
3	D	1	Total	C	O	0	0
			25	16	9		
3	E	1	Total	C	O	0	0
			25	16	9		
3	G	1	Total	C	O	0	0
			25	16	9		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	145	Total	O	0	0
			145	145		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	153	Total 153	O 153	0	0
4	C	156	Total 156	O 156	0	0
4	D	144	Total 144	O 144	0	0
4	E	147	Total 147	O 147	0	0
4	F	150	Total 150	O 150	0	0
4	G	149	Total 149	O 149	0	0
4	H	143	Total 143	O 143	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tryptophan synthase alpha chain



- Molecule 1: Tryptophan synthase alpha chain




- Molecule 1: Tryptophan synthase alpha chain







• Molecule 1: Tryptophan synthase alpha chain




• Molecule 1: Tryptophan synthase alpha chain




• Molecule 1: Tryptophan synthase alpha chain



• Molecule 1: Tryptophan synthase alpha chain





- Molecule 1: Tryptophan synthase alpha chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	181.14Å 182.78Å 181.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.79 – 2.60 48.74 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.79-2.60) 96.8 (48.74-2.60)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.31 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.173 , 0.211 0.174 , 0.206	Depositor DCC
R_{free} test set	4603 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtrriage
Anisotropy	0.168	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 16.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.208 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16633	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.46 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1573e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PE8, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	0/1945	1.03	6/2647 (0.2%)
1	B	0.91	0/1945	1.07	13/2647 (0.5%)
1	C	0.98	1/1945 (0.1%)	1.03	4/2647 (0.2%)
1	D	1.00	1/1945 (0.1%)	1.13	5/2647 (0.2%)
1	E	0.94	0/1945	1.05	7/2647 (0.3%)
1	F	0.93	1/1945 (0.1%)	1.04	8/2647 (0.3%)
1	G	0.94	0/1945	1.05	7/2647 (0.3%)
1	H	0.94	1/1945 (0.1%)	1.01	5/2647 (0.2%)
All	All	0.95	4/15560 (0.0%)	1.05	55/21176 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	1
1	D	0	2
1	E	0	2
1	F	0	1
1	G	0	2
All	All	0	11

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	114	GLU	CD-OE1	7.68	1.34	1.25
1	H	56	SER	CB-OG	6.32	1.50	1.42
1	D	32	GLU	CG-CD	5.78	1.60	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	32	GLU	CD-OE1	5.63	1.31	1.25

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	180	ARG	NE-CZ-NH2	-11.72	114.44	120.30
1	D	90	ARG	NE-CZ-NH2	-11.31	114.64	120.30
1	D	90	ARG	NE-CZ-NH1	10.90	125.75	120.30
1	E	90	ARG	NE-CZ-NH2	-9.78	115.41	120.30
1	E	180	ARG	NE-CZ-NH2	-8.73	115.94	120.30

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	170	GLN	Peptide
1	A	2	SER	Peptide
1	B	1	MET	Peptide
1	C	2	SER	Peptide
1	D	49	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1907	0	1918	63	0
1	B	1907	0	1918	79	0
1	C	1907	0	1918	77	0
1	D	1907	0	1918	70	0
1	E	1907	0	1918	83	0
1	F	1907	0	1918	80	0
1	G	1907	0	1918	52	0
1	H	1907	0	1918	72	0
2	A	15	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	1	0
2	D	15	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	10	0	0	1	0
2	F	10	0	0	0	0
2	G	10	0	0	0	0
2	H	10	0	0	2	0
3	B	25	0	34	3	0
3	D	25	0	34	5	0
3	E	25	0	34	2	0
3	G	25	0	34	3	0
4	A	145	0	0	26	0
4	B	153	0	0	26	0
4	C	156	0	0	33	0
4	D	144	0	0	21	0
4	E	147	0	0	42	0
4	F	150	0	0	31	0
4	G	149	0	0	19	0
4	H	143	0	0	23	0
All	All	16633	0	15480	576	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 576 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:ASN:HB2	4:B:532:HOH:O	1.27	1.32
1:B:111:GLY:HA3	4:B:490:HOH:O	1.27	1.30
1:C:217:GLU:HB2	4:C:546:HOH:O	1.14	1.25
1:H:120:GLN:HG3	4:H:483:HOH:O	1.34	1.24
1:H:193:PRO:HB2	4:H:484:HOH:O	1.38	1.23

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/267 (96%)	239 (94%)	16 (6%)	0	100	100
1	B	255/267 (96%)	236 (92%)	17 (7%)	2 (1%)	19	39
1	C	255/267 (96%)	248 (97%)	7 (3%)	0	100	100
1	D	255/267 (96%)	239 (94%)	13 (5%)	3 (1%)	13	27
1	E	255/267 (96%)	240 (94%)	14 (6%)	1 (0%)	34	57
1	F	255/267 (96%)	237 (93%)	16 (6%)	2 (1%)	19	39
1	G	255/267 (96%)	243 (95%)	12 (5%)	0	100	100
1	H	255/267 (96%)	239 (94%)	15 (6%)	1 (0%)	34	57
All	All	2040/2136 (96%)	1921 (94%)	110 (5%)	9 (0%)	34	57

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	134	VAL
1	H	134	VAL
1	F	2	SER
1	D	109	ALA
1	E	3	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	196/201 (98%)	170 (87%)	26 (13%)	4	7
1	B	196/201 (98%)	171 (87%)	25 (13%)	4	8
1	C	196/201 (98%)	170 (87%)	26 (13%)	4	7
1	D	196/201 (98%)	172 (88%)	24 (12%)	5	9
1	E	196/201 (98%)	168 (86%)	28 (14%)	3	5
1	F	196/201 (98%)	170 (87%)	26 (13%)	4	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	196/201 (98%)	172 (88%)	24 (12%)	5	9
1	H	196/201 (98%)	168 (86%)	28 (14%)	3	5
All	All	1568/1608 (98%)	1361 (87%)	207 (13%)	4	7

5 of 207 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	136	GLU
1	E	87	THR
1	H	80	SER
1	D	196	ASN
1	E	25	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	220	GLN
1	E	170	GLN
1	H	105	ASN
1	E	3	ASN
1	E	200	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	D	302	-	4,4,4	0.38	0	6,6,6	1.18	1 (16%)
2	SO4	B	301	-	4,4,4	0.47	0	6,6,6	0.50	0
2	SO4	A	301	-	4,4,4	0.32	0	6,6,6	0.85	0
3	PE8	B	303	-	24,24,24	0.56	0	23,23,23	1.86	7 (30%)
2	SO4	A	302	-	4,4,4	0.53	0	6,6,6	0.44	0
3	PE8	D	304	-	24,24,24	0.67	0	23,23,23	1.61	4 (17%)
3	PE8	E	303	-	24,24,24	0.58	0	23,23,23	1.87	9 (39%)
2	SO4	B	302	-	4,4,4	0.19	0	6,6,6	0.69	0
2	SO4	H	302	-	4,4,4	0.37	0	6,6,6	1.18	1 (16%)
2	SO4	F	302	-	4,4,4	0.64	0	6,6,6	1.10	1 (16%)
2	SO4	G	301	-	4,4,4	0.20	0	6,6,6	0.41	0
2	SO4	F	301	-	4,4,4	0.45	0	6,6,6	0.64	0
2	SO4	C	301	-	4,4,4	0.37	0	6,6,6	0.45	0
2	SO4	E	302	-	4,4,4	0.23	0	6,6,6	1.10	0
2	SO4	H	301	-	4,4,4	0.38	0	6,6,6	0.93	0
2	SO4	D	301	-	4,4,4	0.41	0	6,6,6	0.44	0
2	SO4	C	302	-	4,4,4	0.38	0	6,6,6	0.76	0
2	SO4	G	302	-	4,4,4	0.34	0	6,6,6	0.25	0
2	SO4	D	303	-	4,4,4	0.22	0	6,6,6	0.54	0
2	SO4	A	303	-	4,4,4	0.41	0	6,6,6	1.06	0
3	PE8	G	303	-	24,24,24	0.60	0	23,23,23	1.46	2 (8%)
2	SO4	E	301	-	4,4,4	0.34	0	6,6,6	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PE8	D	304	-	-	15/22/22/22	-
3	PE8	G	303	-	-	14/22/22/22	-
3	PE8	E	303	-	-	13/22/22/22	-
3	PE8	B	303	-	-	13/22/22/22	-

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	303	PE8	O22-C21-C20	3.33	125.41	110.39
3	B	303	PE8	O10-C9-C8	3.28	125.17	110.39
3	B	303	PE8	O13-C12-C11	3.16	124.66	110.39
3	B	303	PE8	O4-C5-C6	2.97	123.78	110.39
3	D	304	PE8	O22-C23-C24	2.72	122.04	110.07

There are no chirality outliers.

5 of 55 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	303	PE8	O10-C11-C12-O13
3	G	303	PE8	O19-C20-C21-O22
3	B	303	PE8	O16-C17-C18-O19
3	G	303	PE8	O16-C17-C18-O19
3	D	304	PE8	O4-C5-C6-O7


There are no ring outliers.

8 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	302	SO4	1	0
3	B	303	PE8	3	0
3	D	304	PE8	5	0
3	E	303	PE8	2	0
2	H	302	SO4	2	0
2	C	301	SO4	1	0
2	E	302	SO4	1	0
3	G	303	PE8	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/267 (97%)	-0.56	0 100 100	17, 28, 54, 63	0
1	B	259/267 (97%)	-0.51	0 100 100	18, 29, 53, 67	0
1	C	259/267 (97%)	-0.53	1 (0%) 92 91	19, 28, 51, 74	0
1	D	259/267 (97%)	-0.50	0 100 100	17, 27, 53, 66	0
1	E	259/267 (97%)	-0.51	0 100 100	16, 28, 53, 67	0
1	F	259/267 (97%)	-0.49	2 (0%) 86 84	18, 28, 59, 80	0
1	G	259/267 (97%)	-0.52	0 100 100	18, 28, 49, 70	0
1	H	259/267 (97%)	-0.51	1 (0%) 92 91	19, 29, 56, 71	0
All	All	2072/2136 (97%)	-0.52	4 (0%) 95 95	16, 28, 54, 80	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	1	MET	3.9
1	H	1	MET	3.4
1	F	2	SER	3.0
1	C	1	MET	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	302	5/5	0.90	0.31	104,106,110,118	0
2	SO4	D	301	5/5	0.92	0.35	85,90,104,114	0
3	PE8	G	303	25/25	0.93	0.20	41,49,60,62	0
3	PE8	B	303	25/25	0.94	0.16	39,51,54,56	0
3	PE8	E	303	25/25	0.95	0.18	39,45,55,56	0
3	PE8	D	304	25/25	0.96	0.17	40,50,54,57	0
2	SO4	F	302	5/5	0.98	0.15	34,36,42,46	0
2	SO4	H	302	5/5	0.99	0.13	31,38,41,41	0
2	SO4	B	301	5/5	0.99	0.14	42,46,47,55	0
2	SO4	G	301	5/5	0.99	0.14	40,41,43,44	0
2	SO4	F	301	5/5	0.99	0.12	23,23,24,28	0
2	SO4	C	301	5/5	0.99	0.15	26,26,29,31	0
2	SO4	E	302	5/5	0.99	0.16	36,39,42,43	0
2	SO4	A	301	5/5	0.99	0.15	23,23,25,28	0
2	SO4	C	302	5/5	0.99	0.14	33,42,44,44	0
2	SO4	G	302	5/5	0.99	0.11	29,29,34,40	0
2	SO4	D	303	5/5	0.99	0.13	21,24,26,28	0
2	SO4	A	303	5/5	0.99	0.15	26,28,29,32	0
2	SO4	D	302	5/5	0.99	0.13	32,41,48,49	0
2	SO4	E	301	5/5	0.99	0.14	26,29,31,35	0
2	SO4	H	301	5/5	1.00	0.12	21,25,26,28	0
2	SO4	B	302	5/5	1.00	0.13	23,23,25,26	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around PE8 G 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around PE8 B 303:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PE8 E 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PE8 D 304:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.