

Full wwPDB X-ray Structure Validation Report (i)

Sep 17, 2024 – 04:06 PM EDT

PDB ID : 8VPG

Title : CamA Adenine Methyltransferase Complexed to Cognate Substrate DNA and

Containing Quinoline-based SGI-1027 Analog 455

Authors : Zhou, J.; Horton, J.R.; Cheng, X.

Deposited on : 2024-01-16

Resolution : 3.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 2022.3.0, CSD as 543be (2022)

Xtriage (Phenix) : 1.20.1

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.002 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.38.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	164625	2258 (3.10-3.02)
Clashscore	180529	2399 (3.10-3.02)
Ramachandran outliers	177936	2269 (3.10-3.02)
Sidechain outliers	177891	2268 (3.10-3.02)
RSRZ outliers	164620	2258 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	577	82%	12%	6%				
			% 	1270	070				
1	В	577	85% 2%	10%	5%				
1	С	577	83%	10%	7%				
2	Е	14	50% 43%		7%				
2	G	14	43% 57%						



Mol	Chain	Length	Quality of chain						
2	I	14	57%	43%					
3	D	14	50%	50%					
3	F	14	50%	50%					
3	Н	14	43%	57%					



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 15124 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Site-specific DNA-methyltransferase (adenine-specific).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	543	Total	С	N	О	S	0	0	0
1	A	040	4434	2882	705	830	17	0	U	
1	D	551	Total	С	N	О	S	0	0	0
1	Б	991	4570	2974	732	847	17	U	U	0
1	С	537	Total	С	N	О	S	0	0	0
1		357	4330	2812	690	811	17	0		

• Molecule 2 is a DNA chain called DNA Strand II.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	Е	13	Total	С	N	О	Р	0	0	0
	12	10	269	129	45	82	13	U		
9	G	14	Total	С	N	О	Р	0	0	0
	G	14	287	139	50	85	13	U		
9	Т	1.4	Total	С	N	О	Р	0	0	0
2	2 1	14	287	139	50	85	13	0		

• Molecule 3 is a DNA chain called DNA Strand I.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	D	1.4	Total	С	N	О	Р	0	0	0
3	ט	$\begin{array}{ c c c c c c }\hline & 14 & & 2 \\ \hline & & 2 \\ \hline \end{array}$	281	136	53	79	13	U	U	
3	E	14	Total	С	N	О	Р	0	0	0
3	I'	14	281	136	53	79	13	U	U	U
2	Н	14	Total	С	N	О	Р	0	0	0
3	11	14	281	136	53	79	13	U	U	U

• Molecule 4 is N-{3-[(2-amino-6-methylpyrimidin-4-yl)amino]-5-[(dimethylamino)methyl]phe nyl}-3-[(quinolin-4-yl)amino]benzamide (three-letter code: A1AC7) (formula: $C_{30}H_{30}N_8O$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	С	N	О	0	0
4	ש	1	39	30	8	1	0	U

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	10	Total O 10 10	0	0
5	В	35	Total O 35 35	0	0
5	С	8	Total O 8 8	0	0
5	Е	3	Total O 3 3	0	0
5	D	2	Total O 2 2	0	0
5	F	3	Total O 3 3	0	0
5	Н	1	Total O 1 1	0	0
5	I	3	Total O 3 3	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Site-specific DNA-methyltransferase (adenine-specific)







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	80.54Å 161.60Å 234.98Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.84 - 3.05	Depositor
Resolution (A)	41.84 - 3.05	EDS
% Data completeness	99.6 (41.84-3.05)	Depositor
(in resolution range)	99.6 (41.84-3.05)	EDS
R_{merge}	0.38	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.24 (at 3.06Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
D D.	0.202 , 0.235	Depositor
R, R_{free}	0.203 , 0.235	DCC
R_{free} test set	57218 reflections (3.38%)	wwPDB-VP
Wilson B-factor (Å ²)	81.2	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 60.6	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	15124	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A1AC7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.24	0/4524	0.43	0/6103
1	В	0.24	0/4664	0.44	0/6276
1	С	0.24	0/4418	0.43	0/5964
2	Е	0.51	0/300	1.06	0/462
2	G	0.52	0/321	1.00	0/495
2	I	0.52	0/321	1.00	0/495
3	D	0.57	0/315	0.96	0/483
3	F	0.50	0/315	0.90	0/483
3	Н	0.47	0/315	0.92	0/483
All	All	0.29	0/15493	0.54	0/21244

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4434	0	4330	38	0
1	В	4570	0	4562	30	0
1	С	4330	0	4190	32	0
2	Ε	269	0	150	5	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	287	0	162	6	0
2	I	287	0	162	3	0
3	D	281	0	159	5	0
3	F	281	0	159	4	0
3	Н	281	0	159	5	0
4	D	39	0	0	4	0
5	A	10	0	0	0	0
5	В	35	0	0	1	0
5	С	8	0	0	1	0
5	D	2	0	0	0	0
5	\mathbf{E}	3	0	0	0	0
5	F	3	0	0	0	0
5	Н	1	0	0	0	0
5	I	3	0	0	0	0
All	All	15124	0	14033	125	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (125) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance } (\text{\AA}) \end{array}$	Clash overlap (Å)
1:A:281:LYS:HE3	1:A:303:GLU:HB3	1.64	0.79
1:A:69:LEU:HD12	1:A:123:LEU:HD21	1.71	0.72
2:G:10:DT:H2"	2:G:11:DT:H72	1.73	0.71
1:A:232:ARG:NH1	1:A:314:ASP:O	2.27	0.68
3:H:3:DC:H2'	3:H:4:DA:C8	2.29	0.67
2:I:2:DT:H2"	2:I:3:DG:C8	2.30	0.67
1:C:43:LYS:HD2	1:C:264:THR:HG21	1.77	0.66
1:C:432:ARG:NH2	3:H:2:DT:OP2	2.29	0.65
1:A:66:GLY:HA3	1:A:123:LEU:HD13	1.80	0.64
3:F:3:DC:H2'	3:F:4:DA:C8	2.33	0.64
2:I:10:DT:H2"	2:I:11:DT:H72	1.81	0.62
1:C:386:ILE:HD13	1:C:550:LEU:HB3	1.82	0.61
3:H:12:DC:H2"	3:H:13:DC:C5	2.36	0.60
1:B:58:ILE:HD13	1:B:76:LEU:HD11	1.84	0.60
1:C:203:LYS:HE3	1:C:207:ILE:HD13	1.84	0.59
1:B:131:LYS:HE2	1:B:133:VAL:HB	1.84	0.59
1:C:168:TYR:CE1	1:C:219:PRO:HD3	2.38	0.59
1:A:225:SER:O	1:A:232:ARG:NH2	2.36	0.59
1:B:69:LEU:HD12	1:B:123:LEU:HD21	1.84	0.58



 $Continued\ from\ previous\ page...$

Continuea from previ		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap $(Å)$
1:B:51:ILE:HD11	1:B:83:ASN:HB3	1.84	0.58
2:E:8:DT:H2"	2:E:9:DT:H71	1.85	0.57
1:A:242:GLN:NE2	1:A:266:ASP:OD2	2.37	0.57
1:A:248:LEU:HD23	1:A:280:ILE:HG22	1.87	0.57
1:A:402:ASN:HD21	1:A:405:THR:HG23	1.69	0.57
1:C:66:GLY:HA3	1:C:123:LEU:HD13	1.85	0.56
1:C:69:LEU:HD12	1:C:123:LEU:HD21	1.86	0.56
1:C:206:ASP:OD1	1:C:234:TYR:OH	2.22	0.56
1:A:465:ASP:OD1	1:A:469:ASN:ND2	2.39	0.55
3:D:13:DC:H2"	3:D:14:DA:C8	2.42	0.54
1:B:292:THR:OG1	1:B:294:GLU:OE1	2.26	0.54
1:C:195:ASP:OD1	1:C:196:LEU:N	2.41	0.54
1:C:401:ASP:OD1	1:C:401:ASP:N	2.41	0.54
1:A:114:ASP:OD1	1:A:115:ILE:N	2.42	0.53
1:A:378:LYS:NZ	2:E:9:DT:OP2	2.29	0.53
3:H:1:DT:H2'	3:H:2:DT:C6	2.44	0.53
1:B:168:TYR:CE1	1:B:219:PRO:HD3	2.43	0.53
1:C:69:LEU:HD22	1:C:110:ILE:HG23	1.91	0.53
1:B:447:GLU:HA	1:B:468:ASN:HB3	1.91	0.52
1:B:386:ILE:HD13	1:B:550:LEU:HB3	1.91	0.52
1:B:195:ASP:OD1	1:B:196:LEU:N	2.42	0.52
2:I:12:DT:H2"	2:I:13:DG:C8	2.45	0.52
3:D:9:DG:H21	4:D:101:A1AC7:C33	2.23	0.52
1:C:552:ASN:OD1	1:C:553:LYS:N	2.42	0.52
3:D:12:DC:O4'	4:D:101:A1AC7:N1	2.43	0.51
2:G:2:DT:H2"	2:G:3:DG:C8	2.45	0.51
1:B:401:ASP:N	1:B:401:ASP:OD1	2.43	0.51
1:C:290:PHE:CG	1:C:296:LEU:HD11	2.46	0.51
1:A:51:ILE:HD11	1:A:83:ASN:HB3	1.94	0.50
2:G:1:DA:H1'	2:G:2:DT:H5'	1.94	0.50
1:C:97:THR:HG23	1:C:100:ASN:H	1.76	0.50
1:C:316:TRP:HB3	1:C:318:LEU:HD13	1.93	0.49
3:D:6:DA:H2"	3:D:7:DA:C8	2.47	0.49
1:A:451:ILE:HG13	1:A:490:TYR:HD1	1.77	0.49
1:B:254:LYS:NZ	2:G:3:DG:OP2	2.45	0.49
1:A:354:LYS:HD2	1:A:354:LYS:H	1.77	0.49
1:A:126:SER:O	1:A:130:LYS:HG2	2.12	0.49
1:A:369:ASP:HB3	1:A:372:PHE:HD1	1.78	0.48
1:A:243:GLU:HA	1:A:276:ASP:O	2.13	0.48
1:A:88:LYS:HE2	1:A:96:TRP:O	2.14	0.48
3:H:5:DA:H2"	3:H:6:DA:C8	2.49	0.48



 $Continued\ from\ previous\ page...$

Atom 1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	$\text{overlap } (\mathring{\mathbf{A}})$
1:A:369:ASP:HB3	1:A:372:PHE:CD1	2.48	0.48
1:C:504:LYS:O	1:C:508:ILE:HG12	2.13	0.48
1:C:40:ILE:O	1:C:44:THR:OG1	2.32	0.47
1:B:383:ASN:ND2	5:B:604:HOH:O	2.35	0.47
1:B:464:ILE:HG13	1:B:494:VAL:HG11	1.97	0.47
1:B:433:LYS:HB2	1:B:436:GLU:HG3	1.97	0.47
3:F:10:DT:H1'	3:F:11:DC:H5'	1.96	0.46
1:C:403:GLU:O	1:C:407:LYS:HG2	2.14	0.46
3:F:11:DC:H1'	3:F:12:DC:H5'	1.97	0.46
1:A:58:ILE:HD12	1:A:76:LEU:HD11	1.98	0.46
1:A:464:ILE:HG13	1:A:494:VAL:HG11	1.96	0.46
1:C:333:LYS:HG2	1:C:575:LEU:HA	1.98	0.46
1:C:234:TYR:O	1:C:238:ASN:HB2	2.15	0.45
1:C:338:LEU:HD23	1:C:338:LEU:HA	1.83	0.45
1:B:124:LYS:HG3	1:B:143:ILE:HD12	1.98	0.45
1:C:275:ILE:HD11	1:C:309:GLN:HB2	1.97	0.45
1:A:236:LYS:NZ	1:A:312:LEU:O	2.44	0.45
1:C:319:VAL:HG11	1:C:508:ILE:HG23	1.99	0.45
1:B:143:ILE:HG22	1:B:145:LEU:HG	1.99	0.45
2:E:11:DT:C6	2:E:11:DT:H5'	2.52	0.45
2:G:4:DG:H2"	2:G:5:DG:C8	2.52	0.44
1:A:143:ILE:HG22	1:A:145:LEU:HG	1.99	0.44
1:A:534:ASN:OD1	1:A:534:ASN:N	2.49	0.44
1:A:431:ILE:HG12	2:G:14:DA:H2"	1.99	0.44
1:A:40:ILE:HG23	1:A:215:SER:HB3	2.00	0.44
1:B:201:TYR:CE2	1:B:218:THR:HG21	2.53	0.44
1:A:319:VAL:HG11	1:A:508:ILE:HG23	1.99	0.43
1:B:175:GLU:HG3	1:B:178:TYR:H	1.83	0.43
1:A:350:THR:HG22	1:A:437:LEU:HD22	2.00	0.43
1:C:326:PHE:CE1	1:C:577:ILE:HD12	2.53	0.43
2:E:3:DG:H2"	2:E:4:DG:C8	2.53	0.43
1:B:384:LYS:HD3	1:B:498:ASN:HB3	2.01	0.43
2:E:5:DG:H21	4:D:101:A1AC7:C37	2.30	0.43
3:F:13:DC:H2"	3:F:14:DA:C8	2.52	0.43
1:C:432:ARG:NH1	5:C:602:HOH:O	2.52	0.43
1:B:114:ASP:OD1	1:B:115:ILE:N	2.52	0.43
1:B:171:HIS:CD2	1:B:193:LYS:HD2	2.54	0.43
1:B:338:LEU:HD23	1:B:338:LEU:HA	1.86	0.43
4:D:101:A1AC7:C17	4:D:101:A1AC7:C13	2.96	0.43
1:A:512:LYS:NZ	1:A:516:ASN:H	2.16	0.43
1:C:447:GLU:HA	1:C:468:ASN:HB3	2.00	0.43



Continued from previous page...

A	A	Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)	
1:A:325:THR:HG22	1:A:329:LYS:HE2	2.00	0.43	
1:A:277:VAL:O	1:A:304:HIS:HA	2.19	0.42	
1:B:350:THR:HG22	1:B:437:LEU:HD22	1.99	0.42	
1:B:345:PHE:CE2	1:B:477:SER:HB2	2.55	0.42	
1:A:503:ASP:OD1	1:A:507:LYS:NZ	2.48	0.42	
1:A:36:ILE:HG21	1:A:217:ILE:HD12	2.01	0.42	
1:B:337:SER:HA	1:B:529:ARG:HA	2.01	0.42	
1:C:331:GLN:OE1	1:C:529:ARG:HG3	2.19	0.42	
3:D:9:DG:H1'	3:D:10:DT:H5"	2.01	0.42	
1:A:337:SER:HA	1:A:529:ARG:HA	2.01	0.42	
1:B:450:LYS:NZ	1:B:465:ASP:OD2	2.51	0.41	
1:C:377:ILE:HD11	1:C:452:MET:HE1	2.01	0.41	
1:A:229:LYS:HA	1:A:232:ARG:NH1	2.35	0.41	
1:B:392:TYR:O	1:B:393:ARG:NH1	2.47	0.41	
1:B:203:LYS:O	1:B:207:ILE:HG12	2.20	0.41	
1:C:174:LEU:HD23	1:C:179:LYS:HG3	2.03	0.41	
1:C:201:TYR:CE2	1:C:218:THR:HG21	2.55	0.41	
1:A:248:LEU:HD21	1:A:279:LYS:HG3	2.03	0.41	
1:C:420:LYS:HA	1:C:420:LYS:HD2	1.89	0.41	
1:C:350:THR:HG22	1:C:437:LEU:HD22	2.03	0.41	
1:A:256:ILE:HG22	1:A:258:VAL:HG12	2.01	0.41	
1:B:36:ILE:HD12	1:B:262:ILE:HD11	2.03	0.41	
1:A:50:ILE:HD11	1:A:80:PHE:CZ	2.57	0.40	
1:B:504:LYS:O	1:B:508:ILE:HG12	2.22	0.40	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	539/577 (93%)	521 (97%)	18 (3%)	0	100 100	



Mol	Chain	Analysed	Favoured	Tavoured Allowed		Perce	Percentiles	
1	В	549/577 (95%)	529 (96%)	20 (4%)	0	100	100	
1	\mathbf{C}	533/577 (92%)	519 (97%)	14 (3%)	0	100	100	
All	All	1621/1731 (94%)	1569 (97%)	52 (3%)	0	100	100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	478/547 (87%)	471 (98%)	7 (2%)	60 77
1	В	503/547 (92%)	496 (99%)	7 (1%)	62 79
1	С	461/547 (84%)	454 (98%)	7 (2%)	60 77
All	All	1442/1641 (88%)	1421 (98%)	21 (2%)	60 77

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	ILE
1	A	183	LEU
1	A	243	GLU
1	A	263	LEU
1	A	317	ILE
1	A	455	TYR
1	A	555	ILE
1	В	97	THR
1	В	141	ILE
1	В	168	TYR
1	В	263	LEU
1	В	401	ASP
1	В	455	TYR
1	В	519	ASP
1	С	38	ASP
1	С	44	THR



Mol	Chain	Res	Type
1	С	168	TYR
1	С	263	LEU
1	С	408	ARG
1	С	455	TYR
1	С	519	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	104	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain	hain Res	Dog	Pog	Ros	Ros	Pos	Pog	Link	Bo	ond leng	${ m ths}$	В	ond ang	eles
	MIOI	Type			LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2						
	4	A1AC7	D	101	-	43,43,43	0.43	0	59,60,60	0.74	4 (6%)						



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A1AC7	D	101	-	-	6/20/20/20	0/5/5/5

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	D	101	A1AC7	C12-N10-C9	2.55	115.87	110.69
4	D	101	A1AC7	C14-N15-C16	2.11	132.17	126.61
4	D	101	A1AC7	C13-C14-N15	2.11	126.98	120.13
4	D	101	A1AC7	C11-N10-C9	2.00	114.76	110.69

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	101	A1AC7	C17-C16-N15-C14
4	D	101	A1AC7	O34-C16-N15-C14
4	D	101	A1AC7	O34-C16-C17-C33
4	D	101	A1AC7	O34-C16-C17-C18
4	D	101	A1AC7	N15-C16-C17-C33
4	D	101	A1AC7	N15-C16-C17-C18

There are no ring outliers.


1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	101	A1AC7	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and



any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	543/577 (94%)	0.15	10 (1%) 67 48	54, 95, 138, 168	0
1	В	551/577 (95%)	-0.12	4 (0%) 84 69	49, 79, 119, 177	0
1	С	537/577 (93%)	0.16	9 (1%) 69 49	66, 112, 172, 225	0
2	E	13/14 (92%)	0.43	0 100 100	85, 96, 211, 223	0
2	G	14/14 (100%)	-0.33	0 100 100	60, 71, 109, 109	0
2	I	14/14 (100%)	-0.13	0 100 100	79, 90, 224, 234	0
3	D	14/14 (100%)	0.32	0 100 100	70, 106, 199, 215	0
3	F	14/14 (100%)	-0.14	0 100 100	58, 68, 123, 125	0
3	Н	14/14 (100%)	0.31	0 100 100	81, 104, 188, 203	0
All	All	1714/1815 (94%)	0.06	23 (1%) 74 56	49, 94, 163, 234	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	HIS	4.1
1	С	317	ILE	3.9
1	В	27	SER	3.6
1	С	84	ILE	3.5
1	A	192	ASP	3.4
1	A	65	CYS	3.2
1	A	255	ASN	3.0
1	В	132	VAL	2.6
1	A	169	ILE	2.6
1	В	133	VAL	2.5
1	С	146	PHE	2.4
1	A	199	CYS	2.4
1	С	85	TYR	2.3
1	В	28	GLY	2.3
1	A	132	VAL	2.3



Mol	Chain	Res	Type	RSRZ
1	A	170	GLY	2.3
1	С	113	ALA	2.3
1	С	56	PRO	2.3
1	С	556	ASP	2.2
1	С	509	THR	2.1
1	A	513	MET	2.0
1	A	141	ILE	2.0
1	С	272	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ m B ext{-}factors(\AA^2)$	Q<0.9
4	A1AC7	D	101	39/39	0.65	0.21	94,127,159,165	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

