

wwPDB X-ray Structure Validation Summary Report (i)

Oct 5, 2023 – 03:57 AM EDT

PDB ID	:	6VUV
Title	:	Scabin (S117A) toxin from Streptomyces scabies
Authors	:	Lyons, B.; Lidster, T.; Merrill, A.R.
Deposited on		
Resolution	:	1.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	FAILED
Xtriage (Phenix)	:	1.13
EDS	:	FAILED
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\hbox{-}RAY\,DIFFRACTION$

The reported resolution of this entry is 1.55 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2653 atoms, of which 1199 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Scabin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	А	165	Total 2502	C 832	Н 1199	N 214	O 252	${f S}{5}$	0	5	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	6	MET	-	expression tag	UNP C9Z6T8
А	7	GLY	-	expression tag	UNP C9Z6T8
А	8	SER	-	expression tag	UNP C9Z6T8
А	9	SER	-	expression tag	UNP C9Z6T8
А	10	HIS	-	expression tag	UNP C9Z6T8
А	11	HIS	-	expression tag	UNP C9Z6T8
A	12	HIS	-	expression tag	UNP C9Z6T8
А	13	HIS	-	expression tag	UNP C9Z6T8
А	14	HIS	-	expression tag	UNP C9Z6T8
А	15	HIS	-	expression tag	UNP C9Z6T8
А	16	SER	-	expression tag	UNP C9Z6T8
A	17	SER	-	expression tag	UNP C9Z6T8
A	18	GLY	-	expression tag	UNP C9Z6T8
А	19	GLU	-	expression tag	UNP C9Z6T8
A	20	ASN	-	expression tag	UNP C9Z6T8
А	21	LEU	-	expression tag	UNP C9Z6T8
A	22	TYR	-	expression tag	UNP C9Z6T8
А	23	PHE	-	expression tag	UNP C9Z6T8
А	24	GLN	-	expression tag	UNP C9Z6T8
А	25	GLY	-	expression tag	UNP C9Z6T8
А	26	SER	-	expression tag	UNP C9Z6T8
А	27	HIS	-	expression tag	UNP C9Z6T8
А	28	MET	-	expression tag	UNP C9Z6T8
А	117	ALA	SER	engineered mutation	UNP C9Z6T8

There are 24 discrepancies between the modelled and reference sequences:

• Molecule 2 is water.



Mo	Chain	Residues	Atoms		ZeroOcc	AltConf
2	А	151	Total 151	O 151	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	88.45Å 60.85 Å 37.95 Å	Depositor	
a, b, c, α , β , γ	90.00° 99.60° 90.00°	Depositor	
Resolution (Å)	43.61 - 1.55	Depositor	
% Data completeness	99.7 (43.61-1.55)	Depositor	
(in resolution range)		-	
R _{merge}	0.04	Depositor	
R _{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	$1.21 (at 1.31 \text{\AA})$	Xtriage	
Refinement program	PHENIX 1.14_3260	Depositor	
R, R_{free}	0.179 , 0.210	Depositor	
Wilson B-factor $(Å^2)$	23.0	Xtriage	
Anisotropy	0.177	Xtriage	
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	2653	wwPDB-VP	
Average B, all atoms $(Å^2)$	33.0	wwPDB-VP	

EDS failed to run properly - this section is therefore incomplete.

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.



4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

