



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 2, 2023 – 11:37 PM EDT

PDB ID : 3VYG
Title : Crystal structure of Thiocyanate hydrolase mutant R136W
Authors : Yamanaka, Y.; Sato, M.; Arakawa, T.; Namima, S.; Hori, S.; Ohtaki, A.;
Noguchi, K.; Katayama, Y.; Yohda, M.; Odaka, M.
Deposited on : 2012-09-25
Resolution : 1.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	126	 84% 9% • 5%
1	D	126	 80% 12% • 6%
1	G	126	 75% 19% • 5%
1	J	126	 80% 14% • 5%
2	B	157	 77% 17% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	E	157	 76% 18% . .
2	H	157	 81% 15% . .
2	K	157	 % 77% 18% . .
3	C	243	 77% 11% . 11%
3	F	243	 79% 10% 11%
3	I	243	 74% 13% . 11%
3	L	243	 74% 12% . 11%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 17343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thiocyanate hydrolase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	120	974	620	162	188	4	15	0	0
1	D	119	973	620	161	188	4	10	1	0
1	G	120	982	626	163	189	4	16	1	0
1	J	120	981	625	163	189	4	8	1	0

- Molecule 2 is a protein called Thiocyanate hydrolase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	152	1244	787	224	227	6	0	2	0
2	E	151	1248	790	226	226	6	0	3	0
2	H	151	1226	775	221	224	6	0	0	0
2	K	151	1224	773	221	224	6	0	0	0

- Molecule 3 is a protein called Thiocyanate hydrolase subunit gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	217	1743	1114	306	315	8	10	2	0
3	F	217	1744	1113	306	317	8	0	2	0
3	I	217	1727	1103	302	314	8	7	0	0
3	L	217	1733	1107	303	315	8	17	1	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	136	TRP	ARG	engineered mutation	UNP O66188
F	136	TRP	ARG	engineered mutation	UNP O66188
I	136	TRP	ARG	engineered mutation	UNP O66188
L	136	TRP	ARG	engineered mutation	UNP O66188

- Molecule 4 is COBALT (III) ION (three-letter code: 3CO) (formula: Co).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Co 1 1	0	0
4	F	1	Total Co 1 1	0	0
4	I	1	Total Co 1 1	0	0
4	L	1	Total Co 1 1	0	0

- Molecule 5 is L(+)-TARTARIC ACID (three-letter code: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total C O 10 4 6	0	0
5	F	1	Total C O 10 4 6	0	0
5	I	1	Total C O 10 4 6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			10	4	6		



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	71	Total	O	0	0
			71	71		
6	B	129	Total	O	0	0
			129	129		
6	C	214	Total	O	0	0
			214	214		
6	D	78	Total	O	0	0
			78	78		
6	E	128	Total	O	0	0
			128	128		
6	F	189	Total	O	0	0
			189	189		
6	G	84	Total	O	0	0
			84	84		
6	H	107	Total	O	0	0
			107	107		
6	I	148	Total	O	0	0
			148	148		
6	J	90	Total	O	0	0
			90	90		
6	K	109	Total	O	0	0
			109	109		
6	L	153	Total	O	0	0
			153	153		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Thiocyanate hydrolase subunit alpha




- Molecule 1: Thiocyanate hydrolase subunit alpha




- Molecule 1: Thiocyanate hydrolase subunit alpha




- Molecule 1: Thiocyanate hydrolase subunit alpha




- Molecule 2: Thiocyanate hydrolase subunit beta




- Molecule 2: Thiocyanate hydrolase subunit beta




- Molecule 2: Thiocyanate hydrolase subunit beta




- Molecule 2: Thiocyanate hydrolase subunit beta



- Molecule 3: Thiocyanate hydrolase subunit gamma




- Molecule 3: Thiocyanate hydrolase subunit gamma




- Molecule 3: Thiocyanate hydrolase subunit gamma





- Molecule 3: Thiocyanate hydrolase subunit gamma

Chain L: 74% 12% 11%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.45Å 172.40Å 172.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.96 – 1.72 33.96 – 1.72	Depositor EDS
% Data completeness (in resolution range)	96.5 (33.96-1.72) 96.2 (33.96-1.72)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.52 (at 1.72Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.187 , 0.218 0.188 , 0.217	Depositor DCC
R_{free} test set	17284 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 31.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.447 for -h,l,k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17343	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 3CO, TLA, CSO, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.46	4/1000 (0.4%)	1.15	3/1354 (0.2%)
1	D	1.48	5/999 (0.5%)	1.17	5/1353 (0.4%)
1	G	1.52	7/1008 (0.7%)	1.36	13/1365 (1.0%)
1	J	1.51	7/1007 (0.7%)	1.32	7/1364 (0.5%)
2	B	1.41	8/1279 (0.6%)	1.35	14/1741 (0.8%)
2	E	1.40	6/1283 (0.5%)	1.27	7/1746 (0.4%)
2	H	1.36	2/1258 (0.2%)	1.27	7/1712 (0.4%)
2	K	1.39	5/1256 (0.4%)	1.22	13/1709 (0.8%)
3	C	1.48	5/1776 (0.3%)	1.30	12/2425 (0.5%)
3	F	1.49	6/1777 (0.3%)	1.26	9/2426 (0.4%)
3	I	1.40	7/1757 (0.4%)	1.26	13/2400 (0.5%)
3	L	1.50	12/1766 (0.7%)	1.22	13/2412 (0.5%)
All	All	1.45	74/16166 (0.5%)	1.26	116/22007 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	E	0	1

The worst 5 of 74 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	50	ARG	CD-NE	-12.79	1.24	1.46
3	L	79	LYS	CD-CE	-10.39	1.25	1.51
1	D	34	LYS	CB-CG	8.68	1.75	1.52
3	F	81	CYS	CB-SG	-8.04	1.68	1.82
3	C	81	CYS	CB-SG	-7.89	1.68	1.82

The worst 5 of 116 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	138	MET	CG-SD-CE	-13.02	79.36	100.20
2	H	91	ARG	NE-CZ-NH2	-11.13	114.73	120.30
3	I	109	ASP	CB-CG-OD2	10.79	128.01	118.30
3	L	109	ASP	CB-CG-OD2	10.70	127.93	118.30
2	E	138	MET	CG-SD-CE	-10.55	83.32	100.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	E	4	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	974	0	926	17	0
1	D	973	0	924	17	0
1	G	982	0	936	14	0
1	J	981	0	934	10	0
2	B	1244	0	1211	22	0
2	E	1248	0	1222	27	0
2	H	1226	0	1193	19	0
2	K	1224	0	1186	24	0
3	C	1743	0	1755	16	0
3	F	1744	0	1750	11	0
3	I	1727	0	1732	30	0
3	L	1733	0	1740	16	0
4	C	1	0	0	0	0
4	F	1	0	0	0	0
4	I	1	0	0	0	0
4	L	1	0	0	0	0
5	C	10	0	4	0	0
5	F	10	0	4	0	0
5	I	10	0	4	0	0
5	L	10	0	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	71	0	0	1	0
6	B	129	0	0	0	0
6	C	214	0	0	3	0
6	D	78	0	0	0	0
6	E	128	0	0	2	0
6	F	189	0	0	0	0
6	G	84	0	0	1	0
6	H	107	0	0	3	0
6	I	148	0	0	4	0
6	J	90	0	0	0	0
6	K	109	0	0	5	0
6	L	153	0	0	2	0
All	All	17343	0	15525	170	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 170 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:I:47:GLU:HG2	3:I:50:ARG:NH2	1.40	1.32
2:K:65:GLU:OE1	2:K:70:LEU:HB2	1.17	1.27
2:B:137:GLU:OE1	2:B:141:ARG:HD2	1.35	1.22
2:B:137:GLU:OE1	2:B:141:ARG:CD	1.97	1.11
1:D:10:TRP:HZ3	6:H:283:HOH:O	1.36	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/126 (94%)	114 (97%)	4 (3%)	0	100	100
1	D	118/126 (94%)	114 (97%)	4 (3%)	0	100	100
1	G	119/126 (94%)	114 (96%)	5 (4%)	0	100	100
1	J	119/126 (94%)	115 (97%)	4 (3%)	0	100	100
2	B	152/157 (97%)	147 (97%)	5 (3%)	0	100	100
2	E	152/157 (97%)	148 (97%)	4 (3%)	0	100	100
2	H	149/157 (95%)	146 (98%)	3 (2%)	0	100	100
2	K	149/157 (95%)	145 (97%)	4 (3%)	0	100	100
3	C	215/243 (88%)	206 (96%)	8 (4%)	1 (0%)	29	13
3	F	215/243 (88%)	209 (97%)	6 (3%)	0	100	100
3	I	213/243 (88%)	207 (97%)	6 (3%)	0	100	100
3	L	214/243 (88%)	209 (98%)	5 (2%)	0	100	100
All	All	1933/2104 (92%)	1874 (97%)	58 (3%)	1 (0%)	51	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	132	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/108 (94%)	100 (98%)	2 (2%)	55	37
1	D	102/108 (94%)	99 (97%)	3 (3%)	42	22
1	G	103/108 (95%)	101 (98%)	2 (2%)	57	39
1	J	103/108 (95%)	100 (97%)	3 (3%)	42	22
2	B	131/134 (98%)	128 (98%)	3 (2%)	50	31
2	E	132/134 (98%)	130 (98%)	2 (2%)	65	49
2	H	129/134 (96%)	125 (97%)	4 (3%)	40	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	K	128/134 (96%)	124 (97%)	4 (3%)	40	19
3	C	190/212 (90%)	189 (100%)	1 (0%)	88	83
3	F	190/212 (90%)	188 (99%)	2 (1%)	73	62
3	I	188/212 (89%)	180 (96%)	8 (4%)	29	10
3	L	189/212 (89%)	183 (97%)	6 (3%)	39	18
All	All	1687/1816 (93%)	1647 (98%)	40 (2%)	50	29

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	26	GLN
3	L	141[A]	GLN
1	J	34	LYS
2	K	58	ILE
3	L	170	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	26	GLN
2	K	100	GLN
1	J	84	ASN
2	K	28	HIS
3	L	175	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	CSO	F	133	3,4	3,6,7	1.02	0	0,6,8	-	-
3	CSD	I	131	3,4	3,7,8	0.23	0	1,8,10	1.49	0
3	CSD	F	131	3,4	3,7,8	1.13	0	1,8,10	0.80	0
3	CSD	L	131	3,4	3,7,8	0.85	0	1,8,10	1.53	0
3	CSO	C	133	3,4	3,6,7	1.11	0	0,6,8	-	-
3	CSO	L	133	3,4	3,6,7	1.10	0	0,6,8	-	-
3	CSD	C	131	3,4	3,7,8	0.98	0	1,8,10	2.43	1 (100%)
3	CSO	I	133	3,4	3,6,7	1.42	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CSO	F	133	3,4	-	0/1/5/7	-
3	CSD	I	131	3,4	-	0/2/6/8	-
3	CSD	F	131	3,4	-	0/2/6/8	-
3	CSD	L	131	3,4	-	0/2/6/8	-
3	CSO	C	133	3,4	-	0/1/5/7	-
3	CSO	L	133	3,4	-	0/1/5/7	-
3	CSD	C	131	3,4	-	0/2/6/8	-
3	CSO	I	133	3,4	-	0/1/5/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	C	131	CSD	OD1-SG-CB	-2.43	100.91	105.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	TLA	C	302	-	9,9,9	1.40	2 (22%)	12,12,12	1.63	2 (16%)
5	TLA	F	302	-	9,9,9	1.15	0	12,12,12	1.34	2 (16%)
5	TLA	I	302	-	9,9,9	1.16	0	12,12,12	0.91	1 (8%)
5	TLA	L	302	-	9,9,9	1.19	1 (11%)	12,12,12	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TLA	C	302	-	-	0/12/12/12	-
5	TLA	F	302	-	-	0/12/12/12	-
5	TLA	I	302	-	-	0/12/12/12	-
5	TLA	L	302	-	-	0/12/12/12	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	L	302	TLA	O41-C4	-2.48	1.22	1.30
5	C	302	TLA	O1-C1	2.27	1.29	1.22
5	C	302	TLA	C2-C1	2.23	1.55	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
5	C	302	TLA	O11-C1-O1	3.57	132.19	124.09
5	F	302	TLA	O41-C4-C3	2.37	119.67	113.27
5	C	302	TLA	O41-C4-C3	2.31	119.52	113.27
5	F	302	TLA	O11-C1-O1	2.24	129.17	124.09
5	I	302	TLA	O3-C3-C2	-2.03	106.20	110.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	120/126 (95%)	-0.27	1 (0%) 86 89	13, 20, 30, 38	4 (3%)
1	D	119/126 (94%)	-0.29	0 100 100	14, 20, 30, 37	3 (2%)
1	G	120/126 (95%)	-0.30	1 (0%) 86 89	12, 16, 29, 43	5 (4%)
1	J	120/126 (95%)	-0.32	0 100 100	11, 16, 29, 36	2 (1%)
2	B	152/157 (96%)	-0.31	0 100 100	12, 16, 30, 39	0
2	E	151/157 (96%)	-0.32	0 100 100	12, 16, 31, 37	0
2	H	151/157 (96%)	-0.24	0 100 100	12, 19, 39, 48	0
2	K	151/157 (96%)	-0.15	1 (0%) 87 90	12, 18, 39, 50	0
3	C	215/243 (88%)	-0.30	0 100 100	12, 17, 28, 37	4 (1%)
3	F	215/243 (88%)	-0.32	0 100 100	12, 17, 28, 45	0
3	I	215/243 (88%)	-0.23	1 (0%) 91 92	12, 19, 35, 49	2 (0%)
3	L	215/243 (88%)	-0.17	1 (0%) 91 92	12, 19, 35, 53	5 (2%)
All	All	1944/2104 (92%)	-0.26	5 (0%) 94 95	11, 18, 33, 53	25 (1%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	L	239	VAL	6.8
1	G	126	ALA	3.0
3	I	239	VAL	2.3
1	A	126	ALA	2.3
2	K	153	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CSO	F	133	7/8	0.96	0.10	13,15,20,31	1
3	CSO	C	133	7/8	0.97	0.09	13,14,20,29	1
3	CSO	I	133	7/8	0.97	0.09	12,14,21,32	1
3	CSO	L	133	7/8	0.97	0.09	13,14,21,33	1
3	CSD	C	131	8/9	0.98	0.07	12,15,18,18	2
3	CSD	L	131	8/9	0.98	0.07	12,15,17,17	2
3	CSD	I	131	8/9	0.98	0.09	12,16,17,18	2
3	CSD	F	131	8/9	0.99	0.07	12,15,18,19	2

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	TLA	C	302	10/10	0.94	0.11	18,27,34,37	0
5	TLA	F	302	10/10	0.95	0.08	20,24,34,35	0
5	TLA	I	302	10/10	0.95	0.08	18,21,30,32	0
5	TLA	L	302	10/10	0.96	0.09	20,22,28,32	0
4	3CO	F	301	1/1	0.98	0.08	12,12,12,12	1
4	3CO	C	301	1/1	0.99	0.09	12,12,12,12	1
4	3CO	I	301	1/1	0.99	0.10	13,13,13,13	1
4	3CO	L	301	1/1	1.00	0.08	13,13,13,13	1

6.5 Other polymers [i](#)

There are no such residues in this entry.