WORLDWIDE wwPDB PROTEIN DATA BANK

wwPDB X-ray Structure Validation Summary Report (i)

Dec 10, 2023 – 03:52 pm GMT

PDB ID	:	2VZE
Title	:	Crystal structure of human acyl-CoA synthetase medium-chain family member 2A (L64P mutation) in complex with AMP
Authors	:	Yue, W.W.; Kochan, G.T.; Pilka, E.S.; Bhatia, C.; von Delft, F.; Arrowsmith, C.H.; Edwards, A.M.; Wikstrom, M.; Bountra, C.; Oppermann, U.
Deposited on	:	2008-07-31
Resolution	:	2.45 Å(reported)

The following versions of software and data (see references (i)) were used in the production of this report:

|		
MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1   Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percentile Ranks	Value
Rfree		0.254
Clashscore		6
Ramachandran outliers		0.1%
Sidechain outliers		1.9%
RSRZ outliers		1.1%

Worse   *Better*

▌ Percentile relative to all X-ray structures

▯ Percentile relative to X-ray structures of similar resolution

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R$_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	570	83% 9% · 6%
1	B	570	84% 9% · 6%
1	C	570	82% 11% · 6%

2 Entry composition ⓘ

There are 4 unique types of molecules in this entry. The entry contains 13191 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACYL-COENZYME A SYNTHETASE ACSM2A, MITO-CHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	533	Total	C	N	O	S	0	6	0
			4175	2668	703	776	28			
1	B	533	Total	C	N	O	S	0	2	0
			4120	2636	688	768	28			
1	C	536	Total	C	N	O	S	0	5	0
			4171	2665	700	778	28			

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	expression tag	UNP Q08AH3
A	9	GLY	-	expression tag	UNP Q08AH3
A	10	HIS	-	expression tag	UNP Q08AH3
A	11	HIS	-	expression tag	UNP Q08AH3
A	12	HIS	-	expression tag	UNP Q08AH3
A	13	HIS	-	expression tag	UNP Q08AH3
A	14	HIS	-	expression tag	UNP Q08AH3
A	15	HIS	-	expression tag	UNP Q08AH3
A	16	SER	-	expression tag	UNP Q08AH3
A	17	SER	-	expression tag	UNP Q08AH3
A	18	GLY	-	expression tag	UNP Q08AH3
A	19	VAL	-	expression tag	UNP Q08AH3
A	20	ASP	-	expression tag	UNP Q08AH3
A	21	LEU	-	expression tag	UNP Q08AH3
A	22	GLY	-	expression tag	UNP Q08AH3
A	23	THR	-	expression tag	UNP Q08AH3
A	24	GLU	-	expression tag	UNP Q08AH3
A	25	ASN	-	expression tag	UNP Q08AH3
A	26	LEU	-	expression tag	UNP Q08AH3
A	27	TYR	-	expression tag	UNP Q08AH3
A	28	PHE	-	expression tag	UNP Q08AH3
A	29	GLN	-	expression tag	UNP Q08AH3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	SER	-	expression tag	UNP Q08AH3
A	31	MET	-	expression tag	UNP Q08AH3
A	64	PRO	LEU	engineered mutation	UNP Q08AH3
A	463	ASP	ASN	conflict	UNP Q08AH3
B	8	MET	-	expression tag	UNP Q08AH3
B	9	GLY	-	expression tag	UNP Q08AH3
B	10	HIS	-	expression tag	UNP Q08AH3
B	11	HIS	-	expression tag	UNP Q08AH3
B	12	HIS	-	expression tag	UNP Q08AH3
B	13	HIS	-	expression tag	UNP Q08AH3
B	14	HIS	-	expression tag	UNP Q08AH3
B	15	HIS	-	expression tag	UNP Q08AH3
B	16	SER	-	expression tag	UNP Q08AH3
B	17	SER	-	expression tag	UNP Q08AH3
B	18	GLY	-	expression tag	UNP Q08AH3
B	19	VAL	-	expression tag	UNP Q08AH3
B	20	ASP	-	expression tag	UNP Q08AH3
B	21	LEU	-	expression tag	UNP Q08AH3
B	22	GLY	-	expression tag	UNP Q08AH3
B	23	THR	-	expression tag	UNP Q08AH3
B	24	GLU	-	expression tag	UNP Q08AH3
B	25	ASN	-	expression tag	UNP Q08AH3
B	26	LEU	-	expression tag	UNP Q08AH3
B	27	TYR	-	expression tag	UNP Q08AH3
B	28	PHE	-	expression tag	UNP Q08AH3
B	29	GLN	-	expression tag	UNP Q08AH3
B	30	SER	-	expression tag	UNP Q08AH3
B	31	MET	-	expression tag	UNP Q08AH3
B	64	PRO	LEU	engineered mutation	UNP Q08AH3
B	463	ASP	ASN	conflict	UNP Q08AH3
C	8	MET	-	expression tag	UNP Q08AH3
C	9	GLY	-	expression tag	UNP Q08AH3
C	10	HIS	-	expression tag	UNP Q08AH3
C	11	HIS	-	expression tag	UNP Q08AH3
C	12	HIS	-	expression tag	UNP Q08AH3
C	13	HIS	-	expression tag	UNP Q08AH3
C	14	HIS	-	expression tag	UNP Q08AH3
C	15	HIS	-	expression tag	UNP Q08AH3
C	16	SER	-	expression tag	UNP Q08AH3
C	17	SER	-	expression tag	UNP Q08AH3
C	18	GLY	-	expression tag	UNP Q08AH3
C	19	VAL	-	expression tag	UNP Q08AH3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	20	ASP	-	expression tag	UNP Q08AH3
C	21	LEU	-	expression tag	UNP Q08AH3
C	22	GLY	-	expression tag	UNP Q08AH3
C	23	THR	-	expression tag	UNP Q08AH3
C	24	GLU	-	expression tag	UNP Q08AH3
C	25	ASN	-	expression tag	UNP Q08AH3
C	26	LEU	-	expression tag	UNP Q08AH3
C	27	TYR	-	expression tag	UNP Q08AH3
C	28	PHE	-	expression tag	UNP Q08AH3
C	29	GLN	-	expression tag	UNP Q08AH3
C	30	SER	-	expression tag	UNP Q08AH3
C	31	MET	-	expression tag	UNP Q08AH3
C	64	PRO	LEU	engineered mutation	UNP Q08AH3
C	463	ASP	ASN	conflict	UNP Q08AH3

- Molecule 2 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: $C_{10}H_{14}N_5O_7P$).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
2	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	1	Total 1	Mg 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	256	Total 256	O 256	0	0
4	B	234	Total 234	O 234	0	0
4	C	165	Total 165	O 165	0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACYL-COENZYME A SYNTHETASE ACSM2A, MITOCHONDRIAL

• Molecule 1: ACYL-COENZYME A SYNTHETASE ACSM2A, MITOCHONDRIAL

• Molecule 1: ACYL-COENZYME A SYNTHETASE ACSM2A, MITOCHONDRIAL

I395 I410 G411 I412 F423 N429 L443 I449 K450 R461 I465 Y471 E484 V489 E490 P497 F508 V509 V510 P537 L550 P551 K552 L563 W568 K569 MET SER GLY LYS ALA ARG ALA GLN

4   Data and refinement statistics ⓘ

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α, β, γ	97.73Å 97.73Å 384.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.16 – 2.45 38.00 – 2.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.16-2.45) 98.8 (38.00-2.45)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.30 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.243 0.208 , 0.254	Depositor DCC
R_{free} test set	3451 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å^2)	25.9	Xtriage
Anisotropy	0.111	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 22.0	EDS
L-test for twinning[2]	$< |L| > = 0.45, < L^2 > = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.93	EDS
Total number of atoms	13191	wwPDB-VP
Average B, all atoms (Å^2)	2.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.70% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$|Z|$>5	RMSZ	#$|Z|$>5
1	A	0.66	1/4287 (0.0%)	0.70	0/5817
1	B	0.60	0/4222	0.67	0/5735
1	C	0.60	2/4281 (0.0%)	0.65	0/5817
All	All	0.62	3/12790 (0.0%)	0.67	0/17369

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	471	TYR	CD1-CE1	-5.67	1.30	1.39
1	A	271	CYS	CB-SG	-5.37	1.73	1.81
1	C	484	GLU	CG-CD	5.30	1.59	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	470	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4175	0	4171	53	0
1	B	4120	0	4085	45	0
1	C	4171	0	4127	51	0
2	A	23	0	12	0	0
2	B	23	0	12	0	0
2	C	23	0	12	0	0
3	C	1	0	0	0	0
4	A	256	0	0	13	1
4	B	234	0	0	12	1
4	C	165	0	0	6	0
All	All	13191	0	12419	149	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:GLU:HG2	4:A:2067:HOH:O	1.43	1.14
1:C:371:MET:HE3	1:C:387:ALA:HB2	1.40	1.03
1:A:368:LEU:HD23	1:A:371:MET:HE1	1.44	0.99
1:A:151[B]:MET:HE2	1:A:230:MET:HG3	1.48	0.91
1:C:368:LEU:HD21	1:C:371:MET:HE2	1.55	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2144:HOH:O	4:B:2166:HOH:O[7_555]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	537/570 (94%)	524 (98%)	13 (2%)	0	100	100
1	B	533/570 (94%)	518 (97%)	15 (3%)	0	100	100
1	C	539/570 (95%)	522 (97%)	16 (3%)	1 (0%)	47	57
All	All	1609/1710 (94%)	1564 (97%)	44 (3%)	1 (0%)	51	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	36	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	453/485 (93%)	445 (98%)	8 (2%)	59	71
1	B	441/485 (91%)	434 (98%)	7 (2%)	62	74
1	C	448/485 (92%)	438 (98%)	10 (2%)	52	64
All	All	1342/1455 (92%)	1317 (98%)	25 (2%)	57	69

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	552	LYS
1	C	163	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	429	ASN
1	C	138	ILE
1	C	164	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	268	ASN
1	B	332	GLN
1	C	400	ASN
1	C	268	ASN
1	A	268	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#|Z| > 2$	Counts	RMSZ	$\#|Z| > 2$
2	AMP	B	1570	-	22,25,25	1.03	1 (4%)	25,38,38	1.44	5 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	#|Z| > 2	Counts	RMSZ	#|Z| > 2
2	AMP	A	1570	-	22,25,25	1.06	1 (4%)	25,38,38	1.35	2 (8%)
2	AMP	C	1571	-	22,25,25	1.00	0	25,38,38	1.28	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	B	1570	-	-	0/6/26/26	0/3/3/3
2	AMP	A	1570	-	-	0/6/26/26	0/3/3/3
2	AMP	C	1571	-	-	2/6/26/26	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1570	AMP	C5-C4	2.50	1.47	1.40
2	B	1570	AMP	C5-C4	2.42	1.47	1.40

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1570	AMP	N3-C2-N1	-4.10	122.27	128.68
2	A	1570	AMP	C4-C5-N7	-3.59	105.65	109.40
2	A	1570	AMP	N3-C2-N1	-3.20	123.68	128.68
2	C	1571	AMP	N3-C2-N1	-3.14	123.78	128.68
2	B	1570	AMP	O3P-P-O2P	2.39	116.77	107.64

There are no chirality outliers.

All (2) torsion outliers are listed below:

| Mol	Chain	Res	Type	Atoms
2	C	1571	AMP	C3'-C4'-C5'-O5'
2	C	1571	AMP	C5'-O5'-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand AMP B 1570

Bond lengths

Bond angles

Torsions

Rings

Ligand AMP A 1570

Bond lengths

Bond angles

Torsions

Rings

Ligand AMP C 1571

Bond lengths

Bond angles

Torsions

Rings

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	533/570 (93%)	-0.42	4 (0%)	86	86	2, 2, 4, 11	0
1	B	533/570 (93%)	-0.34	5 (0%)	84	85	2, 2, 4, 9	0
1	C	536/570 (94%)	-0.15	8 (1%)	73	71	2, 2, 4, 17	0
All	All	1602/1710 (93%)	-0.30	17 (1%)	80	80	2, 2, 4, 17	0

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	39	GLU	5.2
1	B	37	HIS	4.4
1	C	81	TRP	3.9
1	B	38	GLN	3.7
1	A	37	HIS	3.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($Å^2$)	Q<0.9
3	MG	C	1570	1/1	0.93	0.26	13,13,13,13	0
2	AMP	B	1570	23/23	0.98	0.11	2,2,7,14	0
2	AMP	C	1571	23/23	0.98	0.09	2,3,7,11	0
2	AMP	A	1570	23/23	0.98	0.11	2,2,3,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around AMP B 1570:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around AMP C 1571:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around AMP A 1570:

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

6.5 Other polymers ⓘ

There are no such residues in this entry.