



wwPDB X-ray Structure Validation Summary Report

Oct 5, 2023 – 07:20 AM EDT


PDB ID : 6VZD
Title : N-terminal domain of mouse surfactant protein B (K46E/R51E mutant) with bound lipid
Authors : Rapoport, T.A.; Bodnar, N.O.
Deposited on : 2020-02-28
Resolution : 1.88 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6494 atoms, of which 3176 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pulmonary surfactant-associated protein B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	79	1279	410	638	102	122	7	0	1	0
1	E	85	1369	435	683	113	131	7	0	2	0
1	B	78	1286	412	642	103	122	7	0	2	0
1	C	78	1266	407	629	101	122	7	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP P50405
A	46	GLU	LYS	engineered mutation	UNP P50405
A	51	GLU	ARG	engineered mutation	UNP P50405
E	1	SER	-	expression tag	UNP P50405
E	46	GLU	LYS	engineered mutation	UNP P50405
E	51	GLU	ARG	engineered mutation	UNP P50405
B	1	SER	-	expression tag	UNP P50405
B	46	GLU	LYS	engineered mutation	UNP P50405
B	51	GLU	ARG	engineered mutation	UNP P50405
C	1	SER	-	expression tag	UNP P50405
C	46	GLU	LYS	engineered mutation	UNP P50405
C	51	GLU	ARG	engineered mutation	UNP P50405

- Molecule 2 is (7Z,19R,22R)-25-amino-22-hydroxy-16,22-dioxo-17,21,23-trioxa-22lambda 5 -phosphapentacos-7-en-19-yl (9Z)-octadec-9-enoate (three-letter code: RXY) (formula: C₃₉H₇₄NO₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	A	1	244	78	146	2	16	2	0	1
2	A	1	122	39	73	1	8	1	0	0
2	E	1	122	39	73	1	8	1	0	0
2	B	1	122	39	73	1	8	1	0	0
2	B	1	244	78	146	2	16	2	0	1
2	B	1	122	39	73	1	8	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	94	Total	O	0	0
			94	94		
3	E	100	Total	O	0	0
			100	100		
3	B	80	Total	O	0	0
			80	80		
3	C	44	Total	O	0	0
			44	44		

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.79Å 66.12Å 59.62Å 90.00° 97.37° 90.00°	Depositor
Resolution (Å)	59.13 – 1.88	Depositor
% Data completeness (in resolution range)	91.8 (59.13-1.88)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.88Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.170 , 0.217	Depositor
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.001	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6494	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RXY	A	102	-	48,48,48	1.07	3 (6%)	51,53,53	0.90	2 (3%)
2	RXY	E	201	-	48,48,48	1.05	3 (6%)	51,53,53	0.90	2 (3%)
2	RXY	B	403	-	48,48,48	1.09	3 (6%)	51,53,53	0.91	2 (3%)
2	RXY	B	401	-	48,48,48	1.05	3 (6%)	51,53,53	0.82	2 (3%)
2	RXY	B	402[B]	-	48,48,48	1.07	3 (6%)	51,53,53	0.84	2 (3%)
2	RXY	A	101[A]	-	48,48,48	1.08	3 (6%)	51,53,53	0.84	2 (3%)
2	RXY	A	101[B]	-	48,48,48	1.06	3 (6%)	51,53,53	0.79	2 (3%)
2	RXY	B	402[A]	-	48,48,48	1.09	3 (6%)	51,53,53	0.93	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RXY	A	102	-	-	32/52/52/52	-
2	RXY	E	201	-	-	21/52/52/52	-
2	RXY	B	403	-	-	16/52/52/52	-
2	RXY	B	401	-	-	26/52/52/52	-
2	RXY	B	402[B]	-	-	26/52/52/52	-
2	RXY	A	101[A]	-	-	21/52/52/52	-
2	RXY	A	101[B]	-	-	24/52/52/52	-
2	RXY	B	402[A]	-	-	21/52/52/52	-

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	403	RXY	OAU-CAT	3.36	1.43	1.33
2	B	402[A]	RXY	OAU-CAT	3.28	1.42	1.33
2	B	401	RXY	OAU-CAT	3.28	1.42	1.33
2	A	102	RXY	OAU-CAT	3.26	1.42	1.33
2	A	101[A]	RXY	OAU-CAT	3.26	1.42	1.33

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	402[A]	RXY	OBW-CBV-CBU	4.10	120.34	111.50
2	A	102	RXY	OBW-CBV-CBU	4.08	120.30	111.50
2	E	201	RXY	OBW-CBV-CBU	4.03	120.19	111.50
2	A	101[A]	RXY	OBW-CBV-CBU	4.02	120.16	111.50
2	B	403	RXY	OBW-CBV-CBU	4.01	120.13	111.50

There are no chirality outliers.


5 of 187 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
2	A	101[A]	RXY	CBB-OBA-PAZ-OAY
2	A	101[A]	RXY	CBB-OBA-PAZ-OBX
2	A	101[A]	RXY	CBB-OBA-PAZ-OAB
2	A	101[A]	RXY	OBA-CBB-CBC-NBD
2	A	101[B]	RXY	CBB-OBA-PAZ-OAB




There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.