



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 5, 2023 – 12:31 AM EDT

PDB ID : 6VZY  
Title : Crystal structure of SznF from *Streptomyces achromogenes* var. *streptozoticus* NRRL 2697 with a diiron(II) central domain cofactor  
Authors : McBride, M.J.; Boal, A.K.  
Deposited on : 2020-02-28  
Resolution : 1.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**  
Xtrriage (Phenix) : 1.13  
EDS : **FAILED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.66 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7916 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cupin domain-containing diiron protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	464	3765	2395	653	705	12	0	0	0
1	B	460	3734	2377	645	701	11	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP A0A411MR89
A	-18	GLY	-	expression tag	UNP A0A411MR89
A	-17	SER	-	expression tag	UNP A0A411MR89
A	-16	SER	-	expression tag	UNP A0A411MR89
A	-15	HIS	-	expression tag	UNP A0A411MR89
A	-14	HIS	-	expression tag	UNP A0A411MR89
A	-13	HIS	-	expression tag	UNP A0A411MR89
A	-12	HIS	-	expression tag	UNP A0A411MR89
A	-11	HIS	-	expression tag	UNP A0A411MR89
A	-10	HIS	-	expression tag	UNP A0A411MR89
A	-9	SER	-	expression tag	UNP A0A411MR89
A	-8	SER	-	expression tag	UNP A0A411MR89
A	-7	GLY	-	expression tag	UNP A0A411MR89
A	-6	LEU	-	expression tag	UNP A0A411MR89
A	-5	VAL	-	expression tag	UNP A0A411MR89
A	-4	PRO	-	expression tag	UNP A0A411MR89
A	-3	ARG	-	expression tag	UNP A0A411MR89
A	-2	GLY	-	expression tag	UNP A0A411MR89
A	-1	SER	-	expression tag	UNP A0A411MR89
A	0	HIS	-	expression tag	UNP A0A411MR89
B	-19	MET	-	expression tag	UNP A0A411MR89
B	-18	GLY	-	expression tag	UNP A0A411MR89
B	-17	SER	-	expression tag	UNP A0A411MR89
B	-16	SER	-	expression tag	UNP A0A411MR89
B	-15	HIS	-	expression tag	UNP A0A411MR89

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	expression tag	UNP A0A411MR89
B	-13	HIS	-	expression tag	UNP A0A411MR89
B	-12	HIS	-	expression tag	UNP A0A411MR89
B	-11	HIS	-	expression tag	UNP A0A411MR89
B	-10	HIS	-	expression tag	UNP A0A411MR89
B	-9	SER	-	expression tag	UNP A0A411MR89
B	-8	SER	-	expression tag	UNP A0A411MR89
B	-7	GLY	-	expression tag	UNP A0A411MR89
B	-6	LEU	-	expression tag	UNP A0A411MR89
B	-5	VAL	-	expression tag	UNP A0A411MR89
B	-4	PRO	-	expression tag	UNP A0A411MR89
B	-3	ARG	-	expression tag	UNP A0A411MR89
B	-2	GLY	-	expression tag	UNP A0A411MR89
B	-1	SER	-	expression tag	UNP A0A411MR89
B	0	HIS	-	expression tag	UNP A0A411MR89

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Fe 3 3	0	0
2	B	3	Total Fe 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	202	Total O 202 202	0	0
3	B	209	Total O 209 209	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

### 3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.95Å 105.85Å 150.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.42 – 1.66	Depositor
% Data completeness (in resolution range)	96.1 (36.42-1.66)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 1.66Å)	Xtriage
Refinement program	REFMAC 1.14_3260, PHENIX 1.14_3260	Depositor
R, $R_{free}$	0.190 , 0.222	Depositor
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.044	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7916	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.15 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3159e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality [i](#)

### 4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.3 Torsion angles [i](#)

#### 4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

#### 4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

### 4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 4.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

#### 4.7 Other polymers [i](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data

### 5.1 Protein, DNA and RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.2 Non-standard residues in protein, DNA, RNA chains

EDS failed to run properly - this section is therefore empty.

### 5.3 Carbohydrates

EDS failed to run properly - this section is therefore empty.

### 5.4 Ligands

EDS failed to run properly - this section is therefore empty.

### 5.5 Other polymers

EDS failed to run properly - this section is therefore empty.