



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 09:11 pm BST

PDB ID : 1W2X
Title : Crystal structure of the carboxyltransferase domain of acetyl- coenzyme A
carboxylase in complex with CP-640186
Authors : Zhang, H.; Tweel, B.; Li, J.; Tong, L.
Deposited on : 2004-07-09
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	758	 3% 57% 31% 10%
1	B	758	 4% 52% 35% 11%
1	C	758	 3% 53% 31% 12%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 16588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYL-COA CARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	682	Total	C	N	O	S	0	0	1
			5425	3459	931	1016	19			
1	B	676	Total	C	N	O	S	0	0	1
			5377	3427	924	1007	19			
1	C	666	Total	C	N	O	S	0	0	1
			5299	3374	913	993	19			

- Molecule 2 is (3R)-1'-(9-ANTHRYLCARBONYL)-3-(MORPHOLIN-4-YLCARBONYL)-1,4'-BIPIPERIDINE (three-letter code: RCP) (formula: C₃₀H₃₅N₃O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total	C	N	O	0	0
			36	30	3	3		
2	B	1	Total	C	N	O	0	0
			36	30	3	3		
2	C	1	Total	C	N	O	0	0
			36	30	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	147	Total 147	O 147	0	0
3	B	128	Total 128	O 128	0	0
3	C	104	Total 104	O 104	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ACETYL-COA CARBOXYLASE




- Molecule 1: ACETYL-COA CARBOXYLASE





● Molecule 1: ACETYL-COA CARBOXYLASE



GLU	R2128	ASP	L1950	W1845	V1740
VAL	E2132	LYS	W1953	E1848	R1741
ILE	R1954	TYR	R1954	G1849	L1742
LYS	I2135	ARG	G1955	G1849	G1743
MET	I2136	GLU	R1961	E1851	Q1744
LEU	R2137	LEU	Q1960	T1852	R1745
THR	R2138	ARG	R1961	E1853	A1746
SER	H2141	SER	V1967	S1854	I1747
ASP	Q2142	GLN	L1968	G1855	Q1752
ASP	V2143	LEU	L1978	Y1858	I1755
LYS	G2144	SER	Y1979	L1756	L1756
GLU	E2145	ASN	D1980	T1757	T1757
LYS	A2146	LYS	Y1981	G1758	A1759
LEU	R2151	LEU	K1982	P1760	P1760
LEU	I2152	ALA	Q1983	K1764	K1764
THR	A2153	PRO	V1984	V1770	V1770
LEU	R2154	GLU	I1985	Y1771	Y1771
LEU	I2155	VAL	L1986	T1772	T1772
LYS	P2160	HIS	I1986	S1773	S1773
	V2163	GLN	P1991	M1774	M1774
	D2164	ILE	T1992	L1775	L1775
	R2169	LEU	G1993	Q1776	Q1776
	Q2170	ALA	E1994	T1780	T1780
	R2180	LEU	L1995	Q1781	Q1781
	K2185	ASP	R1996	I1782	I1782
	L2189	ARG	R1997	M1783	M1783
LEU	K2190	GLU	G1997	Y1784	Y1784
GLU	LEU	GLU	S1999	M1785	M1785
SER	SER	ARG	W2000	G1787	G1787
PHE	PHE	ARG	V2001	V1788	V1788
ALA	ALA	ARG	D2004	T1792	T1792
GLN	GLN	GLU	P2005	K1802	K1802
ASP	ASP	ARG	T2006	R1814	R1814
LEU	LEU	ASP	D2017	N1815	N1815
ALA	ALA	ASP	L2025	V1818	V1818
LYS	LYS	ASP	Q2028	L1821	L1821
LYS	LYS	ASP	V2031	K1824	K1824
ILE	ILE	ASP	G2032	W1827	W1827
ARG	ARG	ASP	I2033	D1828	D1828
SER	SER	ASP	K2034	R1829	R1829
ASP	ASP	ASP	F2035	F1833	F1833
ASP	ASP	ASP	R2036	D1838	D1838
HIS	HIS	ASP	R2037	F1838	F1838
ASP	ASP	ASP	L2041	K1824	K1824
ASN	ASN	ASP	D2042	W1827	W1827
ALA	ALA	ASP	T2043	D1828	D1828
ILE	ILE	ASP	W2044	R1829	R1829
ASP	ASP	ASP	E2114	F1833	F1833
GLY	GLY	ASP	W2115	D1838	D1838
LEU	LEU	ASP	W2124	F1833	F1833
SER	SER	ASP		D1838	D1838

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	246.59Å 124.60Å 145.60Å 90.00° 93.85° 90.00°	Depositor
Resolution (Å)	29.66 – 2.80 29.66 – 2.78	Depositor EDS
% Data completeness (in resolution range)	85.7 (29.66-2.80) 88.7 (29.66-2.78)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.76Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.197 , 0.234 0.200 , 0.238	Depositor DCC
R_{free} test set	10186 reflections (9.92%)	wwPDB-VP
Wilson B-factor (Å ²)	48.3	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16588	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: RCP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/5547	0.61	0/7516
1	B	0.39	0/5498	0.62	0/7451
1	C	0.39	0/5416	0.60	0/7337
All	All	0.39	0/16461	0.61	0/22304

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5425	0	5365	209	0
1	B	5377	0	5316	236	0
1	C	5299	0	5234	248	0
2	A	36	0	35	4	0
2	B	36	0	35	2	0
2	C	36	0	35	3	0
3	A	147	0	0	4	0
3	B	128	0	0	7	0
3	C	104	0	0	1	0
All	All	16588	0	16020	669	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 669 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1730:CYS:HA	1:B:1752:GLN:HE21	1.21	1.05
1:A:1936:ILE:HG12	1:A:1947:MET:HE1	1.41	1.02
1:C:1772:THR:H	1:C:1776:GLN:NE2	1.59	1.01
1:C:1773:SER:H	1:C:1776:GLN:HE21	1.09	1.00
1:B:1815:ASN:H	1:B:1944:GLN:HE22	0.98	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	678/758 (89%)	618 (91%)	52 (8%)	8 (1%)	13	39
1	B	672/758 (89%)	611 (91%)	51 (8%)	10 (2%)	10	33
1	C	662/758 (87%)	588 (89%)	64 (10%)	10 (2%)	10	33
All	All	2012/2274 (88%)	1817 (90%)	167 (8%)	28 (1%)	11	34

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2145	GLU
1	B	2189	LEU
1	C	1644	ASN
1	A	1684	GLY
1	A	1997	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	577/648 (89%)	548 (95%)	29 (5%)	24	56
1	B	572/648 (88%)	534 (93%)	38 (7%)	16	44
1	C	563/648 (87%)	522 (93%)	41 (7%)	14	38
All	All	1712/1944 (88%)	1604 (94%)	108 (6%)	18	46

5 of 108 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1791	LEU
1	B	2035	PHE
1	C	1980	ASP
1	B	1797	LEU
1	B	1909	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 79 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1748	GLN
1	B	2028	GLN
1	C	2011	GLN
1	B	1752	GLN
1	B	1909	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RCP	C	3000	-	41,41,41	1.03	1 (2%)	58,58,58	1.30	6 (10%)
2	RCP	B	3000	-	41,41,41	1.03	1 (2%)	58,58,58	1.27	6 (10%)
2	RCP	A	3000	-	41,41,41	1.03	1 (2%)	58,58,58	1.27	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RCP	C	3000	-	-	4/20/48/48	0/6/6/6
2	RCP	B	3000	-	-	4/20/48/48	0/6/6/6
2	RCP	A	3000	-	-	0/20/48/48	0/6/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3000	RCP	C14-C15	3.84	1.53	1.50
2	B	3000	RCP	C14-C15	3.76	1.53	1.50
2	A	3000	RCP	C14-C15	3.73	1.53	1.50

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	3000	RCP	C27-N26-C29	4.31	120.92	112.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	3000	RCP	C27-N26-C29	3.99	120.31	112.62
2	A	3000	RCP	C16-N15-C18	3.72	119.78	112.62
2	A	3000	RCP	C27-N26-C29	3.71	119.76	112.62
2	B	3000	RCP	C16-N15-C18	3.60	119.56	112.62

There are no chirality outliers.

5 of 8 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	C	3000	RCP	O26-C26-N26-C29
2	C	3000	RCP	C25-C26-N26-C29
2	B	3000	RCP	O26-C26-N26-C29
2	B	3000	RCP	C25-C26-N26-C29
2	C	3000	RCP	O26-C26-N26-C27


There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	3000	RCP	3	0
2	B	3000	RCP	2	0
2	A	3000	RCP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	682/758 (89%)	-0.24	21 (3%) 49 39	22, 47, 98, 99	0
1	B	676/758 (89%)	-0.20	28 (4%) 37 27	24, 49, 99, 99	0
1	C	666/758 (87%)	-0.27	21 (3%) 47 37	22, 49, 99, 99	0
All	All	2024/2274 (89%)	-0.23	70 (3%) 44 34	22, 48, 99, 99	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2082	LEU	6.2
1	C	1644	ASN	5.8
1	B	1492	TRP	5.2
1	B	1680	THR	5.2
1	B	1683	ASN	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	RCP	C	3000	36/36	0.90	0.24	73,87,94,95	0
2	RCP	B	3000	36/36	0.94	0.20	58,63,71,71	0
2	RCP	A	3000	36/36	0.94	0.17	60,68,69,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around RCP B 3000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around RCP A 3000:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.