



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2023 – 04:52 AM EDT

PDB ID : 3W6N
Title : Crystal structure of human Dlp1 in complex with GMP-PN.Pi
Authors : Kishida, H.; Sugio, S.
Deposited on : 2013-02-17
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	 10% 83% 9% • 7%
1	B	364	 5% 87% 6% • 6%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5876 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dynamin-1-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	Total	C	N	O	S	0	0	0
			2639	1662	471	497	9			
1	B	342	Total	C	N	O	S	0	0	0
			2682	1688	480	505	9			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP O00429
A	0	PRO	-	expression tag	UNP O00429
A	330	HIS	-	linker	UNP O00429
A	331	GLY	-	linker	UNP O00429
A	332	THR	-	linker	UNP O00429
A	333	ASP	-	linker	UNP O00429
A	334	SER	-	linker	UNP O00429
B	-1	GLY	-	expression tag	UNP O00429
B	0	PRO	-	expression tag	UNP O00429
B	330	HIS	-	linker	UNP O00429
B	331	GLY	-	linker	UNP O00429
B	332	THR	-	linker	UNP O00429
B	333	ASP	-	linker	UNP O00429
B	334	SER	-	linker	UNP O00429

- Molecule 2 is AMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNH) (formula: C₁₀H₁₆N₆O₁₀P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	28	10	6	10	2	0	0
2	B	1	28	10	6	10	2	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O P		
3	A	1	5	4 1	0	0
3	B	1	5	4 1	0	0


- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Ca 1 1	0	0

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C O 13 8 5	0	0
6	B	1	Total C O 13 8 5	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	203	Total O 203 203	0	0
7	B	257	Total O 257 257	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dynammin-1-like protein



- Molecule 1: Dynammin-1-like protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.34Å 109.02Å 128.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 46.45 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-2.00) 99.9 (46.45-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.97 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.194 , 0.235 0.195 , 0.235	Depositor DCC
R_{free} test set	2517 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtrriage
Anisotropy	0.191	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5876	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.0355e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PO4, GNH, PG4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2673	0.58	0/3616
1	B	0.45	0/2716	0.61	1/3673 (0.0%)
All	All	0.44	0/5389	0.59	1/7289 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	318	LEU	CA-CB-CG	5.89	128.85	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2639	0	2749	22	0
1	B	2682	0	2794	12	0
2	A	28	0	14	1	0
2	B	28	0	14	3	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	1	0	0	0	0
6	B	26	0	36	0	0
7	A	203	0	0	2	0
7	B	257	0	0	2	0
All	All	5876	0	5607	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:SER:HA	2:B:801:GNH:HN32	1.53	0.72
2:B:801:GNH:N3B	7:B:1070:HOH:O	2.25	0.67
1:B:22:ILE:HD11	1:B:307:ARG:HG2	1.80	0.64
1:A:35:SER:HA	2:A:801:GNH:HN32	1.65	0.61
1:A:725:GLN:HE21	1:A:725:GLN:HA	1.68	0.58
1:A:0:PRO:HG2	1:A:44:SER:O	2.02	0.58
1:A:95:HIS:CD2	1:A:96:THR:HG23	2.39	0.57
1:B:260:ASP:OD1	1:B:263:ARG:NH2	2.39	0.56
1:A:320:ASN:HB2	1:A:710:ARG:HE	1.72	0.55
1:A:50:LEU:H	1:A:50:LEU:HD12	1.73	0.53
1:A:1:MET:HG3	1:A:286:ALA:HB1	1.91	0.53
1:A:201:ARG:HD3	7:A:961:HOH:O	2.09	0.52
1:A:725:GLN:HA	1:A:725:GLN:NE2	2.25	0.51
1:A:95:HIS:CE1	1:A:114:GLU:CG	2.93	0.51
1:A:176:ASN:ND2	1:A:178:ASN:H	2.09	0.51
1:A:95:HIS:CE1	1:A:114:GLU:HG3	2.46	0.51
1:B:22:ILE:CD1	1:B:307:ARG:HG2	2.43	0.48
1:B:61:ARG:NH1	1:B:114:GLU:HG3	2.30	0.47
1:B:85:VAL:HG12	1:B:87:ALA:H	1.80	0.46
1:B:11:LEU:HD13	1:B:730:ILE:HD13	1.98	0.45
1:A:176:ASN:HD22	1:A:177:PRO:HD2	1.82	0.45
1:B:140:VAL:HG23	1:B:142:LEU:HB2	1.98	0.44
1:A:294:MET:HB2	1:A:294:MET:HE2	1.90	0.44
1:B:35:SER:HA	2:B:801:GNH:N3B	2.27	0.44
1:A:197:LEU:O	1:A:201:ARG:HG3	2.18	0.44
1:A:15:PHE:CD1	1:A:22:ILE:HG13	2.53	0.42
1:A:15:PHE:CE1	1:A:304:LEU:HD11	2.53	0.42
1:A:43:GLU:HB3	1:A:51:LEU:HD12	2.01	0.42
1:B:86:GLU:H	1:B:86:GLU:HG2	1.67	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:LEU:HB2	7:A:927:HOH:O	2.20	0.42
1:B:303:GLU:HG3	7:B:949:HOH:O	2.19	0.42
1:A:278:ASN:HD22	1:A:278:ASN:N	2.17	0.41
1:A:4:LEU:CD2	1:A:294:MET:HE1	2.50	0.41
1:A:278:ASN:HD22	1:A:278:ASN:H	1.69	0.41
1:B:70:VAL:HG21	1:B:135:PHE:HB3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/364 (91%)	319 (96%)	12 (4%)	0	100	100
1	B	336/364 (92%)	328 (98%)	8 (2%)	0	100	100
All	All	667/728 (92%)	647 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/320 (93%)	284 (95%)	14 (5%)	26	22
1	B	302/320 (94%)	289 (96%)	13 (4%)	29	26

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	600/640 (94%)	573 (96%)	27 (4%)	27	24

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	LEU
1	A	50	LEU
1	A	67	LEU
1	A	89	GLU
1	A	97	LYS
1	A	140	VAL
1	A	172	ARG
1	A	176	ASN
1	A	230	LEU
1	A	278	ASN
1	A	300	CYS
1	A	709	ARG
1	A	725	GLN
1	A	735	LEU
1	B	11	LEU
1	B	18	VAL
1	B	45	LEU
1	B	50	LEU
1	B	67	LEU
1	B	86	GLU
1	B	142	LEU
1	B	171	LEU
1	B	176	ASN
1	B	237	VAL
1	B	271	LYS
1	B	318	LEU
1	B	712	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	GLN
1	A	122	ASN
1	A	131	HIS
1	A	176	ASN
1	A	278	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	320	ASN
1	A	725	GLN
1	B	109	GLN
1	B	122	ASN
1	B	165	GLN
1	B	176	ASN
1	B	320	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 3 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GNH	B	801	4	24,30,30	1.96	6 (25%)	27,47,47	2.23	8 (29%)
6	PG4	B	805	-	12,12,12	0.39	0	11,11,11	0.46	0
3	PO4	B	802	4	4,4,4	0.92	0	6,6,6	1.28	1 (16%)
2	GNH	A	801	4	24,30,30	1.83	5 (20%)	27,47,47	2.12	7 (25%)
3	PO4	A	802	4	4,4,4	0.76	0	6,6,6	1.23	1 (16%)
6	PG4	B	806	-	12,12,12	0.53	0	11,11,11	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GNH	B	801	4	-	0/9/32/32	0/3/3/3
6	PG4	B	805	-	-	3/10/10/10	-
6	PG4	B	806	-	-	6/10/10/10	-
2	GNH	A	801	4	-	0/9/32/32	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	GNH	PB-O1B	6.06	1.55	1.46
2	B	801	GNH	PB-O1B	5.79	1.55	1.46
2	B	801	GNH	O6-C6	4.38	1.35	1.24
2	A	801	GNH	O6-C6	3.82	1.34	1.24
2	B	801	GNH	C5-C6	3.80	1.47	1.41
2	A	801	GNH	C5-C6	3.00	1.46	1.41
2	B	801	GNH	PB-O3A	2.43	1.62	1.59
2	B	801	GNH	O4'-C1'	2.15	1.44	1.41
2	A	801	GNH	O4'-C1'	2.09	1.44	1.41
2	A	801	GNH	PB-O2B	-2.02	1.51	1.56
2	B	801	GNH	C6-N1	2.00	1.36	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	GNH	C2-N3-C4	5.33	121.44	115.36
2	A	801	GNH	C2-N3-C4	5.30	121.41	115.36
2	B	801	GNH	N3-C2-N1	-5.08	120.45	127.22
2	A	801	GNH	N3-C2-N1	-4.78	120.85	127.22
2	B	801	GNH	C4-C5-C6	-4.40	116.59	120.80
2	A	801	GNH	C4-C5-C6	-4.25	116.74	120.80
2	B	801	GNH	O2B-PB-O1B	4.02	120.68	110.25
2	A	801	GNH	O2B-PB-O1B	3.50	119.32	110.25
2	B	801	GNH	C2-N1-C6	3.44	121.40	115.93
2	B	801	GNH	C5-C6-N1	-3.39	118.79	123.43
2	A	801	GNH	C2-N1-C6	3.18	120.99	115.93
2	A	801	GNH	C5-C6-N1	-3.06	119.24	123.43
2	B	801	GNH	N2-C2-N1	2.45	121.07	117.25
3	A	802	PO4	O4-P-O3	2.42	115.74	107.97
3	B	802	PO4	O4-P-O3	2.17	114.94	107.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	GNH	PB-O3A-PA	-2.12	125.84	132.56
2	B	801	GNH	C1'-N9-C4	-2.03	123.08	126.64

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	B	806	PG4	O2-C3-C4-O3
6	B	806	PG4	O4-C7-C8-O5
6	B	806	PG4	O3-C5-C6-O4
6	B	806	PG4	O1-C1-C2-O2
6	B	805	PG4	O1-C1-C2-O2
6	B	805	PG4	O4-C7-C8-O5
6	B	806	PG4	C5-C6-O4-C7
6	B	806	PG4	C3-C4-O3-C5
6	B	805	PG4	C8-C7-O4-C6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	GNH	3	0
2	A	801	GNH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	337/364 (92%)	0.38	37 (10%) 5 5	8, 21, 63, 78	0
1	B	342/364 (93%)	0.06	20 (5%) 23 22	7, 17, 45, 64	0
All	All	679/728 (93%)	0.22	57 (8%) 11 10	7, 19, 52, 78	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	321	SER	9.3
1	A	736	TRP	7.2
1	A	315	TYR	7.2
1	B	318	LEU	6.5
1	A	320	ASN	6.2
1	A	709	ARG	6.1
1	A	711	LYS	5.7
1	A	716	MET	5.4
1	A	73	GLU	5.3
1	A	307	ARG	5.2
1	A	714	ALA	4.8
1	A	717	LEU	4.8
1	A	313	ALA	4.7
1	A	710	ARG	4.6
1	B	319	LEU	4.6
1	A	715	ASP	4.5
1	B	711	LYS	4.5
1	A	312	ALA	4.2
1	A	97	LYS	4.1
1	B	710	ARG	4.0
1	A	311	LEU	3.9
1	B	87	ALA	3.9
1	B	709	ARG	3.8
1	B	736	TRP	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	55	THR	3.8
1	A	18	VAL	3.8
1	B	85	VAL	3.7
1	B	312	ALA	3.5
1	B	73	GLU	3.5
1	A	316	GLN	3.5
1	A	720	LEU	3.4
1	A	308	ILE	3.4
1	A	88	GLU	3.3
1	A	713	ALA	3.3
1	A	98	ASN	3.2
1	B	313	ALA	3.1
1	B	315	TYR	3.0
1	A	718	LYS	2.9
1	B	715	ASP	2.8
1	B	712	GLU	2.7
1	B	86	GLU	2.5
1	A	319	LEU	2.5
1	B	713	ALA	2.5
1	A	719	ALA	2.4
1	B	317	SER	2.4
1	A	303	GLU	2.4
1	A	21	ASP	2.4
1	B	56	GLY	2.3
1	A	24	GLN	2.3
1	A	310	VAL	2.3
1	A	712	GLU	2.2
1	A	317	SER	2.2
1	A	318	LEU	2.2
1	B	718	LYS	2.2
1	A	309	ASN	2.1
1	A	721	GLN	2.1
1	A	20	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	PG4	B	806	13/13	0.82	0.17	37,38,40,41	0
6	PG4	B	805	13/13	0.89	0.15	27,30,35,35	0
3	PO4	A	802	5/5	0.96	0.12	18,19,23,24	0
3	PO4	B	802	5/5	0.96	0.12	16,19,20,24	0
4	MG	A	803	1/1	0.97	0.08	5,5,5,5	0
2	GNH	A	801	28/28	0.98	0.10	5,11,12,14	0
4	MG	B	803	1/1	0.98	0.09	5,5,5,5	0
2	GNH	B	801	28/28	0.99	0.09	7,10,13,14	0
5	CA	B	804	1/1	1.00	0.08	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.