



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2023 – 09:34 PM EST

PDB ID : 5W6T
Title : Crystal structure of the A/Puerto Rico/8/1934 (H1N1) influenza virus hemagglutinin in complex with cyclic peptide CP151070 (P7)
Authors : Wilson, I.A.; Kadam, R.U.
Deposited on : 2017-06-16
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	
2	B	176	
3	F	12	
4	C	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	406	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	322	2542	1603	443	483	13	0	0	0


- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	171	1380	866	235	272	7	0	0	0

- Molecule 3 is a protein called ACE-PH8-ORN-MLE-GLU-TYR-ZCL-GLU-TRP-LEU-SER-9WV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	Cl	N	O			
3	F	12	115	79	2	15	19	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	C	2	28	16	2	10	0	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
5	A	1	14	8	1	5	0	0
5	A	1	14	8	1	5	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
6	A	1	6	3	3	0	0
6	A	1	6	3	3	0	0

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Cl 1 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	48	Total O 48 48	0	0
8	B	19	Total O 19 19	0	0
8	F	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin



- Molecule 2: Hemagglutinin



- Molecule 3: ACE-PH8-ORN-MLE-GLU-TYR-ZCL-GLU-TRP-LEU-SER-9WV



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, α , β , γ	164.33Å 164.33Å 164.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.08 – 2.59 41.08 – 2.59	Depositor EDS
% Data completeness (in resolution range)	92.2 (41.08-2.59) 92.2 (41.08-2.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.58Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.219 , 0.256 0.220 , 0.257	Depositor DCC
R_{free} test set	1072 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	55.4	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.026 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4174	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZCL, 9WV, MLE, ACE, ORN, PH8, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/2606	0.60	0/3544
2	B	0.46	0/1407	0.54	0/1891
3	F	2.68	8/59 (13.6%)	1.63	1/78 (1.3%)
All	All	0.59	8/4072 (0.2%)	0.61	1/5513 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	F	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	6	TYR	CB-CG	-7.53	1.40	1.51
3	F	9	TRP	CD2-CE2	-6.52	1.33	1.41
3	F	9	TRP	CG-CD2	-6.10	1.33	1.43
3	F	10	LEU	CA-C	-6.08	1.37	1.52
3	F	11	SER	CA-C	-5.74	1.38	1.52
3	F	5	GLU	CA-C	-5.50	1.38	1.52
3	F	9	TRP	CA-C	-5.42	1.38	1.52
3	F	6	TYR	CA-C	-5.18	1.39	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	9	TRP	CD1-CG-CD2	5.67	110.83	106.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	F	11	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2542	0	2469	36	0
2	B	1380	0	1310	14	0
3	F	115	0	88	3	0
4	C	28	0	25	0	0
5	A	28	0	26	0	0
6	A	12	0	15	7	0
7	B	1	0	0	0	0
8	A	48	0	0	1	0
8	B	19	0	0	0	0
8	F	1	0	0	0	0
All	All	4174	0	3933	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:50:ASN:O	2:B:54:THR:CG2	2.21	0.88
2:B:50:ASN:O	2:B:54:THR:HG23	1.72	0.86
1:A:73:ASN:ND2	6:A:406:GOL:O1	2.07	0.86
1:A:73:ASN:HD21	6:A:406:GOL:HO1	1.24	0.84
1:A:76:CYS:O	1:A:149:ARG:NH2	2.14	0.80
1:A:37:THR:OG1	1:A:38:HIS:ND1	2.15	0.76
1:A:37:THR:HG1	1:A:38:HIS:CE1	2.10	0.68
1:A:73:ASN:ND2	6:A:406:GOL:HO1	1.93	0.64
2:B:50:ASN:O	2:B:54:THR:HG22	1.97	0.64
1:A:38:HIS:CE1	2:B:21:TRP:HE1	2.17	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:ILE:HG22	1:A:289:ASN:O	2.01	0.61
1:A:95(A):GLY:H	6:A:406:GOL:H31	1.66	0.60
3:F:11:SER:O	3:F:12:9WV:NDV	2.36	0.59
1:A:71:LEU:O	1:A:148:TYR:HB3	2.07	0.55
1:A:103:ILE:HG13	1:A:233:TYR:CE2	2.42	0.54
1:A:83:ARG:NH1	8:A:503:HOH:O	2.40	0.53
1:A:43:LEU:HB2	1:A:314:LEU:HB2	1.92	0.52
1:A:92:ASN:OD1	1:A:92:ASN:N	2.42	0.52
1:A:301:THR:HB	1:A:305:CYS:SG	2.51	0.51
1:A:53:ARG:NH1	1:A:55:GLY:O	2.44	0.50
1:A:95:ASN:HD22	6:A:406:GOL:H12	1.75	0.50
1:A:117:SER:HB3	1:A:261:SER:OG	2.12	0.49
1:A:211:ARG:HG2	1:A:213:PHE:CZ	2.49	0.48
1:A:95:ASN:HA	6:A:406:GOL:H2	1.95	0.48
1:A:74:PRO:HB2	1:A:141:HIS:HB2	1.95	0.47
2:B:17:MET:SD	2:B:23:GLY:HA3	2.54	0.47
3:F:6:TYR:H	3:F:10:LEU:HD12	1.79	0.47
1:A:66:ILE:HG12	1:A:89:GLU:OE1	2.15	0.46
1:A:125(A):LYS:HE2	1:A:152:LEU:HD11	1.96	0.46
2:B:119:TYR:CE1	2:B:136:GLY:HA2	2.51	0.46
1:A:125(B):GLU:O	1:A:125(C):SER:HB2	2.16	0.45
2:B:49:THR:OG1	3:F:4:MLE:HN2	2.17	0.45
2:B:97:GLU:O	2:B:101:LEU:HG	2.16	0.45
1:A:15:ILE:HD11	2:B:122:VAL:HG21	1.99	0.44
1:A:316:MET:HE1	2:B:55:VAL:HG11	2.01	0.43
2:B:163:SER:H	2:B:163:SER:HG	1.54	0.43
1:A:202:VAL:HG13	1:A:247:ALA:HB2	2.00	0.43
1:A:89:GLU:O	1:A:269:THR:HA	2.19	0.43
2:B:170:ARG:O	2:B:171:GLU:HG2	2.19	0.42
1:A:220:ARG:NH1	1:A:228:GLY:HA2	2.34	0.42
1:A:98:TYR:O	1:A:232:TYR:OH	2.30	0.42
1:A:71:LEU:HD22	1:A:151:LEU:HD11	2.01	0.42
1:A:24:ASP:O	1:A:35:THR:HA	2.20	0.41
1:A:97:CYS:HA	6:A:406:GOL:H32	2.03	0.41
1:A:74:PRO:HG3	1:A:147:PHE:O	2.21	0.41
2:B:4:GLY:O	2:B:8:GLY:HA3	2.21	0.41
1:A:28:THR:HG22	2:B:104:ASN:HB3	2.03	0.41
1:A:37:THR:OG1	1:A:38:HIS:CE1	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/326 (98%)	307 (96%)	11 (3%)	2 (1%)	25	47
2	B	169/176 (96%)	163 (96%)	6 (4%)	0	100	100
3	F	6/12 (50%)	5 (83%)	0	1 (17%)	0	0
All	All	495/514 (96%)	475 (96%)	17 (3%)	3 (1%)	25	47

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125(C)	SER
3	F	11	SER
1	A	125(A)	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/289 (99%)	279 (98%)	6 (2%)	53	77
2	B	147/151 (97%)	146 (99%)	1 (1%)	84	94
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	438/446 (98%)	431 (98%)	7 (2%)	62	82

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	SER
1	A	102	PHE
1	A	156	GLU
1	A	261	SER
1	A	290	SER
1	A	302	ILE
2	B	54	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ORN	F	3	3	6,7,8	0.43	0	2,7,9	0.59	0
3	9WV	F	12	3	8,10,11	0.78	0	10,11,13	0.97	0
3	PH8	F	2	3	12,13,14	1.62	2 (16%)	10,15,17	1.24	1 (10%)
3	ZCL	F	7	3	12,13,14	1.46	1 (8%)	14,17,19	0.98	1 (7%)
3	MLE	F	4	3	7,8,9	0.56	0	6,9,11	1.46	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ORN	F	3	3	-	0/5/6/8	-
3	9WV	F	12	3	-	4/10/11/13	-
3	PH8	F	2	3	-	3/7/8/10	0/1/1/1
3	ZCL	F	7	3	-	0/5/6/8	0/1/1/1
3	MLE	F	4	3	-	0/5/8/10	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	2	PH8	CJ-CG	-4.97	1.37	1.51
3	F	7	ZCL	CB-CG	-4.88	1.39	1.51
3	F	2	PH8	CB-CA	-2.17	1.50	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	4	MLE	CN-N-CA	2.42	121.18	113.64
3	F	2	PH8	CI-CJ-CG	-2.34	104.86	113.68
3	F	7	ZCL	CE2-CZ-CLZ	2.01	122.44	118.41
3	F	4	MLE	CG-CB-CA	-2.00	110.34	115.34

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	2	PH8	C-CA-CB-CI
3	F	2	PH8	N-CA-CB-CI
3	F	12	9WV	O-C-CA-CB
3	F	12	9WV	CEH-CEG-NDV-CA
3	F	12	9WV	OEK-CEG-NDV-CA
3	F	12	9WV	CEG-CEH-CEI-NEJ
3	F	2	PH8	CA-CB-CI-CJ

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	12	9WV	1	0
3	F	4	MLE	1	0

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	1	4,1	14,14,15	0.29	0	17,19,21	0.58	0
4	NAG	C	2	4	14,14,15	0.28	0	17,19,21	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	A	406	1	5,5,5	0.36	0	5,5,5	0.78	0
5	NAG	A	404	1	14,14,15	0.52	0	17,19,21	0.99	1 (5%)
5	NAG	A	403	1	14,14,15	0.42	0	17,19,21	0.59	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	405	-	5,5,5	0.31	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	406	1	-	1/4/4/4	-
5	NAG	A	404	1	-	2/6/23/26	0/1/1/1
5	NAG	A	403	1	-	2/6/23/26	0/1/1/1
6	GOL	A	405	-	-	4/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	404	NAG	C1-O5-C5	3.54	116.99	112.19
5	A	403	NAG	C1-O5-C5	2.04	114.95	112.19

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	405	GOL	O1-C1-C2-C3
5	A	404	NAG	C4-C5-C6-O6
5	A	404	NAG	O5-C5-C6-O6
5	A	403	NAG	C4-C5-C6-O6
6	A	405	GOL	C1-C2-C3-O3
6	A	406	GOL	O1-C1-C2-C3
6	A	405	GOL	O1-C1-C2-O2
5	A	403	NAG	O5-C5-C6-O6
6	A	405	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	406	GOL	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/326 (98%)	-0.17	3 (0%) 84 82	25, 43, 77, 97	0
2	B	171/176 (97%)	0.49	13 (7%) 13 10	25, 62, 104, 118	0
3	F	6/12 (50%)	0.24	0 100 100	64, 73, 83, 87	0
All	All	499/514 (97%)	0.06	16 (3%) 47 40	25, 48, 97, 118	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	168	LEU	4.1
1	A	325	SER	4.1
2	B	141	TYR	3.5
2	B	65	ALA	3.1
2	B	62	GLN	2.8
1	A	92	ASN	2.8
2	B	164	GLU	2.6
2	B	61	ILE	2.6
2	B	162	TYR	2.5
2	B	157	TYR	2.5
2	B	166	SER	2.5
2	B	138	PHE	2.4
2	B	158	ASP	2.4
1	A	196	GLN	2.3
2	B	29	GLU	2.1
2	B	147	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9WV	F	12	11/12	0.85	0.22	93,96,102,102	0
3	ORN	F	3	8/9	0.91	0.17	54,62,80,93	0
3	ZCL	F	7	13/14	0.92	0.22	62,65,71,89	0
3	PH8	F	2	13/14	0.94	0.17	53,58,69,73	0
3	MLE	F	4	9/10	0.94	0.21	45,58,65,71	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	1	14/15	0.79	0.29	99,114,127,136	0
4	NAG	C	2	14/15	0.89	0.27	109,138,152,159	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	404	14/15	0.70	0.37	114,132,137,140	0
5	NAG	A	403	14/15	0.79	0.34	99,116,122,124	0
6	GOL	A	406	6/6	0.80	0.34	51,57,61,77	0
6	GOL	A	405	6/6	0.90	0.17	41,54,57,66	0
7	CL	B	201	1/1	0.99	0.35	32,32,32,32	1

6.5 Other polymers [i](#)

There are no such residues in this entry.