

Full wwPDB X-ray Structure Validation Report (i)

Aug 7, 2020 – 10:56 PM BST

PDB ID : 5W7E

Title: Murine acyloxyacyl hydrolase (AOAH), S262A mutant, with dimyristoyl phos-

phatidylcholine

Authors : Gorelik, A.; Illes, K.; Nagar, B.

Deposited on : 2017-06-19

Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned} ext{Similar resolution} \ (\# ext{Entries}, ext{resolution range}(ext{Å})) \end{aligned}$		
R_{free}	130704	4003 (1.86-1.82)		
Clashscore	141614	4233 (1.86-1.82)		
Ramachandran outliers	138981	4185 (1.86-1.82)		
Sidechain outliers	138945	4186 (1.86-1.82)		
RSRZ outliers	127900	3957 (1.86-1.82)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	562	88%	5%	7%			
1	В	562	86%	7%	6%			



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 18152 atoms, of which 8544 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Acyloxyacyl hydrolase.

Mol	Chain	Residues		${f Atoms}$				ZeroOcc	AltConf	Trace	
1	A	521	Total 8382	C 2699	H 4144	N 732	O 783	S 24	0	15	0
1	В	526	Total 8531	C 2740	H 4224	N 745	O 798	S 24	0	21	0

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	ASP	_	expression tag	UNP O35298
A	14	ARG	-	expression tag	UNP O35298
A	15	HIS	-	expression tag	UNP O35298
A	16	HIS	-	expression tag	UNP O35298
A	17	HIS	-	expression tag	UNP O35298
A	18	HIS	-	expression tag	UNP O35298
A	19	HIS	-	expression tag	UNP O35298
A	20	HIS	-	expression tag	UNP O35298
A	21	LYS	-	expression tag	UNP O35298
A	22	LEU	-	expression tag	UNP O35298
A	152	ARG	PRO	conflict	UNP O35298
A	184	VAL	ILE	$\operatorname{conflict}$	UNP O35298
A	262	ALA	SER	engineered mutation	UNP O35298
В	13	ASP	-	expression tag	UNP O35298
В	14	ARG	-	expression tag	UNP O35298
В	15	HIS	-	expression tag	UNP O35298
В	16	HIS	-	expression tag	UNP O35298
В	17	HIS	-	expression tag	UNP O35298
В	18	HIS	-	expression tag	UNP O35298
В	19	HIS	-	expression tag	UNP O35298
В	20	HIS	-	expression tag	UNP O35298
В	21	LYS	-	expression tag	UNP O35298
В	22	LEU	=	expression tag	UNP O35298
В	152	ARG	PRO	conflict	UNP O35298
В	184	VAL	ILE	conflict	UNP O35298




Continued from previous page...

Chain	Residue	Modelled	Actual	${f Comment}$	Reference
В	262	ALA	SER	engineered mutation	UNP O35298

• Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	3	Total Ca 3 3	0	0
2	A	3	Total Ca	0	0


• Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by author).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf			
3	Λ	1	Total	С	Н	N	О	0	0			
)	A	1	28	8	14	1	5	0	0			
3	Λ	Λ	Λ	Λ	1	Total	С	Н	N	О	0	0
3	A	1	28	8	14	1	5	U				
3	D	1	Total	С	Н	N	О	0	0			
3) D	1	28	8	14	1	5	0	U			
3	D	B 1	Total	С	Н	N	О	0	0			
3	Ď		28	8	14	1	5	0				

• Molecule 4 is (2S)-3-hydroxypropane-1,2-diyl ditetradecanoate (three-letter code: FAW) (formula: C₃₁H₆₀O₅) (labeled as "Ligand of Interest" by author).





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf	
4	Λ	Λ	1	Total	С	Н	О	0	0
4	А	1	96	31	60	5	0	U	
1	D	1	Total	С	Н	О	0	0	
4	4 B	1	96	31	60	5	U	U	

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	468	Total O 468 468	0	0
5	В	461	Total O 461 461	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acyloxyacyl hydrolase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	81.42Å 85.37Å 93.40Å	Depositor
a, b, c, α , β , γ	90.00° 103.64° 90.00°	Depositor
Resolution (Å)	40.07 - 1.83	Depositor
resolution (A)	40.07 - 1.83	EDS
% Data completeness	58.9 (40.07-1.83)	Depositor
(in resolution range)	83.0 (40.07-1.83)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.99 (at 1.83Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
D D.	0.157 , 0.198	Depositor
R, R_{free}	0.158 , 0.197	DCC
R_{free} test set	1948 reflections (2.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.4	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.42,63.3	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18152	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, FAW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.42	0/4392	0.56	0/5957	
1	В	0.43	0/4482	0.59	$2/6079 \ (0.0\%)$	
All	All	0.42	0/8874	0.58	$2/12036 \ (0.0\%)$	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	В	187	ASP	CB-CG-OD1	5.86	123.58	118.30
1	В	187	ASP	CB-CG-OD2	-5.40	113.44	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4238	4144	4153	20	1
1	В	4307	4224	4230	26	1
2	A	3	0	0	0	0
2	В	3	0	0	0	0
3	A	28	28	26	0	0
3	В	28	28	26	0	0
4	A	36	60	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
4	В	36	60	0	2	0
5	A	468	0	0	10	5
5	В	461	0	0	11	5
All	All	9608	8544	8435	46	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:314:GLU:OE1	5:A:701:HOH:O	2.00	0.78
1:B:325:ARG:NH2	1:B:547:GLN:OE1	2.18	0.76
1:A:130:GLU:OE2	5:A:702:HOH:O	2.04	0.75
1:A:87:GLN:OE1	5:A:703:HOH:O	2.08	0.71
1:B:438:ASN:OD1	5:B:701:HOH:O	2.09	0.70
1:A:422:PHE:HE1	1:A:518:PRO:HB3	1.65	0.61
1:B:451[A]:ARG:NE	5:B:706:HOH:O	2.29	0.58
1:B:278:GLN:NE2	1:B:278:GLN:O	2.37	0.58
1:A:344:ARG:NH2	1:A:384:GLU:O	2.38	0.57
1:B:275[B]:THR:HG21	5:B:944:HOH:O	2.06	0.54
1:A:510:ASP:OD2	1:A:541:ARG:NH2	2.38	0.54
1:B:144:ARG:O	1:B:145:ARG:HB2	2.08	0.54
1:B:257:ILE:HD13	1:B:353:LEU:HD21	1.93	0.51
1:B:361:HIS:HD2	5:B:1097:HOH:O	1.94	0.50
1:A:422:PHE:CE1	1:A:518:PRO:HB3	2.45	0.50
1:A:126:PRO:HG3	5:A:1067:HOH:O	2.13	0.49
1:B:429[A]:ASN:ND2	5:B:702:HOH:O	2.19	0.48
1:B:159:CYS:O	1:B:165:THR:HG22	2.13	0.48
1:B:451[A]:ARG:NH2	5:B:706:HOH:O	2.45	0.48
1:A:325:ARG:NH1	5:A:720:HOH:O	2.47	0.48
1:B:105[A]:VAL:HG23	5:B:920:HOH:O	2.17	0.45
1:A:418:PRO:HB3	1:A:502:PHE:CD2	2.52	0.45
1:A:275[A]:THR:HG21	5:A:947:HOH:O	2.16	0.45
1:B:262:ALA:HB3	4:B:606:FAW:CA1	2.47	0.44
1:A:127:LEU:HG	1:A:131:ALA:HB3	1.98	0.44
1:A:198:TYR:HA	1:A:201:ARG:O	2.16	0.44
1:A:161:LEU:HD12	1:A:164:LEU:HD12	2.00	0.44
1:A:451:ARG:HD3	5:A:1055:HOH:O	2.18	0.44
1:B:517:GLN:HG3	5:B:830:HOH:O	2.17	0.43
1:B:164:LEU:O	1:B:167:ILE:HG22	2.18	0.43



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}\;({ m \AA})$	$oxed{ ext{overlap } (ext{\AA}) }$
1:A:370:ILE:O	1:A:418:PRO:HD2	2.18	0.43
1:A:540:ASN:HB2	5:A:708:HOH:O	2.19	0.43
1:B:281:VAL:HG23	5:B:875:HOH:O	2.18	0.42
1:B:430:ARG:HD2	5:B:711:HOH:O	2.18	0.42
1:B:484:LYS:NZ	5:B:715:HOH:O	2.40	0.42
1:A:422:PHE:HE1	1:A:518:PRO:CB	2.30	0.41
1:B:534:ALA:O	1:B:538[B]:GLN:HG3	2.21	0.41
1:B:422:PHE:CE1	1:B:518:PRO:HB3	2.55	0.41
1:B:198:TYR:HA	1:B:201:ARG:O	2.21	0.41
1:B:428:HIS:NE2	1:B:429[A]:ASN:OD1	2.54	0.41
1:B:504:PHE:CE2	4:B:606:FAW:C33	3.04	0.40
1:B:272:GLU:HA	1:B:278:GLN:NE2	2.36	0.40
1:B:501:ASP:OD2	1:B:503:ALA:HB2	2.21	0.40
1:A:320[B]:HIS:HD2	5:A:905:HOH:O	2.05	0.40
1:A:451:ARG:NH2	1:A:463:SER:HB2	2.36	0.40
5:A:789:HOH:O	1:B:166:LYS:HE2	2.21	0.40

All (6) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap} & (ext{Å}) \end{aligned}$
5:A:1067:HOH:O	5:B:776:HOH:O[1_655]	1.98	0.22
5:A:1108:HOH:O	5:B:1077:HOH:O[1_655]	2.01	0.19
5:A:1115:HOH:O	5:B:1118:HOH:O[2_454]	2.06	0.14
1:A:238:LYS:HZ1	1:B:239:ASP:O[1_556]	1.48	0.12
5:A:987:HOH:O	5:B:1081:HOH:O[2_454]	2.10	0.10
5:A:1029:HOH:O	5:B:712:HOH:O[1_655]	2.13	0.07

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed Favoured Allowed		Outliers	Percentiles	
1	A	530/562~(94%)	517 (98%)	12 (2%)	1 (0%)	47 33
1	В	542/562~(96%)	529 (98%)	11 (2%)	2 (0%)	34 20
All	All	$1072/1124 \ (95\%)$	1046 (98%)	23 (2%)	3 (0%)	41 27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	371	GLY
1	В	371	GLY
1	В	183	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percent	iles
1	A	478/500 (96%)	477 (100%)	1 (0%)	93 9	2
1	В	$488/500 \ (98\%)$	483 (99%)	5 (1%)	76 6	8
All	All	$966/1000 \; (97\%)$	960 (99%)	6 (1%)	86 8	2

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	HIS
1	В	36	HIS
1	В	129	GLN
1	В	278	GLN
1	В	327	HIS
1	В	501	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

\mathbf{Mol}	Chain	${f Res}$	Type
1	В	278	GLN
1	В	438	ASN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	nd leng	ths	В	ond ang	les
WIOI	Type	Chain	res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	В	605	1	14,14,15	0.24	0	17,19,21	0.35	0
3	NAG	A	605	1	14,14,15	0.35	0	17,19,21	0.85	1 (5%)
4	FAW	A	606	-	35,35,35	0.90	4 (11%)	37,37,37	1.23	3 (8%)
4	FAW	В	606	-	35,35,35	0.87	4 (11%)	37,37,37	1.20	2 (5%)
3	NAG	В	604	1	14,14,15	0.25	0	17,19,21	0.83	1 (5%)
3	NAG	A	604	1	14,14,15	0.31	0	17,19,21	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	В	605	1	-	2/6/23/26	0/1/1/1
3	NAG	A	605	1	-	2/6/23/26	0/1/1/1
4	FAW	A	606	-	-	18/37/37/37	-



 $Continued\ from\ previous\ page...$

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FAW	В	606	_	-	11/37/37/37	-
3	NAG	В	604	1	-	0/6/23/26	0/1/1/1
3	NAG	A	604	1	-	0/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\operatorname{\AA})$
4	A	606	FAW	OG1-CG1	-2.46	1.39	1.45
4	В	606	FAW	OG1-CA1	2.38	1.40	1.33
4	A	606	FAW	OG2-CB1	2.32	1.40	1.34
4	В	606	FAW	OG2-CG2	-2.22	1.41	1.46
4	В	606	FAW	OG2-CB1	2.21	1.40	1.34
4	A	606	FAW	OG2-CG2	-2.19	1.41	1.46
4	A	606	FAW	OG1-CA1	2.19	1.39	1.33
4	В	606	FAW	OG1-CG1	-2.10	1.40	1.45

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
4	В	606	FAW	OG2-CB1-CB2	4.85	121.95	111.50
4	A	606	FAW	OG2-CB1-CB2	4.67	121.57	111.50
3	В	604	NAG	C1-O5-C5	2.87	116.08	112.19
4	A	606	FAW	OG1-CA1-CA2	2.51	119.78	111.91
4	В	606	FAW	CG2-OG2-CB1	-2.44	111.77	117.79
3	A	605	NAG	C1-O5-C5	2.44	115.50	112.19
4	A	606	FAW	CG1-OG1-CA1	-2.16	109.11	117.12

There are no chirality outliers.

All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	606	FAW	OG2-CG2-CG3-OXT
4	A	606	FAW	CB2-CB1-OG2-CG2
4	В	606	FAW	CB2-CB1-OG2-CG2
4	A	606	FAW	OB1-CB1-OG2-CG2
4	В	606	FAW	OB1-CB1-OG2-CG2
4	A	606	FAW	CA2-CA1-OG1-CG1
4	A	606	FAW	OA1-CA1-OG1-CG1
3	A	605	NAG	C4-C5-C6-O6
4	A	606	FAW	CBB-CAB-CB9-CB8
4	A	606	FAW	CA2-CA3-CA4-CA5



Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	В	606	FAW	CB4-CB5-CB6-CB7
4	A	606	FAW	CB3-CB4-CB5-CB6
4	В	606	FAW	CAB-CBB-CCB-C35
4	A	606	FAW	CAB-CBB-CCB-C35
4	В	606	FAW	OG1-CG1-CG2-CG3
4	A	606	FAW	CB7-CB8-CB9-CAB
4	В	606	FAW	CBB-CAB-CB9-CB8
4	A	606	FAW	CAA-CBA-CCA-C33
4	В	606	FAW	CB3-CB4-CB5-CB6
3	В	605	NAG	C4-C5-C6-O6
3	В	605	NAG	O5-C5-C6-O6
3	A	605	NAG	O5-C5-C6-O6
4	A	606	FAW	CG1-CG2-CG3-OXT
4	A	606	FAW	CA8-CA9-CAA-CBA
4	A	606	FAW	CB9-CAB-CBB-CCB
4	A	606	FAW	CB5-CB6-CB7-CB8
4	В	606	FAW	OG1-CG1-CG2-OG2
4	A	606	FAW	CG3-CG2-OG2-CB1
4	В	606	FAW	CA6-CA7-CA8-CA9
4	A	606	FAW	CA9-CAA-CBA-CCA
4	В	606	FAW	C36-C35-CCB-CBB
4	В	606	FAW	C34-C33-CCA-CBA
4	A	606	FAW	OG1-CA1-CA2-CA3


There are no ring outliers.


1 monomer is involved in 2 short contacts:

Mol	Chain	${f Res}$	Type	Clashes	Symm-Clashes
4	В	606	FAW	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	$521/562 \; (92\%)$	-0.21	22 (4%) 36	33	7, 20, 52, 81	0
1	В	$526/562 \ (93\%)$	-0.28	17 (3%) 47	44	6, 21, 52, 90	0
All	All	1047/1124 (93%)	-0.24	39 (3%) 41	38	6, 21, 53, 90	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	GLN	4.5
1	A	35	SER	4.4
1	В	162	PRO	4.4
1	A	140	ARG	4.4
1	A	162	PRO	4.0
1	A	161	LEU	4.0
1	В	161	LEU	3.9
1	A	142	VAL	3.7
1	A	158	ILE	3.7
1	В	504	PHE	3.5
1	В	312	GLY	3.4
1	A	160	SER	3.3
1	A	143	LEU	3.3
1	A	115	ARG	3.3
1	В	237	PRO	3.1
1	В	238	LYS	3.1
1	A	312	GLY	3.1
1	В	233	TRP	2.9
1	В	160	SER	2.9
1	В	145	ARG	2.9
1	A	137	GLU	2.9
1	В	158	ILE	2.9
1	A	165	THR	2.8
1	A	36	HIS	2.7



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	163	PHE	2.6
1	В	235	ILE	2.6
1	A	504	PHE	2.5
1	В	36	HIS	2.5
1	A	164	LEU	2.5
1	A	241	ILE	2.4
1	В	159	CYS	2.3
1	A	138	LYS	2.2
1	A	513	LYS	2.2
1	В	224	HIS	2.2
1	A	159	CYS	2.2
1	A	136	LEU	2.2
1	В	239	ASP	2.2
1	В	219	ASP	2.1
1	A	223	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
3	NAG	A	605	14/15	0.73	0.23	48,64,77,77	0
3	NAG	В	605	14/15	0.74	0.22	55,70,83,86	0
4	FAW	A	606	36/36	0.77	0.22	32,56,111,117	0
4	FAW	В	606	36/36	0.81	0.23	34,65,101,108	0
3	NAG	В	604	14/15	0.87	0.22	39,58,97,97	0
3	NAG	A	604	14/15	0.93	0.08	18,28,36,44	0
2	CA	A	603	1/1	0.98	0.06	19,19,19,19	0
2	CA	A	602	1/1	0.98	0.08	20,20,20,20	0




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q<0.9
2	CA	В	603	1/1	0.98	0.07	23,23,23,23	0
2	CA	В	602	1/1	0.98	0.07	24,24,24,24	0
2	CA	A	601	1/1	0.98	0.08	17,17,17,17	0
2	CA	В	601	1/1	0.99	0.10	19,19,19,19	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















Electron density around CA A 603: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around CA B 603: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around CA B 602:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around CA A 601: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray


and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

