



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 7, 2023 – 04:54 AM EST

PDB ID : 5W7H
Title : Supercharged arPTE variant R5
Authors : Campbell, E.; Grant, J.; Wang, Y.; Sandhu, M.; Williams, R.J.; Nisbet, D.R.;
Perriman, A.; Lupton, D.; Jackson, C.J.
Deposited on : 2017-06-19
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	 3% 84% 11% . .
1	B	341	 2% 84% 11% . .

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5128 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphotriesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2528	1599	463	458	8	0	0	0
1	B	329	2526	1597	463	458	8	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	initiating methionine	UNP Q93LD7
A	80	VAL	ALA	conflict	UNP Q93LD7
A	92	ALA	SER	conflict	UNP Q93LD7
A	148	ARG	GLN	conflict	UNP Q93LD7
A	155	ARG	GLN	conflict	UNP Q93LD7
A	185	ARG	LYS	conflict	UNP Q93LD7
A	204	GLY	ALA	conflict	UNP Q93LD7
A	238	ARG	SER	conflict	UNP Q93LD7
A	263	ARG	GLU	conflict	UNP Q93LD7
A	315	ARG	ASP	conflict	UNP Q93LD7
A	365	SER	-	expression tag	UNP Q93LD7
B	25	MET	-	initiating methionine	UNP Q93LD7
B	80	VAL	ALA	conflict	UNP Q93LD7
B	92	ALA	SER	conflict	UNP Q93LD7
B	148	ARG	GLN	conflict	UNP Q93LD7
B	155	ARG	GLN	conflict	UNP Q93LD7
B	185	ARG	LYS	conflict	UNP Q93LD7
B	204	GLY	ALA	conflict	UNP Q93LD7
B	238	ARG	SER	conflict	UNP Q93LD7
B	263	ARG	GLU	conflict	UNP Q93LD7
B	315	ARG	ASP	conflict	UNP Q93LD7
B	365	SER	-	expression tag	UNP Q93LD7

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	B	2	Total 2	Zn 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	41	Total 41	O 41	0	0
3	B	29	Total 29	O 29	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphotriesterase



- Molecule 1: Phosphotriesterase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.39Å 110.08Å 112.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.88 – 2.75 34.88 – 2.75	Depositor EDS
% Data completeness (in resolution range)	90.7 (34.88-2.75) 90.7 (34.88-2.75)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.55 (at 2.76Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.221 , 0.276 0.221 , 0.276	Depositor DCC
R_{free} test set	703 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtrriage
Anisotropy	1.054	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 16.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.026 for -h,l,k	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5128	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2567	0.44	0/3482
1	B	0.24	0/2565	0.42	0/3479
All	All	0.24	0/5132	0.43	0/6961

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	169	KCX	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2528	0	2555	17	0
1	B	2526	0	2548	17	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	41	0	0	1	0
3	B	29	0	0	1	0
All	All	5128	0	5103	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:275:ARG:HH11	1:A:275:ARG:H	1.48	0.61
1:A:263:ARG:O	1:A:319:ARG:NH2	2.34	0.60
1:A:57:HIS:HB2	1:A:303:LEU:HB3	1.83	0.59
1:A:57:HIS:CE1	1:A:169:KCX:OQ2	2.60	0.55
1:A:108:ARG:NH2	1:A:153:GLU:OE2	2.38	0.54
1:B:96:GLN:NE2	3:B:501:HOH:O	2.37	0.54
1:A:319:ARG:NH1	3:A:502:HOH:O	2.41	0.51
1:B:134:PRO:HG2	1:B:139:ARG:HG3	1.92	0.50
1:B:102:SER:HA	1:B:106:ILE:HD12	1.94	0.50
1:B:188:ALA:HA	1:B:221:LEU:HD22	1.94	0.49
1:A:54:THR:HG21	1:A:297:LEU:HD22	1.95	0.48
1:A:203:SER:O	1:A:206:GLN:N	2.47	0.48
1:B:41:ARG:HD3	1:B:118:ARG:HG2	1.95	0.47
1:A:204:GLY:HA3	1:A:233:ASP:HB2	1.97	0.47
1:B:203:SER:O	1:B:205:SER:N	2.48	0.47
1:A:353:ASN:OD1	1:A:356:ARG:NH1	2.48	0.46
1:B:236:ASP:HB3	1:B:239:TYR:HB3	1.96	0.45
1:A:61:SER:OG	1:A:62:SER:N	2.50	0.45
1:B:202:THR:OG1	1:B:229:GLY:O	2.32	0.44
1:A:181:GLU:HG3	1:A:215:ILE:HD13	2.00	0.44
1:B:88:ARG:NH2	1:B:119:ALA:O	2.49	0.44
1:B:52:THR:HA	1:B:97:THR:O	2.19	0.43
1:A:315:ARG:O	1:A:319:ARG:HG3	2.19	0.43
1:B:57:HIS:HB2	1:B:303:LEU:HB3	2.00	0.43
1:A:56:GLU:OE2	1:A:299:SER:OG	2.35	0.42
1:A:207:ARG:HD3	1:A:207:ARG:HA	1.92	0.42
1:B:105:ASP:OD2	1:B:131:TRP:HB3	2.20	0.42
1:A:52:THR:HA	1:A:97:THR:O	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:HIS:O	1:B:303:LEU:HA	2.20	0.41
1:B:39:THR:OG1	1:B:42:GLY:O	2.27	0.41
1:B:111:ARG:CZ	1:B:161:THR:HG22	2.51	0.41
1:B:264:GLY:HA3	1:B:265:ASN:HA	1.88	0.40
1:B:184:LEU:HD21	1:B:212:GLN:HG2	2.04	0.40
1:A:57:HIS:O	1:A:303:LEU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/341 (96%)	306 (94%)	18 (6%)	2 (1%)	25	42
1	B	326/341 (96%)	305 (94%)	19 (6%)	2 (1%)	25	42
All	All	652/682 (96%)	611 (94%)	37 (6%)	4 (1%)	25	42

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	131	TRP
1	B	204	GLY
1	A	274	THR
1	B	236	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/272 (97%)	248 (94%)	15 (6%)	20	36
1	B	262/272 (96%)	252 (96%)	10 (4%)	33	53
All	All	525/544 (96%)	500 (95%)	25 (5%)	25	44

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	LEU
1	A	51	PHE
1	A	151	LEU
1	A	164	ARG
1	A	185	ARG
1	A	196	VAL
1	A	237	LEU
1	A	271	LEU
1	A	272	PHE
1	A	275	ARG
1	A	303	LEU
1	A	306	PHE
1	A	337	ARG
1	A	344	GLU
1	A	362	VAL
1	B	164	ARG
1	B	175	LYS
1	B	196	VAL
1	B	206	GLN
1	B	262	LEU
1	B	263	ARG
1	B	295	ARG
1	B	303	LEU
1	B	306	PHE
1	B	309	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	A	169	2,1	9,11,12	2.27	2 (22%)	5,12,14	1.55	1 (20%)
1	KCX	B	169	2,1	9,11,12	2.28	2 (22%)	5,12,14	1.49	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	169	2,1	-	2/9/10/12	-
1	KCX	B	169	2,1	-	1/9/10/12	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	169	KCX	CX-NZ	6.02	1.45	1.35
1	B	169	KCX	CX-NZ	5.98	1.45	1.35
1	A	169	KCX	OQ1-CX	2.61	1.26	1.21
1	B	169	KCX	OQ1-CX	2.61	1.26	1.21

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	169	KCX	OQ1-CX-NZ	-2.95	120.39	124.96
1	A	169	KCX	OQ1-CX-NZ	-2.86	120.53	124.96

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	169	KCX	CG-CD-CE-NZ
1	B	169	KCX	CA-CB-CG-CD
1	A	169	KCX	CA-CB-CG-CD

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	169	KCX	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	328/341 (96%)	0.27	10 (3%) 50 59	19, 31, 50, 74	0
1	B	328/341 (96%)	0.28	7 (2%) 63 72	21, 37, 56, 93	0
All	All	656/682 (96%)	0.27	17 (2%) 56 65	19, 34, 53, 93	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	263	ARG	4.5
1	B	33	THR	4.5
1	A	205	SER	4.3
1	B	361	THR	4.2
1	A	206	GLN	3.2
1	A	362	VAL	2.8
1	B	34	GLY	2.8
1	A	202	THR	2.8
1	A	34	GLY	2.5
1	B	205	SER	2.4
1	A	107	GLY	2.4
1	B	342	PRO	2.3
1	A	272	PHE	2.3
1	A	36	LEU	2.2
1	A	155	ARG	2.2
1	A	259	ALA	2.1
1	B	260	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	KCX	A	169	12/13	0.90	0.28	22,26,49,51	0
1	KCX	B	169	12/13	0.94	0.17	21,28,34,37	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZN	B	401	1/1	0.93	0.06	25,25,25,25	0
2	ZN	A	401	1/1	0.94	0.07	28,28,28,28	0
2	ZN	B	402	1/1	0.97	0.03	31,31,31,31	0
2	ZN	A	402	1/1	0.98	0.11	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.