



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 09:27 AM EDT

PDB ID : 6W83
Title : K2P2.1 (TREK-1), 100 mM K+
Authors : Lolicato, M.; Minor, D.L.
Deposited on : 2020-03-20
Resolution : 3.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1002 (4.14-3.66)
Clashscore	141614	1004 (4.12-3.68)
Ramachandran outliers	138981	1021 (4.14-3.66)
Sidechain outliers	138945	1014 (4.14-3.66)
RSRZ outliers	127900	1275 (4.20-3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	
1	B	312	

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 4366 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium channel subfamily K member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	268	2103	1405	332	362	4	0	0	0
1	B	281	2180	1452	345	378	5	0	0	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	expression tag	UNP P97438
A	84	ARG	LYS	engineered mutation	UNP P97438
A	85	GLU	GLN	engineered mutation	UNP P97438
A	86	LYS	THR	engineered mutation	UNP P97438
A	88	LEU	ILE	engineered mutation	UNP P97438
A	89	ARG	ALA	engineered mutation	UNP P97438
A	90	ALA	GLN	engineered mutation	UNP P97438
A	92	PRO	ALA	engineered mutation	UNP P97438
A	95	SER	ASN	engineered mutation	UNP P97438
A	96	ASP	SER	engineered mutation	UNP P97438
A	97	GLN	THR	engineered mutation	UNP P97438
A	119	ALA	ASN	engineered mutation	UNP P97438
A	300	ALA	SER	engineered mutation	UNP P97438
A	306	ALA	GLU	engineered mutation	UNP P97438
A	323	SER	-	expression tag	UNP P97438
A	324	ASN	-	expression tag	UNP P97438
A	325	SER	-	expression tag	UNP P97438
A	326	LEU	-	expression tag	UNP P97438
A	327	GLU	-	expression tag	UNP P97438
A	328	VAL	-	expression tag	UNP P97438
A	329	LEU	-	expression tag	UNP P97438
A	330	PHE	-	expression tag	UNP P97438
A	331	GLN	-	expression tag	UNP P97438
B	20	MET	-	expression tag	UNP P97438
B	84	ARG	LYS	engineered mutation	UNP P97438

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	GLU	GLN	engineered mutation	UNP P97438
B	86	LYS	THR	engineered mutation	UNP P97438
B	88	LEU	ILE	engineered mutation	UNP P97438
B	89	ARG	ALA	engineered mutation	UNP P97438
B	90	ALA	GLN	engineered mutation	UNP P97438
B	92	PRO	ALA	engineered mutation	UNP P97438
B	95	SER	ASN	engineered mutation	UNP P97438
B	96	ASP	SER	engineered mutation	UNP P97438
B	97	GLN	THR	engineered mutation	UNP P97438
B	119	ALA	ASN	engineered mutation	UNP P97438
B	300	ALA	SER	engineered mutation	UNP P97438
B	306	ALA	GLU	engineered mutation	UNP P97438
B	323	SER	-	expression tag	UNP P97438
B	324	ASN	-	expression tag	UNP P97438
B	325	SER	-	expression tag	UNP P97438
B	326	LEU	-	expression tag	UNP P97438
B	327	GLU	-	expression tag	UNP P97438
B	328	VAL	-	expression tag	UNP P97438
B	329	LEU	-	expression tag	UNP P97438
B	330	PHE	-	expression tag	UNP P97438
B	331	GLN	-	expression tag	UNP P97438

- Molecule 2 is CADMIUM ION (three-letter code: CD) (formula: Cd).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Cd 2 2	0	0
2	B	1	Total Cd 1 1	0	0

- Molecule 3 is DODECANE (three-letter code: D12) (formula: C₁₂H₂₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 12 12	0	0
3	A	1	Total C 12 12	0	0

- Molecule 4 is N-OCTANE (three-letter code: OCT) (formula: C_8H_{18}).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C 8 8	0	0
4	A	1	Total C 8 8	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C 8 8	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	5	Total K 5 5	0	0
5	B	1	Total K 1 1	0	0

- Molecule 6 is HEXADECANE (three-letter code: R16) (formula: C₁₆H₃₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C 16 16	0	0

- Molecule 7 is DECANE (three-letter code: D10) (formula: C₁₀H₂₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C 10 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Potassium channel subfamily K member 2



- Molecule 1: Potassium channel subfamily K member 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.22Å 121.16Å 129.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.95 – 3.90 14.95 – 3.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (14.95-3.90) 100.0 (14.95-3.90)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 3.88Å)	Xtrriage
Refinement program	BUSTER 2.10.3 (6-FEB-2020)	Depositor
R, R_{free}	0.242 , 0.312 0.275 , 0.336	Depositor DCC
R_{free} test set	465 reflections (4.51%)	wwPDB-VP
Wilson B-factor (Å ²)	166.9	Xtrriage
Anisotropy	0.246	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 113.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4366	wwPDB-VP
Average B, all atoms (Å ²)	217.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CD, R16, D10, D12, OCT, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/2156	0.44	0/2932
1	B	0.29	0/2232	0.66	5/3035 (0.2%)
All	All	0.28	0/4388	0.56	5/5967 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	311	ARG	NE-CZ-NH2	-21.60	109.50	120.30
1	B	311	ARG	CD-NE-CZ	10.26	137.96	123.60
1	B	311	ARG	NE-CZ-NH1	8.23	124.42	120.30
1	B	311	ARG	CG-CD-NE	6.91	126.31	111.80
1	B	311	ARG	CA-CB-CG	5.58	125.69	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2103	0	2154	25	0
1	B	2180	0	2237	32	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	24	0	52	0	0
4	A	16	0	36	0	0
4	B	8	0	18	0	0
5	A	5	0	0	0	0
5	B	1	0	0	0	0
6	A	16	0	34	0	0
7	B	10	0	22	0	0
All	All	4366	0	4553	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:ARG:HH21	1:B:123:GLN:HB2	1.43	0.84
1:A:258:VAL:HG12	1:A:260:GLY:H	1.62	0.62
1:B:311:ARG:O	1:B:311:ARG:HD3	2.03	0.59
1:A:156:LYS:NZ	1:B:240:ASP:OD1	2.34	0.59
1:A:252:ILE:HD11	1:B:167:ILE:HD11	1.84	0.59
1:A:223:VAL:HA	1:A:245:VAL:HG11	1.88	0.55
1:A:267:LEU:H	1:A:267:LEU:HD23	1.71	0.55
1:B:223:VAL:HA	1:B:245:VAL:HG11	1.88	0.55
1:A:167:ILE:HD11	1:B:252:ILE:HD11	1.88	0.55
1:A:245:VAL:HG13	1:A:277:TRP:HZ2	1.72	0.54
1:B:109:ALA:HB1	1:B:114:ILE:HG21	1.90	0.53
1:B:197:ILE:HA	1:B:201:VAL:HG11	1.91	0.53
1:A:45:LYS:HG2	1:A:47:VAL:H	1.73	0.52
1:B:152:THR:HG22	1:B:154:GLY:H	1.74	0.52
1:B:245:VAL:HG13	1:B:277:TRP:HZ2	1.75	0.52
1:B:147:ASN:ND2	1:B:258:VAL:HG11	2.27	0.50
1:A:202:SER:O	1:A:206:ILE:HG12	2.13	0.49
1:B:258:VAL:HG12	1:B:260:GLY:H	1.76	0.49
1:A:252:ILE:HG12	1:B:142:THR:HA	1.94	0.49
1:B:141:THR:HG22	1:B:170:PHE:CZ	2.48	0.49
1:A:109:ALA:O	1:A:113:GLY:N	2.42	0.48
1:A:152:THR:HG22	1:A:154:GLY:H	1.79	0.48
1:A:62:ALA:HB2	1:B:133:PHE:HA	1.96	0.47
1:A:68:LEU:HD13	1:B:155:GLY:HA2	1.97	0.47
1:A:200:ASN:OD1	1:A:201:VAL:HG23	2.15	0.47
1:A:142:THR:HA	1:B:252:ILE:HG12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91:HIS:O	1:B:94:VAL:HG12	2.15	0.47
1:B:285:PHE:HA	1:B:288:VAL:HB	1.97	0.47
1:A:141:THR:HG22	1:A:170:PHE:CZ	2.50	0.46
1:B:201:VAL:HB	1:B:206:ILE:HD11	1.96	0.46
1:A:99:LEU:O	1:A:103:ILE:HG12	2.16	0.45
1:A:147:ASN:ND2	1:A:258:VAL:HG11	2.31	0.45
1:B:107:VAL:HA	1:B:110:ILE:HD12	1.98	0.45
1:A:193:GLU:HA	1:A:197:ILE:HG13	1.98	0.45
1:A:127:TRP:CZ3	1:B:65:PHE:HB3	2.53	0.44
1:B:114:ILE:HB	1:B:116:PRO:HD3	1.99	0.44
1:B:270:TYR:O	1:B:274:VAL:HG23	2.18	0.44
1:A:133:PHE:HA	1:B:62:ALA:HB2	2.00	0.43
1:A:285:PHE:HA	1:A:288:VAL:HB	2.01	0.43
1:B:37:SER:HA	1:B:40:ASN:OD1	2.19	0.43
1:B:106:ILE:O	1:B:110:ILE:HG13	2.20	0.42
1:B:111:ASN:HB3	1:B:151:ARG:NH2	2.34	0.42
1:B:147:ASN:CG	1:B:148:ILE:HD12	2.41	0.41
1:B:218:GLY:HA3	1:B:284:TYR:CZ	2.55	0.41
1:B:125:SER:HB3	1:B:128:ASP:HB3	2.03	0.41
1:B:298:VAL:O	1:B:302:LYS:HG2	2.21	0.40
1:A:128:ASP:OD1	1:A:131:SER:N	2.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/312 (85%)	256 (97%)	8 (3%)	0	100	100
1	B	277/312 (89%)	260 (94%)	16 (6%)	1 (0%)	34	71
All	All	541/624 (87%)	516 (95%)	24 (4%)	1 (0%)	47	79

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	264	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/260 (85%)	220 (100%)	0	100	100
1	B	229/260 (88%)	228 (100%)	1 (0%)	91	94
All	All	449/520 (86%)	448 (100%)	1 (0%)	93	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	311	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 9 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	D10	B	404	-	9,9,9	0.25	0	8,8,8	0.85	0
4	OCT	B	402	-	7,7,7	0.38	0	6,6,6	0.57	0
6	R16	A	412	-	15,15,15	0.33	0	14,14,14	0.86	0
4	OCT	A	406	-	7,7,7	0.29	0	6,6,6	0.68	0
4	OCT	A	405	-	7,7,7	0.43	0	6,6,6	0.74	0
3	D12	A	404	-	11,11,11	0.33	0	10,10,10	0.84	0
3	D12	A	403	-	11,11,11	0.33	0	10,10,10	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	D10	B	404	-	-	0/7/7/7	-
4	OCT	B	402	-	-	1/5/5/5	-
6	R16	A	412	-	-	3/13/13/13	-
4	OCT	A	406	-	-	0/5/5/5	-
4	OCT	A	405	-	-	0/5/5/5	-
3	D12	A	404	-	-	2/9/9/9	-
3	D12	A	403	-	-	4/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	412	R16	C30-C31-C32-C33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	412	R16	C34-C35-C36-C37
4	B	402	OCT	C3-C4-C5-C6
3	A	403	D12	C11-C10-C9-C8
3	A	403	D12	C5-C6-C7-C8
3	A	403	D12	C6-C7-C8-C9
3	A	404	D12	C3-C4-C5-C6
3	A	403	D12	C7-C8-C9-C10
6	A	412	R16	C27-C28-C29-C30
3	A	404	D12	C4-C5-C6-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/312 (85%)	-0.35	11 (4%) 37 29	132, 214, 294, 300	0
1	B	281/312 (90%)	-0.35	13 (4%) 32 26	117, 216, 296, 300	0
All	All	549/624 (87%)	-0.35	24 (4%) 34 28	117, 214, 295, 300	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	92	PRO	5.6
1	A	265	GLU	4.4
1	B	97	GLN	4.3
1	B	195	THR	4.1
1	A	200	ASN	4.0
1	B	93	CYS	3.7
1	B	196	PHE	3.6
1	B	199	TRP	3.6
1	B	200	ASN	3.5
1	B	203	GLN	3.4
1	B	96	ASP	3.1
1	B	36	ASP	2.8
1	A	199	TRP	2.5
1	A	320	ASN	2.5
1	B	261	GLY	2.5
1	A	195	THR	2.4
1	A	269	PHE	2.4
1	A	203	GLN	2.4
1	A	153	GLU	2.3
1	A	90	ALA	2.3
1	A	263	ASP	2.3
1	B	309	GLU	2.2
1	A	235	GLY	2.1
1	B	262	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	K	A	408	1/1	0.79	0.25	172,172,172,172	0
2	CD	A	402	1/1	0.80	0.16	188,188,188,188	1
3	D12	A	404	12/12	0.81	0.28	108,158,192,193	0
5	K	A	411	1/1	0.83	0.70	176,176,176,176	0
7	D10	B	404	10/10	0.84	0.30	128,149,155,156	0
5	K	A	410	1/1	0.85	0.18	175,175,175,175	0
4	OCT	A	406	8/8	0.86	0.28	147,158,171,173	0
6	R16	A	412	16/16	0.86	0.26	125,156,204,211	0
5	K	A	407	1/1	0.86	0.18	156,156,156,156	0
4	OCT	B	402	8/8	0.88	0.13	111,139,171,177	0
5	K	B	403	1/1	0.90	0.27	157,157,157,157	0
3	D12	A	403	12/12	0.95	0.15	88,94,129,131	0
4	OCT	A	405	8/8	0.96	0.10	94,104,107,112	0
2	CD	B	401	1/1	0.96	0.03	221,221,221,221	0
5	K	A	409	1/1	0.98	0.10	139,139,139,139	0
2	CD	A	401	1/1	0.98	0.03	170,170,170,170	0

6.5 Other polymers [i](#)

There are no such residues in this entry.