



wwPDB X-ray Structure Validation Summary Report

Nov 7, 2023 – 12:17 am GMT

PDB ID : 1WA3
Title : Mechanism of the Class I KDPG aldolase
Authors : Fullerton, S.W.B.; Griffiths, J.S.; Merkel, A.B.; Wymer, N.J.; Hutchins, M.J.;
Fierke, C.A.; Toone, E.J.; Naismith, J.H.
Deposited on : 2004-10-22
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	
1	B	205	
1	C	205	
1	D	205	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9569 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-KETO-3-DEOXY-6-PHOSPHOGLUCONATE ALDOLASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	Total 1536	C 995	N 253	O 279	S 9	0	0	0
1	B	202	Total 1536	C 995	N 253	O 279	S 9	0	0	0
1	C	201	Total 1527	C 989	N 251	O 278	S 9	0	0	0
1	D	203	Total 1541	C 998	N 254	O 280	S 9	0	0	0
1	E	202	Total 1536	C 995	N 253	O 279	S 9	0	0	0
1	F	201	Total 1527	C 989	N 251	O 278	S 9	0	0	0

- Molecule 2 is PYRUVIC ACID (three-letter code: PYR) (formula: C₃H₄O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 5 3 2	0	0
2	B	1	Total C O 5 3 2	0	0
2	C	1	Total C O 5 3 2	0	0
2	D	1	Total C O 5 3 2	0	0
2	E	1	Total C O 5 3 2	0	0
2	F	1	Total C O 5 3 2	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	F	1	Total O S 5 4 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	74	Total O 74 74	0	0
4	B	73	Total O 73 73	0	0
4	C	41	Total O 41 41	0	0
4	D	49	Total O 49 49	0	0
4	E	46	Total O 46 46	0	0
4	F	23	Total O 23 23	0	0

SEQUENCE-PLOTS INFOmissingINFO

3 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.60Å 101.10Å 124.60Å 90.00° 97.20° 90.00°	Depositor
Resolution (Å)	32.44 – 1.90 32.42 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.0 (32.44-1.90) 94.0 (32.42-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.90 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.2.0007	Depositor
R, R_{free}	0.193 , 0.239 0.193 , 0.239	Depositor DCC
R_{free} test set	3900 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtrriage
Anisotropy	0.031	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9569	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PYR, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.86	1/1565 (0.1%)	0.84	2/2108 (0.1%)
1	B	0.83	0/1565	0.82	4/2108 (0.2%)
1	C	0.68	0/1556	0.75	2/2097 (0.1%)
1	D	0.64	0/1570	0.67	0/2115
1	E	0.71	0/1565	0.71	1/2108 (0.0%)
1	F	0.62	0/1556	0.71	2/2097 (0.1%)
All	All	0.73	1/9377 (0.0%)	0.75	11/12633 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	VAL	CB-CG2	-5.12	1.42	1.52

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	172	VAL	CG1-CB-CG2	8.58	124.63	110.90
1	C	111	VAL	CG1-CB-CG2	8.24	124.08	110.90
1	A	172	VAL	CG1-CB-CG2	7.79	123.36	110.90
1	B	31	VAL	CG1-CB-CG2	6.67	121.56	110.90
1	B	173	LEU	CA-CB-CG	6.30	129.79	115.30

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1536	0	1593	15	2
1	B	1536	0	1592	18	1
1	C	1527	0	1579	23	1
1	D	1541	0	1594	21	0
1	E	1536	0	1592	24	0
1	F	1527	0	1579	28	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
4	A	74	0	0	2	0
4	B	73	0	0	1	0
4	C	41	0	0	0	0
4	D	49	0	0	0	0
4	E	46	0	0	0	0
4	F	23	0	0	1	0
All	All	9569	0	9529	120	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 120 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:117:LEU:HD13	1:C:144:MET:HE3	1.21	1.17
1:C:117:LEU:HD13	1:C:144:MET:CE	1.80	1.09
1:E:136:VAL:HG22	1:E:140:PHE:HB2	1.33	1.07
1:C:117:LEU:CD1	1:C:144:MET:HE3	1.87	1.04
1:B:187:ASP:OD2	4:B:2066:HOH:O	1.78	1.01

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:ARG:NH1	1:A:198:GLU:OE2[1_455]	1.82	0.38
1:B:77:ARG:NH1	1:B:162:ASP:OD2[1_455]	2.08	0.12
1:A:19:ASN:OD1	1:C:187:ASP:OD1[2_655]	2.16	0.04

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/205 (98%)	200 (100%)	0	0	100	100
1	B	200/205 (98%)	197 (98%)	3 (2%)	0	100	100
1	C	199/205 (97%)	194 (98%)	5 (2%)	0	100	100
1	D	201/205 (98%)	199 (99%)	2 (1%)	0	100	100
1	E	200/205 (98%)	198 (99%)	2 (1%)	0	100	100
1	F	199/205 (97%)	196 (98%)	3 (2%)	0	100	100
All	All	1199/1230 (98%)	1184 (99%)	15 (1%)	0	100	100

There are no Ramachandran outliers to report.

4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/169 (98%)	159 (96%)	7 (4%)	30	20

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	166/169 (98%)	156 (94%)	10 (6%)	19 9
1	C	165/169 (98%)	154 (93%)	11 (7%)	16 7
1	D	166/169 (98%)	160 (96%)	6 (4%)	35 26
1	E	166/169 (98%)	158 (95%)	8 (5%)	25 16
1	F	165/169 (98%)	146 (88%)	19 (12%)	5 2
All	All	994/1014 (98%)	933 (94%)	61 (6%)	18 9

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	111	VAL
1	F	172	VAL
1	E	31	VAL
1	F	166	GLU
1	F	183	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	98	GLN
1	D	19	ASN
1	E	98	GLN
1	F	37	HIS
1	F	139	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PYR	F	1204	1	4,4,5	1.77	2 (50%)	4,4,6	1.21	0
3	SO4	E	1205	-	4,4,4	0.18	0	6,6,6	0.70	0
3	SO4	C	1205	-	4,4,4	0.10	0	6,6,6	0.65	0
3	SO4	F	1205	-	4,4,4	0.13	0	6,6,6	0.44	0
3	SO4	B	1205	-	4,4,4	0.48	0	6,6,6	0.88	0
2	PYR	A	1204	1	4,4,5	2.06	1 (25%)	4,4,6	1.26	0
2	PYR	D	1205	1	4,4,5	1.89	1 (25%)	4,4,6	1.44	0
2	PYR	E	1204	1	4,4,5	1.74	2 (50%)	4,4,6	1.47	1 (25%)
3	SO4	D	1206	-	4,4,4	0.22	0	6,6,6	0.61	0
2	PYR	C	1204	1	4,4,5	1.39	1 (25%)	4,4,6	1.78	2 (50%)
2	PYR	B	1204	1	4,4,5	1.73	1 (25%)	4,4,6	1.16	0
3	SO4	A	1205	-	4,4,4	0.10	0	6,6,6	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PYR	F	1204	1	-	0/2/2/4	-
2	PYR	A	1204	1	-	1/2/2/4	-
2	PYR	D	1205	1	-	0/2/2/4	-
2	PYR	E	1204	1	-	0/2/2/4	-
2	PYR	C	1204	1	-	0/2/2/4	-
2	PYR	B	1204	1	-	0/2/2/4	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1204	PYR	O-C	3.60	1.34	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1205	PYR	O-C	3.24	1.32	1.22
2	B	1204	PYR	O-C	2.93	1.31	1.22
2	F	1204	PYR	O-C	2.88	1.31	1.22
2	E	1204	PYR	O-C	2.69	1.31	1.22

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1204	PYR	OXT-C-O	-2.60	116.81	123.30
2	C	1204	PYR	OXT-C-CA	2.11	122.43	113.35
2	E	1204	PYR	OXT-C-CA	2.11	122.39	113.35

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1204	PYR	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/205 (98%)	-0.25	1 (0%) 91 92	2, 10, 20, 24	1 (0%)
1	B	202/205 (98%)	-0.25	1 (0%) 91 92	3, 9, 19, 25	0
1	C	201/205 (98%)	0.58	8 (3%) 38 41	2, 10, 22, 32	0
1	D	203/205 (99%)	0.36	3 (1%) 73 76	5, 12, 24, 30	0
1	E	202/205 (98%)	0.15	2 (0%) 82 84	3, 14, 25, 29	1 (0%)
1	F	201/205 (98%)	0.39	8 (3%) 38 41	7, 23, 38, 48	0
All	All	1211/1230 (98%)	0.16	23 (1%) 66 69	2, 12, 29, 48	2 (0%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	60	GLU	3.9
1	C	56	SER	3.4
1	C	165	CYS	2.8
1	F	82	SER	2.8
1	A	2	LYS	2.6

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PYR	E	1204	5/6	0.92	0.10	15,15,17,17	0
2	PYR	C	1204	5/6	0.95	0.09	10,13,13,15	0
2	PYR	D	1205	5/6	0.96	0.09	12,12,14,14	0
2	PYR	F	1204	5/6	0.96	0.08	17,19,20,20	0
3	SO4	D	1206	5/5	0.96	0.09	28,30,32,33	0
3	SO4	A	1205	5/5	0.97	0.07	24,25,29,29	0
3	SO4	C	1205	5/5	0.97	0.10	22,23,24,25	0
2	PYR	B	1204	5/6	0.97	0.09	9,10,12,12	0
3	SO4	E	1205	5/5	0.97	0.07	24,25,28,28	0
3	SO4	B	1205	5/5	0.98	0.07	13,13,17,20	0
2	PYR	A	1204	5/6	0.98	0.06	7,7,9,9	0
3	SO4	F	1205	5/5	0.98	0.07	29,31,32,32	0

5.5 Other polymers [i](#)

There are no such residues in this entry.