



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 5, 2023 – 04:53 am GMT

PDB ID : 1WA5  
Title : Structure of the Cse1:Imp-alpha:RanGTP complex  
Authors : Stewart, M.  
Deposited on : 2004-10-23  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	176	 2% 90% 6%
2	B	530	 8% 80% 8% 12%
3	C	960	 6% 91% 8%

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 26541 atoms, of which 13019 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	170	2815	905	1420	247	239	4	0	1	0

- Molecule 2 is a protein called SRP1 isoform 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
2	B	466	7464	2327	3766	644	711	16	0	8	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	117	GLN	ARG	conflict	UNP A0A6A5Q590

- Molecule 3 is a protein called Importin alpha re-exporter.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
3	C	950	15512	4985	7824	1247	1436	20	0	10	0

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>14</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
4	A	1	41	10	9	5	14	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	101	Total	O	0	0
			101	101		
6	B	166	Total	O	0	0
			166	166		
6	C	441	Total	O	0	0
			441	441		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: GTP-binding nuclear protein Ran




- Molecule 2: SRP1 isoform 1



- Molecule 3: Importin alpha re-exporter





## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	272.67Å 72.26Å 97.64Å 90.00° 92.57° 90.00°	Depositor
Resolution (Å)	34.92 – 2.00 34.92 – 2.00	Depositor EDS
% Data completeness (in resolution range)	93.9 (34.92-2.00) 93.9 (34.92-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, $R_{free}$	0.188 , 0.216 0.188 , 0.217	Depositor DCC
$R_{free}$ test set	7266 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.2	Xtrriage
Anisotropy	0.140	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.014 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	26541	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1433	0.48	0/1932
2	B	0.24	0/3782	0.40	0/5127
3	C	0.25	0/7888	0.41	0/10718
All	All	0.25	0/13103	0.42	0/17777

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1395	1420	1420	7	0
2	B	3698	3766	3735	32	0
3	C	7688	7824	7782	54	0
4	A	32	9	12	0	0
5	A	1	0	0	0	0
6	A	101	0	0	1	0
6	B	166	0	0	9	0
6	C	441	0	0	13	0
All	All	13522	13019	12949	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 3.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:141:LYS:NZ	6:C:1004:HOH:O	2.15	0.80
3:C:766:LEU:O	3:C:778:TYR:OH	2.04	0.75
3:C:87:LYS:O	6:C:1001:HOH:O	2.06	0.74
2:B:192:GLN:OE1	6:B:601:HOH:O	2.07	0.73
3:C:744:GLU:OE2	6:C:1002:HOH:O	2.08	0.72
3:C:33:GLN:NE2	6:C:1007:HOH:O	2.24	0.71
3:C:183:LYS:O	6:C:1003:HOH:O	2.12	0.68
3:C:800[B]:HIS:ND1	6:C:1009:HOH:O	2.27	0.67
2:B:50:GLU:OE1	6:B:602:HOH:O	2.13	0.66
2:B:89:LEU:HD11	2:B:129:ALA:HB2	1.80	0.64
1:A:110:ARG:NH1	6:A:301:HOH:O	2.29	0.64
2:B:422[B]:ARG:NH1	6:B:605:HOH:O	2.32	0.63
3:C:747[B]:MET:HE3	3:C:789:ILE:HD11	1.81	0.62
3:C:369:LEU:HD11	3:C:518:ILE:HD13	1.81	0.62
3:C:193:GLU:O	6:C:1005:HOH:O	2.16	0.60
3:C:137:MET:CE	3:C:185:VAL:HG22	2.33	0.58
3:C:558:LEU:HD23	3:C:567:LEU:HD11	1.86	0.58
3:C:903:GLU:N	3:C:903:GLU:OE1	2.33	0.58
3:C:558:LEU:HD23	3:C:567:LEU:CD1	2.35	0.56
2:B:17:ARG:NH2	3:C:117:PHE:O	2.36	0.56
3:C:176:ALA:HB3	3:C:177:PRO:HD3	1.88	0.55
3:C:958:VAL:HG12	3:C:958:VAL:O	2.07	0.55
2:B:58:PHE:N	6:B:608:HOH:O	2.41	0.54
2:B:373:THR:HG22	2:B:417:ARG:CZ	2.38	0.54
3:C:137:MET:HE1	3:C:205:VAL:HG11	1.90	0.53
2:B:33:ARG:NH1	2:B:367:ASN:O	2.42	0.52
3:C:187:GLU:N	6:C:1003:HOH:O	2.21	0.52
3:C:357:PHE:CZ	3:C:472:ILE:HD13	2.44	0.51
2:B:472:GLU:OE2	6:B:603:HOH:O	2.18	0.51
1:A:56:ARG:NE	1:A:171:ASP:OD2	2.43	0.51
2:B:33:ARG:NH2	2:B:331:ASP:OD1	2.44	0.51
2:B:226:SER:CB	2:B:231:LEU:HD23	2.41	0.49
2:B:393:GLU:HG3	2:B:394:VAL:HG13	1.95	0.49
2:B:89:LEU:HD11	2:B:129:ALA:CB	2.42	0.49
3:C:67:ARG:NH1	6:C:1015:HOH:O	2.32	0.49
3:C:531:THR:O	6:C:1006:HOH:O	2.20	0.49
3:C:137:MET:HE2	3:C:185:VAL:HG22	1.94	0.48
3:C:279:ASP:OD1	3:C:279:ASP:N	2.47	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:743:LEU:HG	3:C:747[B]:MET:HE1	1.95	0.48
2:B:300:ILE:N	2:B:301:PRO:CD	2.76	0.48
3:C:386:LYS:O	3:C:390:GLU:HG3	2.12	0.48
2:B:514:ASP:OD1	3:C:579:ARG:NH1	2.47	0.47
1:A:77:ASP:N	1:A:77:ASP:OD1	2.47	0.47
3:C:44:ILE:HD13	3:C:105:ILE:HD11	1.97	0.47
1:A:174:LEU:HD23	1:A:175:GLU:N	2.29	0.47
1:A:29:ARG:HG2	1:A:157:PHE:CE1	2.50	0.47
2:B:192:GLN:NE2	6:B:606:HOH:O	2.35	0.47
3:C:12:ALA:O	3:C:15:VAL:HG12	2.14	0.47
3:C:91:VAL:HB	3:C:92:PRO:HD3	1.97	0.47
2:B:87:GLN:OE1	2:B:87:GLN:N	2.48	0.46
3:C:137:MET:HE1	3:C:185:VAL:HG22	1.97	0.46
2:B:431:LYS:HB3	2:B:432:PRO:HD3	1.97	0.45
3:C:575:ARG:HG2	3:C:618:TYR:CD2	2.51	0.45
2:B:120:ARG:N	2:B:121:PRO:CD	2.80	0.45
3:C:317:THR:HG23	3:C:387:GLU:HG3	1.98	0.45
3:C:814:TRP:CH2	3:C:841:MET:HG3	2.52	0.45
3:C:614:ARG:NH1	6:C:1022:HOH:O	2.37	0.45
3:C:221:ILE:HG23	3:C:221:ILE:O	2.17	0.45
2:B:343:LEU:HB2	2:B:344:PRO:HD3	1.98	0.44
2:B:390:LYS:O	2:B:394:VAL:HG22	2.17	0.44
1:A:117:ILE:HB	1:A:144:LEU:HD22	1.98	0.44
2:B:419:ASP:OD1	2:B:422[A]:ARG:NH1	2.51	0.44
3:C:915:ASN:ND2	6:C:1018:HOH:O	2.36	0.44
3:C:938:LEU:HD13	3:C:957:LEU:HD13	1.99	0.44
2:B:347:ARG:NE	6:B:617:HOH:O	2.50	0.44
3:C:464:LEU:HD11	3:C:482:ILE:HD11	2.00	0.44
3:C:4:LEU:C	3:C:4:LEU:HD23	2.38	0.43
3:C:299:ILE:HD12	3:C:299:ILE:HA	1.92	0.43
2:B:421:ILE:HG22	2:B:458:MET:HE3	2.00	0.43
3:C:521:GLU:O	3:C:525:THR:HG23	2.18	0.43
3:C:320:THR:HG22	3:C:391:LYS:HG3	2.00	0.43
3:C:422:LYS:HA	3:C:425:TYR:CE2	2.54	0.43
3:C:692:TRP:CZ3	3:C:701:VAL:HG23	2.53	0.43
2:B:247:LYS:N	6:B:614:HOH:O	2.48	0.43
3:C:365:ILE:HD13	3:C:514:THR:HG21	2.01	0.43
2:B:105:LEU:HD11	2:B:146:MET:HG3	2.00	0.42
3:C:784:VAL:HG21	3:C:830:ASP:HB3	2.00	0.42
2:B:396:GLU:OE1	6:B:604:HOH:O	2.22	0.42
1:A:95:ARG:NH1	2:B:490:GLN:OE1	2.49	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:747[B]:MET:SD	3:C:785:PHE:HZ	2.43	0.42
3:C:186:ASP:OD1	3:C:239:TYR:OH	2.17	0.41
2:B:385:ILE:HB	2:B:386:PRO:HD3	2.01	0.41
2:B:421:ILE:HG22	2:B:458:MET:CE	2.51	0.41
3:C:361:PRO:HB2	3:C:510:TYR:CD1	2.56	0.41
2:B:37:ASP:OD1	2:B:37:ASP:N	2.54	0.41
3:C:186:ASP:N	6:C:1003:HOH:O	2.53	0.40
2:B:421:ILE:CG2	2:B:458:MET:HE3	2.51	0.40
3:C:31:GLU:OE1	3:C:67:ARG:NH2	2.54	0.40
3:C:836:ILE:HD12	3:C:921:TYR:CD1	2.57	0.40
3:C:267:GLN:HA	3:C:267:GLN:OE1	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/176 (96%)	164 (97%)	5 (3%)	0	100	100
2	B	466/530 (88%)	460 (99%)	6 (1%)	0	100	100
3	C	952/960 (99%)	933 (98%)	19 (2%)	0	100	100
All	All	1587/1666 (95%)	1557 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	151/153 (99%)	151 (100%)	0	100	100
2	B	409/457 (90%)	409 (100%)	0	100	100
3	C	882/882 (100%)	880 (100%)	2 (0%)	93	95
All	All	1442/1492 (97%)	1440 (100%)	2 (0%)	93	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	515	TYR
3	C	745	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GTP	A	201	5	26,34,34	1.11	2 (7%)	32,54,54	1.40	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	A	201	5	-	6/18/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	201	GTP	C5-C6	-3.79	1.39	1.47
4	A	201	GTP	C2-N3	2.16	1.38	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	201	GTP	C5-C6-N1	3.16	119.54	113.95
4	A	201	GTP	C8-N7-C5	3.12	108.94	102.99
4	A	201	GTP	C2-N1-C6	-2.89	119.78	125.10
4	A	201	GTP	PA-O3A-PB	-2.53	124.13	132.83
4	A	201	GTP	PB-O3B-PG	-2.46	124.39	132.83

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	201	GTP	C5'-O5'-PA-O3A
4	A	201	GTP	C5'-O5'-PA-O1A
4	A	201	GTP	C5'-O5'-PA-O2A
4	A	201	GTP	O4'-C4'-C5'-O5'
4	A	201	GTP	C3'-C4'-C5'-O5'
4	A	201	GTP	PA-O3A-PB-O2B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	170/176 (96%)	0.11	3 (1%) 68 66	18, 33, 69, 120	0
2	B	466/530 (87%)	0.47	44 (9%) 8 8	28, 49, 98, 157	0
3	C	950/960 (98%)	0.28	53 (5%) 24 23	19, 42, 82, 141	0
All	All	1586/1666 (95%)	0.32	100 (6%) 20 19	18, 44, 86, 157	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	PRO	6.9
2	B	109	VAL	5.7
3	C	255	HIS	5.4
2	B	117	GLN	5.3
3	C	400	PHE	5.0
3	C	250	PRO	4.9
2	B	514	ASP	4.6
2	B	101	MET	4.6
2	B	108	THR	4.5
2	B	58	PHE	4.5
3	C	251	ASP	4.5
3	C	417	LYS	4.4
3	C	247	LEU	4.3
2	B	123	ILE	4.2
3	C	958	VAL	4.2
3	C	253	THR	4.1
3	C	374	THR	4.0
3	C	135	ASP	3.9
2	B	102	GLN	3.9
3	C	414	ASP	3.9
3	C	413	SER	3.8
3	C	246	LEU	3.8
2	B	111	PHE	3.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	B	115	LEU	3.7
3	C	880	VAL	3.7
2	B	20	ASN	3.7
3	C	328	ILE	3.7
2	B	120	ARG	3.6
2	B	247	LYS	3.6
2	B	94	GLN	3.5
3	C	355	GLU	3.5
3	C	959	GLY	3.3
3	C	32	THR	3.2
2	B	95	GLN	3.2
3	C	197	ALA	3.2
3	C	34	ASP	3.1
3	C	352	GLU	3.1
2	B	512	GLU	3.1
2	B	107	ALA	3.1
3	C	1	MET	3.1
3	C	350	ARG	3.1
2	B	105	LEU	3.0
3	C	407	PHE	3.0
2	B	38	THR	3.0
3	C	33	GLN	3.0
3	C	252	GLU	2.9
2	B	100	ASP	2.9
3	C	411	TYR	2.9
3	C	353	ASP	2.9
3	C	356	LEU	2.8
3	C	349	LEU	2.8
2	B	99	ASP	2.8
3	C	329	PHE	2.8
2	B	88	GLU	2.7
2	B	511	GLU	2.7
2	B	19	THR	2.7
2	B	113	GLN	2.7
3	C	254	GLU	2.6
2	B	47	LYS	2.6
3	C	199	LEU	2.6
2	B	498	TYR	2.6
2	B	91	GLN	2.6
3	C	412	MET	2.6
2	B	146	MET	2.5
3	C	293[A]	TRP	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	39	GLN	2.5
2	B	280	ALA	2.5
2	B	510	GLY	2.5
1	A	140	ARG	2.4
3	C	248	GLU	2.4
2	B	103	GLU	2.4
2	B	40	GLN	2.4
3	C	472	ILE	2.4
2	B	92	MET	2.3
2	B	114	ILE	2.3
3	C	375	ASP	2.3
3	C	340	GLU	2.2
2	B	187	VAL	2.2
3	C	271	GLN	2.2
3	C	416	SER	2.2
2	B	515	ALA	2.2
2	B	121	PRO	2.1
3	C	137	MET	2.1
3	C	947	GLN	2.1
2	B	126	VAL	2.1
3	C	915	ASN	2.1
3	C	266	ILE	2.1
3	C	468	ASN	2.1
2	B	48	ARG	2.1
2	B	347	ARG	2.1
3	C	351	GLU	2.1
3	C	358	GLU	2.1
3	C	951	VAL	2.1
2	B	439	ILE	2.0
3	C	77	LEU	2.0
2	B	397	TYR	2.0
3	C	201	ILE	2.0
1	A	114	ASN	2.0
3	C	771	GLN	2.0
3	C	303	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MG	A	202	1/1	0.88	0.18	35,35,35,35	0
4	GTP	A	201	32/32	0.98	0.11	16,25,31,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.