



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 14, 2020 – 07:54 pm BST

PDB ID : 2WDV  
Title : E. coli succinate:quinone oxidoreductase (SQR) with an empty quinone- binding pocket  
Authors : Ruprecht, J.; Yankovskaya, V.; Maklashina, E.; Iwata, S.; Cecchini, G.  
Deposited on : 2009-03-26  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	
1	E	588	
1	I	588	
2	B	238	
2	F	238	
2	J	238	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	129	
3	G	129	
3	K	129	
4	D	115	
4	H	115	
4	L	115	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	TEO	E	1589	-	-	X	-
6	TEO	I	1589	-	-	X	-
8	FES	J	302	-	-	X	-
9	SF4	F	303	-	-	X	-

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 24855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	588	Total 4522	C 2812	N 821	O 861	S 28	0	0	0
1	E	588	Total 4522	C 2812	N 821	O 861	S 28	0	0	0
1	I	588	Total 4522	C 2812	N 821	O 861	S 28	0	0	0

- Molecule 2 is a protein called SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	238	Total 1869	C 1172	N 329	O 348	S 20	0	0	0
2	F	238	Total 1869	C 1172	N 329	O 348	S 20	0	0	0
2	J	238	Total 1869	C 1172	N 329	O 348	S 20	0	0	0


- Molecule 3 is a protein called SUCCINATE DEHYDROGENASE CYTOCHROME B556 SUBUNIT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	121	Total 933	C 619	N 151	O 158	S 5	0	0	0
3	G	121	Total 933	C 619	N 151	O 158	S 5	0	0	0
3	K	121	Total 933	C 619	N 151	O 158	S 5	0	0	0

- Molecule 4 is a protein called SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR PROTEIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	105	Total	C	N	O	S	0	0	0
			836	577	123	133	3			
4	H	105	Total	C	N	O	S	0	0	0
			836	577	123	133	3			
4	L	105	Total	C	N	O	S	0	0	0
			836	577	123	133	3			

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
5	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
5	I	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is MALATE LIKE INTERMEDIATE (three-letter code: TEO) (formula:  $C_4H_4O_5$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 9 4 5	0	0
6	E	1	Total C O 9 4 5	0	0
6	I	1	Total C O 9 4 5	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	I	1	Total Na 1 1	0	0
7	A	1	Total Na 1 1	0	0
7	E	1	Total Na 1 1	0	0

- Molecule 8 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			4	2	2		
8	F	1	Total	Fe	S	0	0
			4	2	2		
8	J	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 9 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	Fe	S	0	0
			8	4	4		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	F	1	Total	Fe	S	0	0
			8	4	4		
9	J	1	Total	Fe	S	0	0
			8	4	4		


- Molecule 10 is FE3-S4 CLUSTER (three-letter code: F3S) (formula: Fe<sub>3</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	B	1	Total	Fe	S	0	0
			7	3	4		
10	F	1	Total	Fe	S	0	0
			7	3	4		
10	J	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 11 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
11	C	1	43	34	1	4	4	0	0
11	G	1	43	34	1	4	4	0	0
11	K	1	43	34	1	4	4	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT




- Molecule 1: SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT





- Molecule 1: SUCCINATE DEHYDROGENASE FLAVOPROTEIN SUBUNIT









● Molecule 2: SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT




● Molecule 2: SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT



● Molecule 2: SUCCINATE DEHYDROGENASE IRON-SULFUR SUBUNIT





● Molecule 3: SUCCINATE DEHYDROGENASE CYTOCHROME B556 SUBUNIT









- Molecule 3: SUCCINATE DEHYDROGENASE CYTOCHROME B556 SUBUNIT





- Molecule 3: SUCCINATE DEHYDROGENASE CYTOCHROME B556 SUBUNIT





- Molecule 4: SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR PROTEIN



- Molecule 4: SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR PROTEIN



- Molecule 4: SUCCINATE DEHYDROGENASE HYDROPHOBIC MEMBRANE ANCHOR PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.34Å 184.85Å 204.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.85 – 3.20 49.95 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.7 (51.85-3.20) 99.7 (49.95-3.20)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.99 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.4.0067	Depositor
R, $R_{free}$	0.205 , 0.233 0.207 , 0.235	Depositor DCC
$R_{free}$ test set	3844 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.2	Xtrriage
Anisotropy	0.413	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 93.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	24855	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TEO, NA, SF4, F3S, FES, HEM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	2/4611 (0.0%)	0.64	0/6237
1	E	0.52	1/4611 (0.0%)	0.61	0/6237
1	I	0.43	1/4611 (0.0%)	0.56	0/6237
2	B	0.66	0/1908	0.67	0/2578
2	F	0.59	1/1908 (0.1%)	0.66	0/2578
2	J	0.47	1/1908 (0.1%)	0.60	0/2578
3	C	0.54	0/953	0.57	0/1293
3	G	0.52	0/953	0.56	0/1293
3	K	0.44	0/953	0.50	0/1293
4	D	0.50	0/859	0.53	0/1175
4	H	0.50	0/859	0.53	0/1175
4	L	0.47	0/859	0.53	0/1175
All	All	0.53	6/24993 (0.0%)	0.60	0/33849

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	159	CYS	CB-SG	-6.16	1.71	1.82
1	I	173	GLN	CD-NE2	-5.74	1.18	1.32
1	E	173	GLN	CD-NE2	-5.73	1.18	1.32
2	J	71	ASN	CG-ND2	-5.45	1.19	1.32
1	A	89	CYS	CB-SG	-5.36	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4522	0	4426	44	0
1	E	4522	0	4426	50	0
1	I	4522	0	4426	61	0
2	B	1869	0	1850	22	0
2	F	1869	0	1850	20	0
2	J	1869	0	1850	23	0
3	C	933	0	979	6	0
3	G	933	0	979	9	0
3	K	933	0	979	8	0
4	D	836	0	875	9	0
4	H	836	0	875	10	0
4	L	836	0	875	11	0
5	A	53	0	30	6	0
5	E	53	0	30	4	0
5	I	53	0	29	7	0
6	A	9	0	3	2	0
6	E	9	0	3	4	0
6	I	9	0	3	4	0
7	A	1	0	0	0	0
7	E	1	0	0	0	0
7	I	1	0	0	0	0
8	B	4	0	0	0	0
8	F	4	0	0	0	0
8	J	4	0	0	2	0
9	B	8	0	0	0	0
9	F	8	0	0	2	0
9	J	8	0	0	1	0
10	B	7	0	0	0	0
10	F	7	0	0	1	0
10	J	7	0	0	1	0
11	C	43	0	30	4	0
11	G	43	0	30	5	0
11	K	43	0	30	8	0
All	All	24855	0	24578	280	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 280 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:C:1129:HEM:HBC2	11:C:1129:HEM:HHD	1.39	1.04
11:K:1129:HEM:HBA2	11:K:1129:HEM:HHA	1.39	1.02
1:E:555:CYS:HA	1:E:571:VAL:HG23	1.50	0.94
11:K:1129:HEM:HBB2	4:L:26:LEU:HD13	1.51	0.90
1:I:555:CYS:HA	1:I:571:VAL:HG23	1.52	0.88

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	586/588 (100%)	567 (97%)	18 (3%)	1 (0%)	47 79
1	E	586/588 (100%)	563 (96%)	21 (4%)	2 (0%)	41 74
1	I	586/588 (100%)	562 (96%)	23 (4%)	1 (0%)	47 79
2	B	236/238 (99%)	221 (94%)	14 (6%)	1 (0%)	34 69
2	F	236/238 (99%)	220 (93%)	16 (7%)	0	100 100
2	J	236/238 (99%)	220 (93%)	14 (6%)	2 (1%)	19 58
3	C	119/129 (92%)	115 (97%)	4 (3%)	0	100 100
3	G	119/129 (92%)	115 (97%)	3 (2%)	1 (1%)	19 58
3	K	119/129 (92%)	115 (97%)	4 (3%)	0	100 100
4	D	103/115 (90%)	94 (91%)	8 (8%)	1 (1%)	15 54
4	H	103/115 (90%)	95 (92%)	7 (7%)	1 (1%)	15 54
4	L	103/115 (90%)	95 (92%)	6 (6%)	2 (2%)	8 39
All	All	3132/3210 (98%)	2982 (95%)	138 (4%)	12 (0%)	34 69

5 of 12 Ramachandran outliers are listed below:



Mol	Chain	Res	Type
2	J	56	ARG
2	B	56	ARG
4	H	41	GLY
4	L	41	GLY
4	D	41	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	473/473 (100%)	466 (98%)	7 (2%)	65 85
1	E	473/473 (100%)	464 (98%)	9 (2%)	57 81
1	I	473/473 (100%)	465 (98%)	8 (2%)	60 83
2	B	208/208 (100%)	198 (95%)	10 (5%)	25 61
2	F	208/208 (100%)	198 (95%)	10 (5%)	25 61
2	J	208/208 (100%)	198 (95%)	10 (5%)	25 61
3	C	101/109 (93%)	99 (98%)	2 (2%)	55 80
3	G	101/109 (93%)	99 (98%)	2 (2%)	55 80
3	K	101/109 (93%)	99 (98%)	2 (2%)	55 80
4	D	88/96 (92%)	85 (97%)	3 (3%)	37 70
4	H	88/96 (92%)	85 (97%)	3 (3%)	37 70
4	L	88/96 (92%)	85 (97%)	3 (3%)	37 70
All	All	2610/2658 (98%)	2541 (97%)	69 (3%)	46 76

5 of 69 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	1	MET
2	F	180	ARG
2	J	212	CYS
2	F	2	ARG
2	F	87	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 3 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	FAD	E	601	1	51,58,58	1.57	8 (15%)	60,89,89	2.01	14 (23%)
9	SF4	F	303	2	0,12,12	0.00	-	-	-	-
5	FAD	I	601	1	51,58,58	1.42	7 (13%)	60,89,89	1.58	8 (13%)
10	F3S	J	304	2	0,9,9	0.00	-	-	-	-
10	F3S	B	304	2	0,9,9	0.00	-	-	-	-
8	FES	B	302	2	0,4,4	0.00	-	-	-	-
8	FES	J	302	2	0,4,4	0.00	-	-	-	-
11	HEM	C	1129	3,4	27,50,50	2.28	6 (22%)	17,82,82	1.41	2 (11%)
8	FES	F	302	2	0,4,4	0.00	-	-	-	-
9	SF4	B	303	2	0,12,12	0.00	-	-	-	-
11	HEM	G	1129	3,4	27,50,50	2.32	5 (18%)	17,82,82	1.60	3 (17%)
6	TEO	I	1589	-	1,8,8	0.42	0	0,10,10	0.00	-
6	TEO	E	1589	-	1,8,8	0.58	0	0,10,10	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
9	SF4	J	303	2	0,12,12	0.00	-	-	-	-
11	HEM	K	1129	3,4	27,50,50	2.30	6 (22%)	17,82,82	1.81	5 (29%)
6	TEO	A	1589	-	1,8,8	0.34	0	0,10,10	0.00	-
10	F3S	F	304	2	0,9,9	0.00	-	-	-	-
5	FAD	A	601	1	51,58,58	1.53	9 (17%)	60,89,89	2.14	15 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FAD	E	601	1	-	7/30/50/50	0/6/6/6
10	F3S	J	304	2	-	-	0/3/3/3
9	SF4	F	303	2	-	-	0/6/5/5
5	FAD	I	601	1	-	2/30/50/50	0/6/6/6
10	F3S	B	304	2	-	-	0/3/3/3
8	FES	B	302	2	-	-	0/1/1/1
8	FES	J	302	2	-	-	0/1/1/1
11	HEM	C	1129	3,4	-	0/6/54/54	-
6	TEO	I	1589	-	-	1/2/8/8	-
8	FES	F	302	2	-	-	0/1/1/1
11	HEM	G	1129	3,4	-	0/6/54/54	-
6	TEO	E	1589	-	-	2/2/8/8	-
9	SF4	J	303	2	-	-	0/6/5/5
11	HEM	K	1129	3,4	-	2/6/54/54	-
6	TEO	A	1589	-	-	2/2/8/8	-
10	F3S	F	304	2	-	-	0/3/3/3
9	SF4	B	303	2	-	-	0/6/5/5
5	FAD	A	601	1	-	10/30/50/50	0/6/6/6

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	C	1129	HEM	C3B-C2B	-5.88	1.32	1.40
11	G	1129	HEM	C3B-C2B	-5.82	1.32	1.40
11	K	1129	HEM	C3B-C2B	-5.73	1.32	1.40
11	K	1129	HEM	C3D-C2D	5.26	1.53	1.37
5	A	601	FAD	C4X-N5	5.21	1.40	1.33

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	601	FAD	C1'-N10-C9A	6.05	123.05	118.29
5	I	601	FAD	C4-N3-C2	5.73	119.98	115.14
5	A	601	FAD	C4-C4X-N5	5.45	124.82	118.60
5	E	601	FAD	C4'-C3'-C2'	-5.37	102.19	113.36
5	E	601	FAD	N3A-C2A-N1A	-5.35	120.31	128.68

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	601	FAD	N10-C1'-C2'-O2'
5	E	601	FAD	N10-C1'-C2'-C3'
5	E	601	FAD	O4'-C4'-C5'-O5'
5	I	601	FAD	N10-C1'-C2'-O2'
6	I	1589	TEO	O2-C2-C3-C4


There are no ring outliers.


14 monomers are involved in 50 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	601	FAD	4	0
9	F	303	SF4	2	0
5	I	601	FAD	7	0
10	J	304	F3S	1	0
8	J	302	FES	2	0
11	C	1129	HEM	4	0
11	G	1129	HEM	5	0
6	I	1589	TEO	4	0
6	E	1589	TEO	4	0
9	J	303	SF4	1	0
11	K	1129	HEM	8	0
6	A	1589	TEO	2	0
10	F	304	F3S	1	0
5	A	601	FAD	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	588/588 (100%)	0.16	7 (1%) 79 67	54, 55, 56, 58	0
1	E	588/588 (100%)	0.24	16 (2%) 54 39	54, 55, 56, 58	0
1	I	588/588 (100%)	1.14	126 (21%) 0 1	54, 55, 56, 58	0
2	B	238/238 (100%)	0.21	8 (3%) 45 29	53, 55, 56, 57	0
2	F	238/238 (100%)	0.27	9 (3%) 40 26	54, 55, 56, 57	0
2	J	238/238 (100%)	0.67	27 (11%) 5 3	54, 55, 56, 57	0
3	C	121/129 (93%)	0.26	5 (4%) 37 24	54, 55, 56, 56	0
3	G	121/129 (93%)	0.46	14 (11%) 4 3	54, 55, 56, 56	0
3	K	121/129 (93%)	1.02	22 (18%) 1 1	54, 55, 56, 56	0
4	D	105/115 (91%)	0.07	5 (4%) 30 18	54, 55, 56, 57	0
4	H	105/115 (91%)	0.24	8 (7%) 13 7	54, 55, 56, 57	0
4	L	105/115 (91%)	0.11	7 (6%) 17 10	54, 55, 56, 57	0
All	All	3156/3210 (98%)	0.45	254 (8%) 12 6	53, 55, 56, 58	0

The worst 5 of 254 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	1	MET	7.8
1	I	338	ALA	7.4
1	I	334	SER	6.2
1	I	268	HIS	5.5
1	I	525	ALA	5.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	TEO	I	1589	9/9	0.89	0.26	122,123,124,124	0
5	FAD	I	601	53/53	0.90	0.23	70,80,90,93	0
6	TEO	E	1589	9/9	0.90	0.28	43,45,47,48	0
7	NA	I	1590	1/1	0.93	0.14	41,41,41,41	0
8	FES	J	302	4/4	0.94	0.10	70,71,72,73	0
6	TEO	A	1589	9/9	0.94	0.18	33,35,35,38	0
7	NA	E	1590	1/1	0.96	0.11	9,9,9,9	0
11	HEM	K	1129	43/43	0.96	0.23	54,55,65,65	0
5	FAD	E	601	53/53	0.96	0.14	26,40,50,53	0
11	HEM	G	1129	43/43	0.97	0.23	55,59,63,65	0
9	SF4	J	303	8/8	0.97	0.07	60,61,64,64	0
11	HEM	C	1129	43/43	0.97	0.17	43,46,55,58	0
10	F3S	B	304	7/7	0.97	0.08	43,45,49,49	0
5	FAD	A	601	53/53	0.97	0.12	17,21,30,40	0
7	NA	A	1590	1/1	0.98	0.08	2,2,2,2	0
10	F3S	F	304	7/7	0.98	0.09	47,48,50,53	0
10	F3S	J	304	7/7	0.98	0.11	61,64,66,69	0
9	SF4	B	303	8/8	0.99	0.10	28,28,30,33	0
8	FES	F	302	4/4	0.99	0.10	32,33,40,41	0
9	SF4	F	303	8/8	0.99	0.07	37,38,41,42	0
8	FES	B	302	4/4	0.99	0.14	30,31,34,35	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FAD I 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around HEM K 1129:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD E 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around HEM G 1129:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM C 1129:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around FAD A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers

There are no such residues in this entry.