

# wwPDB X-ray Structure Validation Summary Report (i)

#### Sep 25, 2023 – 09:00 AM EDT

PDB ID	:	5WEW
Title	:	Crystal structure of Klebsiella pneumoniae fosfomycin resistance protein (Fos-
		AKP) with inhibitor (ANY1) bound
Authors	:	Klontz, E.H.; Sundberg, E.J.
Deposited on	:	2017-07-10
Resolution	:	3.18  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)		
EDS	:	2.35.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 3.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ \textbf{(\#Entries)} \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	1467 (3.20-3.16)
Clashscore	141614	1599 (3.20-3.16)
Ramachandran outliers	138981	1574 (3.20-3.16)
Sidechain outliers	138945	1573 (3.20-3.16)
RSRZ outliers	127900	1423 (3.20-3.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	145	% • 78%	18%	
1	В	145	% • 81%	15%	
1	С	145	2% 	13%	• 5%
1	D	145	% 76%	19%	6%
1	Е	145	2% 66% 16%	18%	6



Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	F	145	% <b>7</b> 9%	15%	6%
1	G	145	% <b>81</b> %	14%	••
1	Н	145	% 82%	14%	• •



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 8654 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	140	Total	С	Ν	0	S	0	0	0
1	A	140	1072	689	183	194	6	0	0	0
1	В	141	Total	С	Ν	0	S	0	0	0
	D	141	1085	696	186	197	6	0	0	0
1	С	138	Total	С	Ν	0	S	0	0	0
	U	130	1052	677	178	191	6	0	0	0
1	D	137	Total	С	Ν	0	S	0	0	0
	D	137	1055	678	178	193	6	0		0
1	Е	119	Total	С	Ν	Ο	S	0	0	0
	Ľ	119	921	594	157	164	6	0	0	
1	F	137	Total	С	Ν	Ο	S	0	0	0
	Г	137	1047	674	177	191	5	0	0	0
1	G	141	Total	С	Ν	Ο	S	0	0	0
	G	141	1093	700	187	200	6	0	0	0
1	Н	141	Total	С	Ν	Ο	S	0	0	0
	11	141	1081	694	185	196	6		) 0	

• Molecule 1 is a protein called Fosfomycin resistance protein.

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	140	HIS	-	expression tag	UNP W8UNW6
А	141	HIS	-	expression tag	UNP W8UNW6
А	142	HIS	-	expression tag	UNP W8UNW6
А	143	HIS	-	expression tag	UNP W8UNW6
А	144	HIS	-	expression tag	UNP W8UNW6
А	145	HIS	-	expression tag	UNP W8UNW6
В	140	HIS	-	expression tag	UNP W8UNW6
В	141	HIS	-	expression tag	UNP W8UNW6
В	142	HIS	-	expression tag	UNP W8UNW6
В	143	HIS	-	expression tag	UNP W8UNW6
В	144	HIS	-	expression tag	UNP W8UNW6
В	145	HIS	-	expression tag	UNP W8UNW6
С	140	HIS	-	expression tag	UNP W8UNW6



		vious page	A	Comment	Defenence
Chain	Residue	Modelled	Actual	Comment	Reference
C	141	HIS	-	expression tag	UNP W8UNW6
C	142	HIS	-	expression tag	UNP W8UNW6
С	143	HIS	-	expression tag	UNP W8UNW6
С	144	HIS	-	expression tag	UNP W8UNW6
С	145	HIS	-	expression tag	UNP W8UNW6
D	140	HIS	-	expression tag	UNP W8UNW6
D	141	HIS	-	expression tag	UNP W8UNW6
D	142	HIS	-	expression tag	UNP W8UNW6
D	143	HIS	-	expression tag	UNP W8UNW6
D	144	HIS	-	expression tag	UNP W8UNW6
D	145	HIS	-	expression tag	UNP W8UNW6
E	140	HIS	-	expression tag	UNP W8UNW6
E	141	HIS	-	expression tag	UNP W8UNW6
E	142	HIS	-	expression tag	UNP W8UNW6
Е	143	HIS	-	expression tag	UNP W8UNW6
Е	144	HIS	-	expression tag	UNP W8UNW6
E	145	HIS	-	expression tag	UNP W8UNW6
F	140	HIS	-	expression tag	UNP W8UNW6
F	141	HIS	-	expression tag	UNP W8UNW6
F	142	HIS	-	expression tag	UNP W8UNW6
F	143	HIS	-	expression tag	UNP W8UNW6
F	144	HIS	-	expression tag	UNP W8UNW6
F	145	HIS	-	expression tag	UNP W8UNW6
G	140	HIS	-	expression tag	UNP W8UNW6
G	141	HIS	-	expression tag	UNP W8UNW6
G	142	HIS	-	expression tag	UNP W8UNW6
G	143	HIS	-	expression tag	UNP W8UNW6
G	144	HIS	-	expression tag	UNP W8UNW6
G	145	HIS	-	expression tag	UNP W8UNW6
Н	140	HIS	-	expression tag	UNP W8UNW6
Н	141	HIS	-	expression tag	UNP W8UNW6
Н	142	HIS	-	expression tag	UNP W8UNW6
Н	143	HIS	-	expression tag	UNP W8UNW6
Н	144	HIS	-	expression tag	UNP W8UNW6
Н	145	HIS	-	expression tag	UNP W8UNW6

Continued from previous page...

• Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


$\mathbf{N}$	ſol	Chain	Residues	Atoms	ZeroOcc	AltConf
	2	А	1	Total Mn 1 1	0	0
	2	В	1	Total Mn 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	1	Total Mn 1 1	0	0
2	D	1	Total Mn 1 1	0	0
2	Е	1	Total Mn 1 1	0	0
2	F	1	Total Mn 1 1	0	0
2	G	1	Total Mn 1 1	0	0
2	Н	1	Total Mn 1 1	0	0

• Molecule 3 is 6,6'-(4-nitro-1H-pyrazole-3,5-diyl)bis(3-bromopyrazolo[1,5-a]pyrimidin-2(1H)-one) (three-letter code: A81) (formula:  $C_{15}H_7Br_2N_9O_4$ ).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
3	Δ	1	Total	Br	С	Ν	Ο	0	0
0	А	1	30	2	15	9	4	0	0
3	В	1	Total	Br	С	Ν	0	0	0
0	D	1	30	2	15	9	4	0	0
3	С	1	Total	Br	С	Ν	0	0	0
0	U	1	30	2	15	9	4	0	0
3	Л	1	Total	Br	С	Ν	0	0	0
0	D	1	30	2	15	9	4	0	0
3	E	1	Total	Br	С	Ν	0	0	0
3	Ľ	1	30	2	15	9	4	U	0




Mol	Chain	Residues		Ate	$\mathbf{oms}$		ZeroOcc	AltConf		
3	Б	1	Total	Br	С	Ν	0	0	0	
0	Г	1	30	2	15	9	4	0	0	
3	Б	1	Total	Br	С	Ν	0	0	0	
0	Г	1	30	2	15	9	4	0	0	
2	С	1	Total	Br	С	Ν	Ο	0	0	
5	G	1	30	2	15	9	4	0	0	

Continued from previous page...



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




• Molecule 1: Fosfomycin resistance protein

HIS HIS HIS



• Molecule 1: Fosfomycin resistance protein





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants	120.40Å 197.62Å 117.03Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	37.75 - 3.18	Depositor
Resolution (A)	38.62 - 3.18	EDS
% Data completeness	89.0 (37.75-3.18)	Depositor
(in resolution range)	89.1 (38.62 - 3.18)	EDS
R <sub>merge</sub>	0.33	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.78 (at 3.18 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.10.1_2155	Depositor
$R, R_{free}$	0.221 , $0.262$	Depositor
$\Pi, \Pi_{free}$	0.223 , $0.262$	DCC
$R_{free}$ test set	1106 reflections $(5.19\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	39.3	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33, $30.3$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	0.004  for  1/2 *h-1/2 *k,-3/2 *h-1/2 *k,-1	Xtriage
Estimated twinning fraction	0.047 for $1/2$ *h $+1/2$ *k, $3/2$ *h $-1/2$ *k, $-1$	Attrage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	8654	wwPDB-VP
Average B, all atoms $(Å^2)$	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 21.30 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.2171e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MN,  $\operatorname{A81}$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.25	0/1103	0.43	0/1502	
1	В	0.25	0/1117	0.42	0/1521	
1	С	0.26	0/1082	0.44	0/1474	
1	D	0.26	0/1085	0.43	0/1477	
1	Ε	0.25	0/948	0.43	0/1290	
1	F	0.25	0/1077	0.43	0/1468	
1	G	0.26	0/1125	0.43	0/1531	
1	Н	0.25	0/1113	0.43	0/1516	
All	All	0.25	0/8650	0.43	0/11779	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1072	0	1014	18	0
1	В	1085	0	1023	15	0
1	С	1052	0	997	13	0
1	D	1055	0	1005	17	0
1	Е	921	0	877	16	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
			( /	( )		•
1	F	1047	0	989	14	0
1	G	1093	0	1033	19	0
1	Н	1081	0	1017	15	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
2	С	1	0	0	0	0
2	D	1	0	0	0	0
2	Ε	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	Н	1	0	0	0	0
3	А	30	0	0	0	0
3	В	30	0	0	0	0
3	С	30	0	0	0	0
3	D	30	0	0	0	0
3	Е	30	0	0	0	0
3	F	60	0	0	0	0
3	G	30	0	0	0	0
All	All	8654	0	7955	112	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 112 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2 Interatomic distance (Å)		Clash overlap (Å)
1:G:127:ARG:NH1	1:G:138:GLU:OE2	2.20	0.72
1:E:106:ASP:OD1	1:E:110:HIS:N	2.23	0.71
1:A:127:ARG:NH1	1:A:138:GLU:OE2	2.25	0.69
1:E:104:PHE:HE1	1:E:114:LEU:HG	1.62	0.64
1:H:72:ILE:HD13	1:H:77:PHE:HA	1.83	0.60

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	138/145~(95%)	137~(99%)	1 (1%)	0	100	100
1	В	139/145~(96%)	136~(98%)	2(1%)	1 (1%)	22	60
1	$\mathbf{C}$	136/145~(94%)	135~(99%)	1 (1%)	0	100	100
1	D	135/145~(93%)	134 (99%)	1 (1%)	0	100	100
1	Ε	115/145~(79%)	108 (94%)	7~(6%)	0	100	100
1	F	135/145~(93%)	134 (99%)	1 (1%)	0	100	100
1	G	139/145~(96%)	137~(99%)	2(1%)	0	100	100
1	Н	139/145~(96%)	138 (99%)	1 (1%)	0	100	100
All	All	1076/1160~(93%)	1059 (98%)	16 (2%)	1 (0%)	51	83

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	140	HIS

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	107/119~(90%)	104~(97%)	3~(3%)	43 73		
1	В	109/119~(92%)	106~(97%)	3~(3%)	43 73		
1	$\mathbf{C}$	105/119~(88%)	103~(98%)	2(2%)	57 80		
1	D	107/119~(90%)	106~(99%)	1 (1%)	78 91		
1	Ε	95/119~(80%)	95~(100%)	0	100 100		
1	F	104/119~(87%)	101~(97%)	3~(3%)	42 72		
1	G	111/119~(93%)	109~(98%)	2(2%)	59 81		
1	Н	108/119~(91%)	105~(97%)	3~(3%)	43 73		
All	All	846/952~(89%)	829~(98%)	17~(2%)	55 79		



5	of	17	residues	with	a	non-rotamerie	С	sidechain	are	listed	below:

Mol	Chain	$\mathbf{Res}$	Type
1	Н	35	ASP
1	Н	106	ASP
1	С	106	ASP
1	D	35	ASP
1	F	35	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Link Bond lengths			Bond angles		
	туре		Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ
3	A81	F	202	2	23,34,34	1.60	7 (30%)	12,52,52	1.09	1 (8%)
3	A81	D	202	2	23,34,34	1.60	7 (30%)	12,52,52	1.03	1 (8%)
3	A81	Е	202	2	23,34,34	1.60	7 (30%)	12,52,52	1.01	1 (8%)
3	A81	G	202	2	23,34,34	1.61	7 (30%)	12,52,52	1.24	1 (8%)



Mol	l Type Chain Res Link				Bo	Bond lengths			Bond angles		
10101	туре	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
3	A81	В	202	2	23,34,34	1.59	7 (30%)	12,52,52	1.09	1 (8%)	
3	A81	А	202	2	23,34,34	1.61	7 (30%)	12,52,52	0.99	1 (8%)	
3	A81	С	202	2	23,34,34	1.59	7 (30%)	12,52,52	0.98	0	
3	A81	F	203	2	23,34,34	1.63	7 (30%)	12,52,52	0.97	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A81	F	202	2	-	0/0/12/12	0/5/5/5
3	A81	D	202	2	-	0/0/12/12	0/5/5/5
3	A81	Е	202	2	-	0/0/12/12	0/5/5/5
3	A81	G	202	2	-	0/0/12/12	0/5/5/5
3	A81	В	202	2	-	0/0/12/12	0/5/5/5
3	A81	А	202	2	-	0/0/12/12	0/5/5/5
3	A81	С	202	2	-	0/0/12/12	0/5/5/5
3	A81	F	203	2	-	0/0/12/12	0/5/5/5

The worst 5 of 56 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
3	F	203	A81	BR2-C18	2.94	1.94	1.88
3	D	202	A81	BR-C29	2.94	1.94	1.88
3	В	202	A81	BR-C29	2.89	1.94	1.88
3	G	202	A81	BR2-C18	2.88	1.94	1.88
3	С	202	A81	BR-C29	2.86	1.94	1.88

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	G	202	A81	C05-C04-C08	-2.87	105.35	107.46
3	F	202	A81	C05-C04-C08	-2.38	105.71	107.46
3	В	202	A81	C05-C04-C08	-2.12	105.90	107.46
3	А	202	A81	C05-C04-C08	-2.08	105.93	107.46
3	D	202	A81	BR-C29-C27	2.07	127.53	120.73

There are no chirality outliers.


There are no torsion outliers.




There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	140/145~(96%)	-0.32	1 (0%) 87 81	21, 33, 49, 57	0
1	В	141/145~(97%)	-0.30	2 (1%) 75 63	24, 34, 51, 70	0
1	С	138/145~(95%)	-0.21	3 (2%) 62 48	22, 31, 50, 72	0
1	D	137/145~(94%)	-0.18	1 (0%) 87 81	22, 37, 56, 71	0
1	Ε	119/145~(82%)	0.29	3 (2%) 57 43	31, 58, 75, 92	0
1	F	137/145~(94%)	0.16	2 (1%) 73 61	34, 52, 70, 78	0
1	G	141/145~(97%)	-0.10	2 (1%) 75 63	30, 42, 59, 69	0
1	Н	141/145~(97%)	-0.17	1 (0%) 87 81	29, 42, 62, 80	0
All	All	1094/1160~(94%)	-0.11	15 (1%) 75 63	21, 41, 64, 92	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	1	MET	3.5
1	Ε	24	GLN	3.0
1	G	61	GLU	3.0
1	Е	124	ALA	3.0
1	F	124	ALA	2.9

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
3	A81	Е	202	30/30	0.96	0.18	$39{,}53{,}60{,}63$	0
3	A81	F	203	30/30	0.96	0.17	42,49,62,68	0
3	A81	С	202	30/30	0.97	0.14	$27,\!36,\!45,\!52$	0
3	A81	D	202	30/30	0.97	0.14	27,38,45,53	0
2	MN	F	201	1/1	0.97	0.12	25,25,25,25	0
3	A81	F	202	30/30	0.97	0.17	33,42,52,54	0
2	MN	Н	201	1/1	0.97	0.11	26,26,26,26	0
3	A81	G	202	30/30	0.97	0.16	$36,\!47,\!56,\!59$	0
3	A81	А	202	30/30	0.98	0.13	28,34,42,48	0
3	A81	В	202	30/30	0.98	0.13	22,31,42,48	0
2	MN	А	201	1/1	0.98	0.15	21,21,21,21	0
2	MN	В	201	1/1	0.98	0.15	17,17,17,17	0
2	MN	Е	201	1/1	0.99	0.10	47,47,47,47	0
2	MN	С	201	1/1	0.99	0.10	24,24,24,24	0
2	MN	G	201	1/1	0.99	0.14	23,23,23,23	0
2	MN	D	201	1/1	0.99	0.12	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















## 6.5 Other polymers (i)

There are no such residues in this entry.

