

# Full wwPDB X-ray Structure Validation Report (i)

#### Oct 8, 2023 – 06:15 PM EDT

PDB ID	:	6WHK
Title	:	Minimally mutated anti-influenza broadly neutralizing antibody
Authors	:	Bajic, G.; Schmidt, A.G.
Deposited on		
Resolution	:	2.60  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Xtriage (Phenix) EDS buster-report Percentile statistics Refmac CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA)	: : : : :	20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove) Engh & Huber (2001) Parkinson et al. (1996)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	230	2% <b>8</b> 0%	11%	8%
2	С	240	75%	18%	8%
3	В	219	76%	20%	·
4	D	2	100%		



#### 6 WHK

## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5075 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	211	Total 1706	C 1088	N 292	O 322	$\frac{S}{4}$	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	46	ALA	-	expression tag	UNP B4UPF7
А	47	SER	-	expression tag	UNP B4UPF7
А	268	GLY	-	expression tag	UNP B4UPF7
А	269	SER	-	expression tag	UNP B4UPF7
А	270	LEU	-	expression tag	UNP B4UPF7
А	271	GLU	-	expression tag	UNP B4UPF7
А	272	VAL	-	expression tag	UNP B4UPF7
A	273	LEU	-	expression tag	UNP B4UPF7
А	274	PHE	-	expression tag	UNP B4UPF7
А	275	GLN	-	expression tag	UNP B4UPF7

• Molecule 2 is a protein called Fab Heavy Chain.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	C	222	Total 1678	C 1069	N 282	0 321	S 6	0	0	0

• Molecule 3 is a protein called Fab Light Chain.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	В	211	Total 1557	C 976	N 260	0 317	$\frac{S}{4}$	0	0	0


• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	A	Aton	ns		ZeroOcc	AltConf	Trace
4	D	2	Total 28	C 16	N 2	0 10	0	0	0

• Molecule 5 is 2-{2-[2-(2-{2-[2-(2-ETHOXY-ETHOXY)-ETHOXY]-ETHOXY}PA



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	Δ	1	Total	С	0	0	0
5	A	1	24	16	8	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	38	Total         O           38         38	0	0
6	С	27	TotalO2727	0	0
6	В	17	Total O 17 17	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Hemagglutinin HA1 chain

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

100%



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	69.57Å 76.33Å 126.17Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	63.08 - 2.60	Depositor
Resolution (A)	65.31 - 2.60	EDS
% Data completeness	99.9 (63.08-2.60)	Depositor
(in resolution range)	$100.0\ (65.31-2.60)$	EDS
R <sub>merge</sub>	0.16	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.53 (at 2.62 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
D D.	0.218 , $0.268$	Depositor
$R, R_{free}$	0.218 , $0.267$	DCC
$R_{free}$ test set	1057 reflections $(4.96%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	48.5	Xtriage
Anisotropy	0.432	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34 , 57.7	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5075	wwPDB-VP
Average B, all atoms $(Å^2)$	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PE4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.27	0/1759	0.45	0/2397	
2	С	0.28	0/1719	0.49	0/2340	
3	В	0.29	0/1595	0.50	0/2178	
All	All	0.28	0/5073	0.48	0/6915	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1706	0	1616	16	0
2	С	1678	0	1656	26	0
3	В	1557	0	1518	33	0
4	D	28	0	25	0	0
5	А	24	0	34	0	0
6	А	38	0	0	1	0
6	В	17	0	0	1	0
6	С	27	0	0	1	0
All	All	5075	0	4849	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 7.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
3:B:194:ARG:HH12	3:B:213:PRO:C	1.84	0.81
1:A:58:ASN:ND2	6:A:401:HOH:O	2.15	0.80
2:C:173:LEU:HD21	2:C:196:VAL:HG21	1.62	0.79
3:B:149:VAL:HG12	3:B:202:HIS:HB2	1.72	0.70
1:A:78:GLU:HG2	1:A:115:PHE:H	1.55	0.70
2:C:137:PRO:HD3	2:C:223:LYS:HE2	1.74	0.70
3:B:62:ARG:NH1	3:B:78:GLY:O	2.25	0.69
3:B:187:PRO:O	3:B:191:LYS:NZ	2.29	0.66
1:A:86:THR:HB	1:A:87:PRO:HD2	1.79	0.65
2:C:74:ARG:NH1	6:C:303:HOH:O	2.29	0.65
3:B:154:LYS:HG2	3:B:197:SER:HB2	1.78	0.65
3:B:194:ARG:NH1	3:B:213:PRO:C	2.50	0.65
1:A:112:VAL:HG11	1:A:115:PHE:HB2	1.78	0.64
3:B:118:PRO:HD3	3:B:202:HIS:HD2	1.62	0.63
1:A:114:SER:HB3	1:A:261:SER:HB2	1.81	0.62
3:B:14:PRO:HA	3:B:79:LEU:HB3	1.83	0.59
1:A:176:VAL:HG22	1:A:237:LEU:HB3	1.85	0.58
2:C:173:LEU:HD12	2:C:175:SER:OG	2.03	0.58
3:B:113:GLN:NE2	3:B:145:TYR:CE1	2.71	0.58
3:B:113:GLN:NE2	3:B:145:TYR:CZ	2.72	0.57
2:C:20:LEU:HB2	2:C:83:LEU:HB3	1.85	0.56
3:B:118:PRO:HD3	3:B:202:HIS:CD2	2.40	0.56
3:B:146:PRO:HB2	3:B:202:HIS:HE1	1.70	0.56
3:B:28:ASN:O	3:B:32:ASN:HB2	2.05	0.56
2:C:20:LEU:HD12	2:C:83:LEU:HD23	1.87	0.56
3:B:141:ILE:HG21	3:B:200:VAL:HG11	1.89	0.55
3:B:141:ILE:O	3:B:179:ALA:N	2.38	0.55
3:B:164:VAL:HG22	3:B:183:LEU:HD12	1.89	0.54
1:A:188:ILE:HD11	1:A:198:GLU:HG2	1.90	0.53
2:C:133:PRO:HB3	2:C:159:TYR:HB3	1.90	0.53
3:B:14:PRO:HD3	3:B:111:LEU:O	2.08	0.53
3:B:194:ARG:O	3:B:194:ARG:HG3	2.09	0.53
3:B:38:GLN:HB2	3:B:48:LEU:HD11	1.91	0.52
3:B:170:SER:OG	3:B:171:LYS:N	2.40	0.52
3:B:167:THR:HG21	3:B:179:ALA:HA	1.94	0.50
2:C:102:TRP:HB2	2:C:111:ALA:HB3	1.94	0.50
2:C:13:GLN:HB2	2:C:16:ARG:HG3	1.95	0.49
1:A:113:SER:H	1:A:263:GLY:H	1.61	0.48

Continued on next page...



Continued from previous page Interatomic Clash									
Atom-1	Atom-2	distance (Å)	overlap (Å)						
2:C:205:THR:HG22	2:C:206:GLN:OE1	2.15	0.47						
3:B:41:PRO:HG2	3:B:171:LYS:HB2	1.97	0.46						
2:C:11:LEU:HD11	2:C:126:SER:HB3	1.98	0.46						
3:B:79:LEU:HD12	3:B:79:LEU:HA	1.80	0.46						
2:C:8:GLY:O	2:C:18:LEU:HD21	2.17	0.45						
2:C:99:THR:HB	2:C:114:LEU:HD12	1.99	0.45						
3:B:166:THR:HG23	3:B:181:SER:HB2	2.00	0.44						
3:B:190:TRP:CZ2	3:B:213:PRO:HA	2.53	0.43						
1:A:146:SER:OG	1:A:147:PHE:N	2.49	0.43						
2:C:50:PHE:CE2	2:C:52:ARG:HG3	2.53	0.43						
2:C:6:GLU:H	2:C:119:GLN:HE22	1.66	0.43						
1:A:194:ILE:HA	2:C:107:ILE:HG21	2.00	0.42						
3:B:36:TRP:CD2	3:B:74:LEU:HB2	2.55	0.42						
2:C:201:SER:O	2:C:205:THR:OG1	2.37	0.42						
2:C:66:VAL:HG13	2:C:70:PHE:CG	2.55	0.42						
2:C:36:TRP:CE2	2:C:83:LEU:HB2	2.55	0.41						
3:B:116:GLY:O	3:B:202:HIS:NE2	2.53	0.41						
1:A:108:GLN:OE1	1:A:262:ARG:HD2	2.20	0.41						
2:C:47:TRP:CG	3:B:100:VAL:HB	2.55	0.41						
1:A:77:LYS:HG2	1:A:78:GLU:N	2.35	0.41						
1:A:114:SER:O	1:A:260:LEU:HA	2.20	0.41						
2:C:159:TYR:CE1	2:C:190:TYR:HB2	2.56	0.41						
2:C:200:SER:O	2:C:203:LEU:HG	2.21	0.41						
3:B:93:ASP:HB3	3:B:98:ALA:HB3	2.03	0.41						
1:A:70:GLU:HB2	1:A:91:ASN:HD22	1.86	0.41						
3:B:33:TYR:CD1	3:B:51:ASP:HA	2.55	0.41						
3:B:194:ARG:HD2	3:B:213:PRO:HD2	2.03	0.41						
2:C:176:GLY:O	2:C:196:VAL:HA	2.20	0.41						
3:B:213:PRO:O	6:B:301:HOH:O	2.22	0.41						
2:C:165:THR:HG1	2:C:213:ASN:HB3	1.86	0.40						
2:C:202:SER:HA	2:C:205:THR:OG1	2.20	0.40						
2:C:152:LEU:HG	2:C:196:VAL:HG13	2.03	0.40						
1:A:81:SER:O	1:A:111:SER:HB2	2.22	0.40						
1:A:213:PHE:O	1:A:215:PRO:HD3	2.22	0.40						
3:B:113:GLN:NE2	3:B:145:TYR:OH	2.54	0.40						

Continued from previous page...

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	209/230~(91%)	205~(98%)	4(2%)	0	100 100
2	$\mathbf{C}$	218/240~(91%)	215~(99%)	3~(1%)	0	100 100
3	В	209/219~(95%)	200 (96%)	9 (4%)	0	100 100
All	All	636/689~(92%)	620 (98%)	16 (2%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	187/200~(94%)	184 (98%)	3~(2%)	62 82
2	С	185/202~(92%)	183~(99%)	2(1%)	73 88
3	В	175/182~(96%)	170 (97%)	5(3%)	42 68
All	All	547/584~(94%)	537~(98%)	10 (2%)	59 80

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	111	SER
1	А	140	SER
1	А	261	SER
2	С	220	LYS
2	С	224	ARG

Continued on next page...



Continued from previous page...

	Ű	-	10
Mol	Chain	$\mathbf{Res}$	Type
3	В	11	SER
3	В	17	LYS
3	В	23	SER
3	В	170	SER
3	В	175	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	С	178	HIS
3	В	113	GLN
3	В	174	ASN
3	В	175	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bo	ond leng	$\mathbf{ths}$	B	ond ang	les
WIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
4	NAG	D	1	1,4	14,14,15	0.34	0	17,19,21	0.45	0
4	NAG	D	2	4	14,14,15	0.30	0	17,19,21	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	2	NAG	O5-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
4	D	2	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Type	Chain	Res	Link	Bond lengths			Bond angles		
	туре			LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
5	PE4	А	303	-	23,23,23	0.55	0	22,22,22	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PE4	А	303	-	-	13/21/21/21	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	А	303	PE4	O7-C13-C14-O8
5	А	303	PE4	O6-C11-C12-O7
5	А	303	PE4	O3-C5-C6-O4
5	А	303	PE4	C12-C11-O6-C10
5	А	303	PE4	O2-C3-C4-O3
5	А	303	PE4	C5-C6-O4-C7
5	А	303	PE4	C7-C8-O5-C9
5	А	303	PE4	C9-C10-O6-C11
5	А	303	PE4	O1-C1-C2-O2
5	А	303	PE4	C14-C13-O7-C12
5	А	303	PE4	O6-C10-C9-O5
5	А	303	PE4	C16-C15-O8-C14
5	А	303	PE4	O4-C7-C8-O5

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient must be highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	$\# \mathbf{RS}$	RZ>	>2	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	А	211/230~(91%)	0.00	4 (1%)	66	62	27, 44, 80, 108	0
2	С	222/240~(92%)	0.11	7 (3%)	47	40	29, 46, 91, 105	0
3	В	211/219 (96%)	0.27	7 (3%)	46	39	34, 59, 89, 96	0
All	All	644/689~(93%)	0.13	18 (2%)	53	46	27, 49, 88, 108	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	С	171	GLY	4.9
2	С	205	THR	4.5
2	С	172	ALA	4.2
1	А	87	PRO	4.0
2	С	203	LEU	3.6
1	А	88	ASN	3.4
3	В	145	TYR	3.2
1	А	89	PRO	3.1
3	В	117	ALA	2.5
3	В	96	LEU	2.5
1	А	73	SER	2.4
3	В	161	LYS	2.4
3	В	147	GLY	2.3
2	С	173	LEU	2.2
3	В	13	ALA	2.1
2	С	1	GLU	2.1
2	С	224	ARG	2.0
3	В	41	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.




### 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

M	ol Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
4	NAG	D	2	14/15	0.61	0.35	55,78,99,100	0
4	NAG	D	1	14/15	0.77	0.20	41,74,87,89	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



### 6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q < 0.9
5	PE4	А	303	24/24	0.83	0.17	$32,\!62,\!76,\!78$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



### 6.5 Other polymers (i)

There are no such residues in this entry.

