

Full wwPDB X-ray Structure Validation Report (i)

Aug 9, 2020 – 09:35 PM BST

PDB ID	:	3WN5
Title	:	Crystal structure of asymmetrically engineered Fc variant in complex with
		FcgRIIIa
Authors	:	Kadono, S.; Mimoto, F.; Katada, H.; Igawa, T.; Kamikawa, T.; Hattori, K.
Deposited on	:	2013-12-05
Resolution	:	2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575(2.80-2.76)
Ramachandran outliers	138981	4487(2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
-1	Δ	000	2%	
	A	230	85%	7% 8%
	5		% •	
1	D	230	86%	6% 8%
	_		2%	
2	В	230	78%	12% • 9%
			5%	
2	E	230	83%	8% 9%
			2%	
3	C	197	80%	6% 14%
			2%	
3	F	197	80%	6% 14%



Mol	Chain	Length	Quality of chain							
4	G	9	44%	33%	22%					
4	Н	9	11%	78%	11%					
4	L	9	56%		44%					
5	Ι	5	60%		40%					
6	J	2		100%						
6	М	2		100%						
7	K	8	38%	63%						



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 9947 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	010	Total	С	Ν	Ο	S	0	0	0
	A		1664	1057	277	323	7			
1	р	919	Total	С	Ν	Ο	S	0	0	0
			1646	1046	274	319	7		U	

• Molecule 1 is a protein called Ig gamma-1 chain C region.

Chain	Residue	Modelled	Actual	Actual Comment	
А	220	SER	CYS	engineered mutation	UNP P01857
А	270	GLU	ASP	engineered mutation	UNP P01857
А	326	ASP	LYS	engineered mutation	UNP P01857
А	330	LYS	ALA	engineered mutation	UNP P01857
А	334	GLU	LYS	engineered mutation	UNP P01857
А	356	CYS	ASP	engineered mutation	UNP P01857
А	366	SER	THR	engineered mutation	UNP P01857
А	368	ALA	LEU	engineered mutation	UNP P01857
А	407	VAL	TYR	engineered mutation	UNP P01857
D	220	SER	CYS	engineered mutation	UNP P01857
D	270	GLU	ASP	engineered mutation	UNP P01857
D	326	ASP	LYS	engineered mutation	UNP P01857
D	330	LYS	ALA	engineered mutation	UNP P01857
D	334	GLU	LYS	engineered mutation	UNP P01857
D	356	CYS	ASP	engineered mutation	UNP P01857
D	366	SER	THR	engineered mutation	UNP P01857
D	368	ALA	LEU	engineered mutation	UNP P01857
D	407	VAL	TYR	engineered mutation	UNP P01857

There are 18 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	В	210	Total 1664	m C 1065	N 271	O 320	S 8	0	0	0



Continued from previous page...

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
2	Е	210	Total	C	N	0	S	0	0	0
			1644	1056	267	313	8			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	220	SER	CYS	engineered mutation	UNP P01857
В	234	TYR	LEU	engineered mutation	UNP P01857
В	235	TYR	LEU	engineered mutation	UNP P01857
В	236	TRP	GLY	engineered mutation	UNP P01857
В	239	MET	SER	engineered mutation	UNP P01857
В	268	ASP	HIS	engineered mutation	UNP P01857
В	298	ALA	SER	engineered mutation	UNP P01857
В	327	ASP	ALA	engineered mutation	UNP P01857
В	349	CYS	TYR	engineered mutation	UNP P01857
В	366	TRP	THR	engineered mutation	UNP P01857
Е	220	SER	CYS	engineered mutation	UNP P01857
Е	234	TYR	LEU	engineered mutation	UNP P01857
Е	235	TYR	LEU	engineered mutation	UNP P01857
Е	236	TRP	GLY	engineered mutation	UNP P01857
Е	239	MET	SER	engineered mutation	UNP P01857
Е	268	ASP	HIS	engineered mutation	UNP P01857
Е	298	ALA	SER	engineered mutation	UNP P01857
Е	327	ASP	ALA	engineered mutation	UNP P01857
Е	349	CYS	TYR	engineered mutation	UNP P01857
E	366	TRP	THR	engineered mutation	UNP P01857

• Molecule 3 is a protein called Low affinity immunoglobulin gamma Fc region receptor III-A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	С	160	Total	С	Ν	Ο	S	0	0	0	0
່ <u>ບ</u>		109	1358	867	232	255	4		0	0	
2	Б	170	Total	С	Ν	Ο	S	0	0	0	
0	Г	170	1354	862	230	258	4			U	

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	35	GLN	ASN	engineered mutation	UNP P08637
С	71	GLN	ASN	engineered mutation	UNP P08637
С	155	VAL	PHE	SEE REMARK 999	UNP P08637
С	166	GLN	ASN	engineered mutation	UNP P08637



3WN5	
------	--

Chain	Residue	Modelled	Actual	Comment	Reference
С	188	HIS	-	expression tag	UNP P08637
С	189	HIS	-	expression tag	UNP P08637
С	190	HIS	-	expression tag	UNP P08637
С	191	HIS	-	expression tag	UNP P08637
С	192	HIS	-	expression tag	UNP P08637
С	193	HIS	-	expression tag	UNP P08637
F	35	GLN	ASN	engineered mutation	UNP P08637
F	71	GLN	ASN	engineered mutation	UNP P08637
F	155	VAL	PHE	SEE REMARK 999	UNP P08637
F	166	GLN	ASN	engineered mutation	UNP P08637
F	188	HIS	-	expression tag	UNP P08637
F	189	HIS	-	expression tag	UNP P08637
F	190	HIS	-	expression tag	UNP P08637
F	191	HIS	-	expression tag	UNP P08637
F	192	HIS	-	expression tag	UNP P08637
F	193	HIS	-	expression tag	UNP P08637

• Molecule 4 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-gluc opyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	G	9	Total C N O 110 62 4 44	0	0	0
4	Н	9	Total C N O 110 62 4 44	0	0	0
4	L	9	Total C N O 110 62 4 44	0	0	0

• Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyran ose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
5	Ι	5	Total 61	С 34	N 2	O 25	0	0	0

• Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
6	J	2	Total C 28 16	C N 5 2	O 10	0	0	0
6	М	2	Total C 28 16	C N 5 2	O 10	0	0	0

• Molecule 7 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alp ha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-man nopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
7	K	8	Total C N O 99 56 4 39	0	0	0

• Molecule 8 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	D	1	Total I 1 1	0	0
8	Е	4	Total I 4 4	0	0
8	В	3	Total I 3 3	0	0
8	С	2	Total I 2 2	0	0
8	А	2	Total I 2 2	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	F	4	Total I 4 4	0	0

• Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	F	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	А	7	Total O 7 7	0	0
10	В	11	Total O 11 11	0	0
10	С	6	Total O 6 6	0	0
10	D	4	$\begin{array}{cc} \text{Total} & \text{O} \\ 4 & 4 \end{array}$	0	0
10	Ε	3	Total O 3 3	0	0
10	F	10	Total O 10 10	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Ig gamma-1 chain C region

• Molecule 3: Low affinity immunoglobulin gamma Fc region receptor III-A



• Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:	44%	33%	22%
IAG 1 IAG 2 IAG 2 IAN 4 IAN 7 IAN 7			

 $\label{eq:main_optimal_states} \bullet \mbox{Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose$

Chain H:	11%	78%	11%
NAG1 NAG2 BMA3 MAN4 NAG5 GAL6 MAN7 NAG8	4 months and a month and a		

• Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:	56%	44%
NAG1 NAG2 BMA3 MAN4 NAG5 GAL6 MAN7 NAG8 FUL9		

 $\label{eq:mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]} beta-D-mannopyranose-(1-6)] beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2$

Chain I:

60%

40%



NAG1 NAG2 BMA3 MAN4 MAN5

• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:

100%

100%

NAG1 NAG2

• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:

NAG1 NAG2

 $\label{eq:constraint} \bullet \mbox{Molecule 7: } 2\mbox{-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-gl$

Chain K:	38%	63%
NAG1 NAG2 MAA3 MAN4 MAN6 NAG5 FUL8		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	75.03Å 72.49Å 163.48Å	Deneiten
a, b, c, α , β , γ	90.00° 91.15° 90.00°	Depositor
$\mathbf{P}_{\text{acclution}}(\hat{\mathbf{A}})$	25.00 - 2.78	Depositor
Resolution (A)	44.51 - 2.78	EDS
% Data completeness	98.1 (25.00-2.78)	Depositor
(in resolution range)	98.1(44.51-2.78)	EDS
R _{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.03 (at 2.77 \text{\AA})$	Xtriage
Refinement program	REFMAC	Depositor
B B.	0.236 , 0.274	Depositor
It, Itfree	0.236 , 0.274	DCC
R_{free} test set	2192 reflections (5.03%)	wwPDB-VP
Wilson B-factor $(Å^2)$	54.5	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , 44.0	EDS
L-test for $twinning^2$	$ L > = 0.49, < L^2 > = 0.32$	Xtriage
	0.015 for k,h,-l	
Estimated twinning fraction	0.016 for -k,-h,-l	Xtriage
	0.020 for h,-k,-l	
F_o, F_c correlation	0.91	EDS
Total number of atoms	9947	wwPDB-VP
Average B, all atoms $(Å^2)$	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, GAL, FUL, IOD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.28	0/1709	0.46	0/2334	
1	D	0.28	0/1691	0.46	0/2313	
2	В	0.28	0/1713	0.46	0/2343	
2	Е	0.28	0/1693	0.44	0/2318	
3	С	0.30	0/1395	0.49	0/1897	
3	F	0.29	0/1391	0.48	0/1895	
All	All	0.28	0/9592	0.46	0/13100	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1664	0	1594	8	0
1	D	1646	0	1567	11	0
2	В	1664	0	1580	16	0
2	Е	1644	0	1547	11	0
3	С	1358	0	1299	7	0
3	F	1354	0	1273	4	0
4	G	110	0	94	1	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Н	110	0	94	2	0
4	L	110	0	94	0	0
5	Ι	61	0	52	0	0
6	J	28	0	25	0	0
6	М	28	0	25	0	0
7	K	99	0	85	0	0
8	А	2	0	0	0	0
8	В	3	0	0	1	0
8	С	2	0	0	0	0
8	D	1	0	0	0	0
8	Ε	4	0	0	1	0
8	F	4	0	0	1	0
9	F	14	0	13	0	0
10	А	7	0	0	0	0
10	В	11	0	0	0	0
10	С	6	0	0	0	0
10	D	4	0	0	0	0
10	Е	3	0	0	0	0
10	F	10	0	0	0	0
All	All	9947	0	$93\overline{42}$	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:255:ARG:HG2	1:D:255:ARG:HH11	1.47	0.80
1:D:255:ARG:CG	1:D:255:ARG:HH11	2.04	0.71
2:B:296:TYR:H	4:H:9:FUL:H61	1.57	0.69
3:F:152:ARG:HB3	3:F:161:SER:HB3	1.75	0.68
2:E:351:LEU:HB2	2:E:366:TRP:HB2	1.81	0.61
2:B:276:ASN:HB2	2:B:322:LYS:HB3	1.83	0.61
1:D:367:CYS:HB2	1:D:381:TRP:CZ2	2.39	0.58
4:G:5:NAG:H4	4:G:6:GAL:O2	2.03	0.58
3:C:65:GLU:HG2	3:C:80:GLN:HE22	1.69	0.57
1:A:351:LEU:HB2	1:A:366:SER:HB2	1.85	0.57
1:D:276:ASN:HB2	1:D:322:LYS:HB3	1.87	0.56
2:B:318:GLU:HG3	2:B:335:THR:CG2	2.36	0.56
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.88	0.55
1:D:255:ARG:CG	1:D:255:ARG:NH1	2.70	0.54



Interatomic Clash					
Atom-1	Atom-2	distance $(Å)$	overlap (Å)		
1:A:367:CYS:HB2	1:A:381:TRP:CZ2	2.43	0.54		
2:E:276:ASN:HB2	2:E:322:LYS:HB3	1.91	0.52		
2:B:351:LEU:HB2	2:B:366:TRP:HB2	1.94	0.50		
2:E:388:GLU:HG2	2:E:410:LEU:HD11	1.94	0.50		
2:E:429:HIS:CD2	2:E:431:ALA:H	2.30	0.49		
4:H:2:NAG:H82	4:H:9:FUL:H2	1.94	0.49		
2:B:361:ASN:N	2:B:361:ASN:OD1	2.37	0.49		
2:E:367:CYS:HB2	2:E:381:TRP:CZ2	2.47	0.48		
2:B:429:HIS:CD2	2:B:431:ALA:H	2.31	0.48		
2:B:355:ARG:HA	2:B:358:LEU:HD12	1.96	0.48		
2:B:245:PRO:HD3	2:B:259:VAL:HG22	1.96	0.48		
1:A:258:GLU:HG2	1:A:307:THR:HG22	1.96	0.48		
2:B:346:PRO:HB3	2:B:372:PHE:HB3	1.95	0.48		
2:E:379:VAL:HG22	2:E:427:VAL:HG22	1.95	0.48		
2:B:414:LYS:HE3	2:B:418:GLN:HE21	1.79	0.47		
1:A:414:LYS:O	1:A:418:GLN:HG2	2.14	0.47		
1:D:278:TYR:HB2	1:D:320:LYS:HB3	1.95	0.47		
1:A:266:VAL:HB	1:A:300:TYR:HB2	1.96	0.47		
2:B:350:THR:OG1	8:B:1010:IOD:I	3.01	0.47		
2:E:254:SER:OG	8:E:1011:IOD:I	2.93	0.47		
3:F:87:TRP:CZ3	3:F:155:VAL:HG11	2.49	0.46		
3:C:118:VAL:HG21	3:C:133:ASN:HA	1.98	0.46		
1:D:253:ILE:HD12	1:D:253:ILE:H	1.81	0.45		
1:A:236:GLY:HA2	3:C:155:VAL:HG12	1.99	0.45		
3:C:65:GLU:HG2	3:C:80:GLN:NE2	2.32	0.45		
3:F:26:CYS:HB2	3:F:39:TRP:CZ2	2.52	0.44		
2:E:429:HIS:HD2	2:E:431:ALA:H	1.65	0.44		
2:E:261:CYS:HB2	2:E:277:TRP:CH2	2.52	0.43		
2:B:240:VAL:HG22	2:B:263:VAL:HG22	2.01	0.43		
2:B:368:LEU:HD13	2:B:407:TYR:CZ	2.52	0.43		
3:C:89:LEU:HD23	3:C:91:GLN:HE21	1.84	0.43		
3:C:150:PHE:HB3	3:C:164:THR:HG22	2.01	0.43		
3:C:18:GLU:O	3:C:19:LYS:HB2	2.17	0.43		
1:D:261:CYS:HB2	1:D:277:TRP:CZ2	2.54	0.42		
2:E:391:TYR:HB3	2:E:410:LEU:HD12	2.00	0.42		
1:A:362:GLN:HG2	1:A:413:ASP:HA	2.02	0.41		
1:D:394:THR:HG22	2:E:397:VAL:HG21	2.02	0.41		
3:F:30:TYR:O	8:F:206:IOD:I	3.08	0.41		
1:D:343:PRO:HA	1:D:373:TYR:O	2.21	0.41		
2:B:261:CYS:HB2	2:B:277:TRP:CH2	2.55	0.41		
2:B:382:GLU:HG2	2:B:424:SER:OG	2.21	0.41		



Atom-1	Atom-2	${f Interatomic}\ {f distance}\ ({ m \AA})$	Clash overlap (Å)			
2:B:318:GLU:HG3	2:B:335:THR:HG22	2.02	0.41			
1:D:378:ALA:HB3	1:D:428:MET:HB2	2.02	0.40			

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	\mathbf{ntiles}
1	А	210/230~(91%)	202~(96%)	8 (4%)	0	100	100
1	D	210/230~(91%)	202~(96%)	8 (4%)	0	100	100
2	В	208/230~(90%)	201~(97%)	6 (3%)	1 (0%)	29	58
2	Е	208/230~(90%)	201 (97%)	6 (3%)	1 (0%)	29	58
3	С	165/197~(84%)	162~(98%)	3 (2%)	0	100	100
3	F	166/197~(84%)	162 (98%)	4 (2%)	0	100	100
All	All	1167/1314 (89%)	1130 (97%)	35 (3%)	2(0%)	47	76

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	В	271	PRO
2	Е	271	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	190/214~(89%)	188~(99%)	2(1%)	73	90
1	D	187/214~(87%)	186 (100%)	1 (0%)	88	95
2	В	188/215~(87%)	184 (98%)	4 (2%)	53	81
2	Ε	181/215~(84%)	181~(100%)	0	100	100
3	С	150/178~(84%)	149~(99%)	1 (1%)	84	94
3	F	148/178~(83%)	143~(97%)	5(3%)	37	68
All	All	1044/1214~(86%)	1031 (99%)	13 (1%)	71	90

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	361	ASN
1	А	376	ASP
2	В	325	ASN
2	В	335	THR
2	В	361	ASN
2	В	363	VAL
3	С	38	GLN
1	D	255	ARG
3	F	43	GLU
3	F	45	LEU
3	F	46	ILE
3	F	94	ARG
3	F	159	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	315	ASN
1	А	362	GLN
2	В	325	ASN
2	В	418	GLN
2	В	429	HIS
3	С	69	GLN
3	С	71	GLN
3	С	80	GLN
3	С	171	GLN
1	D	315	ASN
1	D	347	GLN
2	Е	315	ASN



Continued from previous page...

Mol	Chain	Res	Type
2	Е	347	GLN
2	Е	418	GLN
2	Е	429	HIS
3	F	12	GLN
3	F	69	GLN
3	F	108	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

44 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	True	Chain	Dec	T in le	Bo	ond leng	ths	B	ond ang	les
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
4	NAG	G	1	1,4	14,14,15	0.47	0	$17,\!19,\!21$	1.29	3 (17%)
4	NAG	G	2	4	14,14,15	0.58	0	17,19,21	0.81	0
4	BMA	G	3	4	11,11,12	0.45	0	$15,\!15,\!17$	0.76	0
4	MAN	G	4	4	11,11,12	0.56	0	$15,\!15,\!17$	0.95	1(6%)
4	NAG	G	5	4	14,14,15	0.53	0	$17,\!19,\!21$	0.83	1(5%)
4	GAL	G	6	4	11,11,12	0.59	0	$15,\!15,\!17$	1.22	1(6%)
4	MAN	G	7	4	11,11,12	0.57	0	$15,\!15,\!17$	0.90	0
4	NAG	G	8	4	14,14,15	0.46	0	$17,\!19,\!21$	0.76	0
4	FUL	G	9	4	10,10,11	0.64	0	$14,\!14,\!16$	1.48	3 (21%)
4	NAG	Н	1	2,4	14,14,15	0.59	0	17,19,21	0.95	1 (5%)
4	NAG	Н	2	4	14,14,15	0.51	0	17,19,21	0.89	0



Mal	Tune	Chain	Dog	Tink	Bo	ond leng	ths	Bond angles		
	туре	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	BMA	Н	3	4	11,11,12	0.56	0	$15,\!15,\!17$	1.19	2 (13%)
4	MAN	Н	4	4	11,11,12	0.55	0	$15,\!15,\!17$	0.84	1 (6%)
4	NAG	Н	5	4	14,14,15	0.59	0	17,19,21	1.22	1 (5%)
4	GAL	Н	6	4	11,11,12	0.59	0	$15,\!15,\!17$	0.79	0
4	MAN	Н	7	4	11,11,12	0.50	0	$15,\!15,\!17$	1.67	3 (20%)
4	NAG	Н	8	4	14,14,15	0.60	0	17,19,21	1.07	1 (5%)
4	FUL	Н	9	4	10, 10, 11	0.43	0	14, 14, 16	0.89	1 (7%)
5	NAG	Ι	1	3,5	14,14,15	0.48	0	17,19,21	1.89	2 (11%)
5	NAG	Ι	2	5	14,14,15	0.49	0	17,19,21	0.79	0
5	BMA	Ι	3	5	11,11,12	0.44	0	$15,\!15,\!17$	0.77	0
5	MAN	Ι	4	5	11, 11, 12	0.60	0	$15,\!15,\!17$	0.97	0
5	MAN	Ι	5	5	11,11,12	0.59	0	$15,\!15,\!17$	1.18	2(13%)
6	NAG	J	1	3,6	14,14,15	0.48	0	$17,\!19,\!21$	0.94	1(5%)
6	NAG	J	2	6	14,14,15	0.45	0	$17,\!19,\!21$	0.94	1 (5%)
7	NAG	K	1	1,7	14,14,15	0.53	0	17,19,21	1.12	2 (11%)
7	NAG	K	2	7	14,14,15	0.54	0	17,19,21	0.70	0
7	BMA	K	3	7	11,11,12	0.43	0	$15,\!15,\!17$	1.14	2 (13%)
7	MAN	K	4	7	11,11,12	0.52	0	$15,\!15,\!17$	0.85	1(6%)
7	NAG	K	5	7	14,14,15	0.49	0	17,19,21	0.73	0
7	MAN	K	6	7	11,11,12	0.60	0	$15,\!15,\!17$	0.87	1 (6%)
7	NAG	K	7	7	14,14,15	0.50	0	17,19,21	0.82	1(5%)
7	FUL	K	8	7	10,10,11	0.45	0	14,14,16	0.59	0
4	NAG	L	1	2,4	14,14,15	0.44	0	17,19,21	1.35	1 (5%)
4	NAG	L	2	4	14,14,15	0.47	0	17,19,21	0.83	0
4	BMA	L	3	4	11, 11, 12	0.46	0	$15,\!15,\!17$	0.79	0
4	MAN	L	4	4	11, 11, 12	0.57	0	$15,\!15,\!17$	0.92	1(6%)
4	NAG	L	5	4	14, 14, 15	0.54	0	17,19,21	0.83	0
4	GAL	L	6	4	11,11,12	0.64	0	$15,\!15,\!17$	0.73	0
4	MAN	L	7	4	11,11,12	0.48	0	$15,\!15,\!17$	1.19	1 (6%)
4	NAG	L	8	4	14,14,15	0.58	0	$17,\!19,\!21$	1.09	1(5%)
4	FUL	L	9	4	$10,\!10,\!11$	0.49	0	$14,\!14,\!16$	0.64	0
6	NAG	М	1	3,6	14,14,15	0.51	0	17,19,21	0.95	1 (5%)
6	NAG	M	2	6	14,14,15	0.48	0	17,19,21	0.93	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	0/6/23/26	0/1/1/1
4	BMA	G	3	4	-	0/2/19/22	0/1/1/1
4	MAN	G	4	4	-	0/2/19/22	0/1/1/1
4	NAG	G	5	4	-	2/6/23/26	0/1/1/1
4	GAL	G	6	4	-	2/2/19/22	0/1/1/1
4	MAN	G	7	4	-	0/2/19/22	0/1/1/1
4	NAG	G	8	4	-	0/6/23/26	0/1/1/1
4	FUL	G	9	4	-	_	0/1/1/1
4	NAG	Н	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	Н	2	4	-	0/6/23/26	0/1/1/1
4	BMA	Н	3	4	-	1/2/19/22	0/1/1/1
4	MAN	Н	4	4	_	0/2/19/22	0/1/1/1
4	NAG	Н	5	4	-	0/6/23/26	0/1/1/1
4	GAL	H	6	4	-	0/2/19/22	0/1/1/1
4	MAN	Н	7	4	-	0/2/19/22	0/1/1/1
4	NAG	Н	8	4	-	0/6/23/26	0/1/1/1
4	FUL	Н	9	4	-	-	0/1/1/1
5	NAG	I	1	3,5	-	3/6/23/26	0/1/1/1
5	NAG	Ι	2	5	_	0/6/23/26	0/1/1/1
5	BMA	I	3	5	-	1/2/19/22	0/1/1/1
5	MAN	Ι	4	5	-	2/2/19/22	0/1/1/1
5	MAN	Ι	5	5	-	2/2/19/22	0/1/1/1
6	NAG	J	1	$3,\!6$	-	1/6/23/26	0/1/1/1
6	NAG	J	2	6	-	0/6/23/26	0/1/1/1
7	NAG	Κ	1	1,7	-	2/6/23/26	0/1/1/1
7	NAG	K	2	7	-	0/6/23/26	0/1/1/1
7	BMA	K	3	7	-	0/2/19/22	0/1/1/1
7	MAN	Κ	4	7	-	0/2/19/22	0/1/1/1
7	NAG	Κ	5	7	-	0/6/23/26	0/1/1/1
7	MAN	K	6	7	-	0/2/19/22	0/1/1/1
7	NAG	K	7	7	-	0/6/23/26	0/1/1/1
7	FUL	K	8	7	-	-	0/1/1/1
4	NAG	L	1	2,4	-	2/6/23/26	0/1/1/1
4	NAG	L	2	4	-	0/6/23/26	0/1/1/1
4	BMA	L	3	4	-	0/2/19/22	0/1/1/1
4	MAN	L	4	4	-	0/2/19/22	0/1/1/1
4	NAG	L	5	4	-	2/6/23/26	0/1/1/1
4	GAL	L	6	4	-	2/2/19/22	0/1/1/1
4	MAN	L	7	4	-	0/2/19/22	0/1/1/1
4	NAG	L	8	4	-	1/6/23/26	0/1/1/1



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	FUL	L	9	4	-	-	0/1/1/1
6	NAG	М	1	3,6	-	0/6/23/26	0/1/1/1
6	NAG	М	2	6	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	Ι	1	NAG	C1-O5-C5	5.68	119.89	112.19
4	L	1	NAG	C1-O5-C5	4.09	117.73	112.19
4	Н	5	NAG	C4-C3-C2	3.67	116.40	111.02
4	G	6	GAL	C1-O5-C5	3.50	116.94	112.19
4	G	9	FUL	C3-C4-C5	3.48	115.19	109.77
5	Ι	1	NAG	C2-N2-C7	3.42	127.77	122.90
4	Н	7	MAN	C1-O5-C5	3.28	116.64	112.19
7	К	3	BMA	C1-O5-C5	3.27	116.62	112.19
4	L	7	MAN	C1-O5-C5	3.25	116.59	112.19
4	G	1	NAG	O5-C5-C6	3.23	112.26	107.20
5	Ι	5	MAN	C1-O5-C5	3.13	116.43	112.19
4	Н	7	MAN	O2-C2-C1	2.88	115.05	109.15
4	Н	7	MAN	O2-C2-C3	-2.88	104.36	110.14
4	G	4	MAN	C1-O5-C5	2.84	116.04	112.19
4	Н	3	BMA	C1-C2-C3	2.78	113.08	109.67
4	Н	8	NAG	C4-C3-C2	2.76	115.06	111.02
4	G	1	NAG	C1-O5-C5	2.73	115.89	112.19
4	L	4	MAN	C1-O5-C5	2.68	115.83	112.19
4	Н	3	BMA	C1-O5-C5	2.66	115.80	112.19
6	J	2	NAG	C1-O5-C5	2.59	115.71	112.19
6	М	2	NAG	C1-O5-C5	2.57	115.67	112.19
7	Κ	3	BMA	C1-C2-C3	2.56	112.82	109.67
6	J	1	NAG	C1-O5-C5	2.56	115.66	112.19
4	G	5	NAG	O5-C5-C6	2.53	111.17	107.20
7	K	4	MAN	C1-O5-C5	2.44	115.50	112.19
4	L	8	NAG	C3-C4-C5	2.43	114.57	110.24
4	G	9	FUL	O5-C5-C4	2.40	113.82	109.52
4	Н	1	NAG	C1-O5-C5	2.29	115.29	112.19
7	Κ	6	MAN	C1-O5-C5	2.25	115.24	112.19
7	Κ	7	NAG	O5-C5-C6	2.25	110.73	107.20
4	Н	4	MAN	C1-O5-C5	2.23	115.21	112.19
5	Ι	5	MAN	C3-C4-C5	2.20	114.16	110.24
6	М	1	NAG	C1-O5-C5	2.19	115.17	112.19
4	Н	9	FUL	C1-C2-C3	2.16	112.32	109.67



Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
4	G	9	FUL	C1-O5-C5	2.15	117.66	112.78
7	Κ	1	NAG	O5-C5-C6	2.13	110.54	107.20
7	Κ	1	NAG	O5-C1-C2	-2.04	108.07	111.29
4	G	1	NAG	C6-C5-C4	-2.02	108.27	113.00

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Н	1	NAG	O5-C5-C6-O6
4	L	6	GAL	O5-C5-C6-O6
4	G	1	NAG	O5-C5-C6-O6
5	Ι	4	MAN	O5-C5-C6-O6
4	G	6	GAL	O5-C5-C6-O6
4	L	6	GAL	C4-C5-C6-O6
4	Н	1	NAG	C4-C5-C6-O6
5	Ι	1	NAG	C4-C5-C6-O6
4	G	1	NAG	C4-C5-C6-O6
4	L	1	NAG	O5-C5-C6-O6
5	Ι	4	MAN	C4-C5-C6-O6
4	G	5	NAG	O5-C5-C6-O6
4	G	5	NAG	C4-C5-C6-O6
5	Ι	3	BMA	O5-C5-C6-O6
7	K	1	NAG	O5-C5-C6-O6
4	G	6	GAL	C4-C5-C6-O6
4	L	5	NAG	C4-C5-C6-O6
5	Ι	1	NAG	O5-C5-C6-O6
4	L	5	NAG	O5-C5-C6-O6
5	Ι	5	MAN	C4-C5-C6-O6
4	L	1	NAG	C4-C5-C6-O6
6	J	1	NAG	C4-C5-C6-O6
6	М	2	NAG	O5-C5-C6-O6
4	L	8	NAG	O5-C5-C6-O6
7	Κ	1	NAG	C4-C5-C6-O6
5	Ι	5	MAN	O5-C5-C6-O6
5	I	1	NAG	C3-C2-N2-C7
4	Н	3	BMA	O5-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 3 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Н	9	FUL	2	0
4	G	5	NAG	1	0
4	G	6	GAL	1	0
4	Н	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



























5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 16 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Dog	Link	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
	rybe	Ullaili	II Res		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	NAG	F	201	3	14, 14, 15	0.50	0	17,19,21	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	NAG	F	201	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	F	201	NAG	C4-C5-C6-O6
9	F	201	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2			$OWAB(Å^2)$	$Q{<}0.9$
1	А	212/230~(92%)	0.36	5(2%)	59	54	38, 52, 68, 79	0
1	D	212/230~(92%)	0.26	3 (1%)	75	73	37, 57, 84, 98	0
2	В	210/230~(91%)	0.36	5(2%)	59	54	31, 48, 77, 94	0
2	Ε	210/230~(91%)	0.59	12 (5%)	23	18	44, 70, 111, 121	0
3	С	169/197~(85%)	0.22	3~(1%)	68	65	30, 44, 73, 98	0
3	F	170/197~(86%)	0.29	4 (2%)	59	54	30, 45, 68, 94	0
All	All	1183/1314~(90%)	0.35	32(2%)	54	49	30, 53, 85, 121	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	С	47	SER	5.8
1	D	420	GLY	3.9
1	D	443	LEU	3.9
2	Ε	443	LEU	3.7
3	С	48	SER	3.5
3	F	171	GLN	3.2
2	Е	296	TYR	3.0
2	Ε	444	SER	3.0
3	F	143	LEU	3.0
2	Ε	415	SER	2.7
2	В	444	SER	2.6
1	А	253	ILE	2.5
2	Ε	387	PRO	2.5
3	С	171	GLN	2.5
2	Ε	411	THR	2.4
2	В	278	TYR	2.4
3	F	155	VAL	2.4
3	F	172	GLY	2.3
1	А	443	LEU	2.3



		1	1 0	
Mol	Chain	\mathbf{Res}	Type	RSRZ
1	А	309	LEU	2.2
2	В	277	TRP	2.2
2	Е	252	MET	2.2
2	Е	420	GLY	2.2
2	Е	358	LEU	2.2
1	А	420	GLY	2.2
1	А	431	ALA	2.2
2	Е	359	THR	2.2
2	Е	391	TYR	2.1
2	Е	303	VAL	2.1
2	В	305	VAL	2.1
2	В	284	VAL	2.1
1	D	444	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
4	NAG	L	8	14/15	0.66	0.34	$96,\!100,\!107,\!108$	0
4	GAL	G	6	11/12	0.70	0.39	77,83,85,86	0
5	MAN	Ι	5	11/12	0.71	0.23	84,86,90,93	0
4	NAG	Н	8	14/15	0.75	0.25	$88,\!91,\!93,\!93$	0
4	FUL	L	9	10/11	0.80	0.24	75,78,79,79	0
7	NAG	K	5	14/15	0.83	0.23	71,74,75,76	0
4	FUL	Н	9	10/11	0.83	0.23	$63,\!65,\!66,\!67$	0
6	NAG	М	2	14/15	0.84	0.30	$65,\!69,\!71,\!71$	0
6	NAG	J	2	14/15	0.86	0.19	$65,\!68,\!71,\!72$	0
5	MAN	Ι	4	11/12	0.86	0.26	82,84,86,87	0
4	FUL	G	9	10/11	0.87	0.24	$46,\!47,\!48,\!51$	0
4	MAN	Н	7	11/12	0.88	0.17	$68,\!72,\!75,\!83$	0
7	FUL	K	8	10/11	0.88	0.19	59,60,61,62	0
4	NAG	L	1	14/15	0.89	0.18	$61,\!64,\!68,\!72$	0
4	GAL	L	6	11/12	0.89	0.21	64,66,67,67	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
4	NAG	L	5	14/15	0.91	0.15	$56,\!59,\!61,\!62$	0
4	GAL	Н	6	11/12	0.91	0.22	73,74,76,78	0
5	BMA	Ι	3	11/12	0.91	0.13	71,74,78,80	0
7	NAG	K	7	14/15	0.91	0.20	$54,\!56,\!58,\!58$	0
4	NAG	G	8	14/15	0.91	0.20	$52,\!53,\!56,\!59$	0
4	MAN	L	7	11/12	0.91	0.16	76,80,82,89	0
4	NAG	Н	5	14/15	0.92	0.18	$67,\!68,\!70,\!72$	0
7	MAN	K	4	11/12	0.92	0.14	$62,\!64,\!65,\!68$	0
7	NAG	K	1	14/15	0.93	0.18	$49,\!51,\!54,\!59$	0
4	NAG	Н	2	14/15	0.93	0.17	$50,\!54,\!56,\!57$	0
4	MAN	G	7	11/12	0.93	0.17	$51,\!53,\!55,\!55$	0
4	BMA	Н	3	11/12	0.93	0.17	$57,\!59,\!61,\!67$	0
5	NAG	Ι	1	14/15	0.94	0.20	$47,\!49,\!53,\!54$	0
4	MAN	G	4	11/12	0.94	0.18	$53,\!54,\!55,\!56$	0
6	NAG	М	1	14/15	0.94	0.17	$46,\!49,\!53,\!58$	0
4	NAG	G	5	14/15	0.94	0.21	$55,\!58,\!62,\!68$	0
4	BMA	L	3	11/12	0.94	0.15	$60,\!63,\!65,\!71$	0
4	MAN	L	4	11/12	0.94	0.12	$55,\!58,\!59,\!60$	0
4	MAN	Н	4	11/12	0.95	0.16	$62,\!63,\!64,\!65$	0
4	NAG	G	2	14/15	0.96	0.18	$42,\!43,\!44,\!44$	0
5	NAG	Ι	2	14/15	0.96	0.14	$51,\!57,\!61,\!67$	0
4	NAG	G	1	14/15	0.96	0.12	$43,\!44,\!45,\!46$	0
7	NAG	K	2	14/15	0.96	0.19	$47,\!50,\!53,\!53$	0
4	NAG	Н	1	14/15	0.96	0.17	$45,\!50,\!56,\!61$	0
6	NAG	J	1	14/15	0.96	0.20	$57,\!60,\!63,\!6\overline{5}$	0
4	BMA	G	3	11/12	0.97	0.18	$44,\!46,\!50,\!51$	0
7	BMA	K	3	11/12	0.97	0.14	49,52,54,58	0
7	MAN	K	6	11/12	0.97	0.15	$47,\!48,\!49,\!52$	0
4	NAG	L	2	14/15	0.97	0.14	$55,\!60,\!61,\!63$	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





























6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	NAG	F	201	14/15	0.84	0.27	$64,\!74,\!77,\!78$	0
8	IOD	А	1010	1/1	0.89	0.14	71,71,71,71	1
8	IOD	D	1009	1/1	0.91	0.12	86,86,86,86	1
8	IOD	А	1011	1/1	0.91	0.10	75,75,75,75	1
8	IOD	В	1012	1/1	0.93	0.11	80,80,80,80	1
8	IOD	F	207	1/1	0.95	0.10	81,81,81,81	1
8	IOD	Е	1013	1/1	0.96	0.07	99,99,99,99	0
8	IOD	F	206	1/1	0.97	0.09	$58,\!58,\!58,\!58$	1



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	$Q{<}0.9$
8	IOD	Ε	1011	1/1	0.97	0.09	$85,\!85,\!85,\!85$	0
8	IOD	С	209	1/1	0.98	0.09	91,91,91,91	0
8	IOD	Ε	1012	1/1	0.98	0.06	$93,\!93,\!93,\!93$	0
8	IOD	F	205	1/1	0.99	0.09	$68,\!68,\!68,\!68$	0
8	IOD	В	1010	1/1	0.99	0.13	$49,\!49,\!49,\!49$	0
8	IOD	В	1011	1/1	0.99	0.07	83,83,83,83	0
8	IOD	Е	1010	1/1	0.99	0.08	$102,\!102,\!102,\!102$	0
8	IOD	С	208	1/1	1.00	0.16	$36,\!36,\!36,\!36$	0
8	IOD	F	204	1/1	1.00	0.17	$43,\!43,\!43,\!43$	0

6.5 Other polymers (i)

There are no such residues in this entry.

