

Jan 17, 2023 - 02:11 PM JST

PDB ID : 7WNZ EMDB ID : EMD-32636 Title : CryoEM structure of human alpha-synuclein A53T fibril : Wu, K.-P.; Huang, J.Y.-C. Authors Deposited on 2022-01-20 : 3.40 Å(reported) Resolution : Based on initial model 6CU7:

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


:	0.0.1. dev 43
:	4.02b-467
:	20191225.v01 (using entries in the PDB archive December 25th 2019)
:	1.9.9
:	Engh & Huber (2001)
:	Parkinson et al. (1996)
:	2.31.3
	: : : :

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	EM structures
	$(\# { m Entries})$	$(\# { m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
- 1		05	11%	
1	A	65	89%	11%
1	р	CT.	12%	
1	В	65	88%	12%
			11%	
1	С	65	88%	12%
			12%	
1	D	65	88%	12%
			12%	
1	Ε	65	91%	9%
			11%	
1	F	65	89%	11%
			12%	
1	G	65	88%	12%
			12%	
1	Н	65	88%	12%

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	Ι	65	88%	12%
1	J	65	89%	11%



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 9300 atoms, of which 4780 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			AltConf	Trace
1	C	65	Total	С	Η	Ν	0	0	0
1	G	65	930	283	478	78	91	0	0
1	А	65	Total	С	Η	Ν	0	0	0
1	A	05	930	283	478	78	91	0	0
1	В	65	Total	С	Η	Ν	0	0	0
	D	05	930	283	478	78	91	0	0
1	С	65	Total	С	Η	Ν	0	0	0
	C	65	930	283	478	78	91	0	0
1	D	65	Total	С	Η	Ν	0	0	0
	D	65	930	283	478	78	91		0
1	Е	65	Total	С	Η	Ν	0	0	0
	E	05	930	283	478	78	91	0	
1	F	65	Total	С	Н	Ν	0	0	0
	Г	05	930	283	478	78	91	0	0
1	Н	65	Total	С	Η	Ν	0	0	0
	п	05	930	283	478	78	91	0	0
1	Ι	65	Total	С	Η	Ν	0	0	0
		65	930	283	478	78	91		0
1	J	65	Total	С	Η	Ν	0	0	0
	J	00	930	283	478	78	91	U	0

• Molecule 1 is a protein called Alpha-synuclein.

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	53	THR	ALA	variant	UNP P37840
А	53	THR	ALA	variant	UNP P37840
В	53	THR	ALA	variant	UNP P37840
С	53	THR	ALA	variant	UNP P37840
D	53	THR	ALA	variant	UNP P37840
E	53	THR	ALA	variant	UNP P37840
F	53	THR	ALA	variant	UNP P37840
Н	53	THR	ALA	variant	UNP P37840
Ι	53	THR	ALA	variant	UNP P37840
J	53	THR	ALA	variant	UNP P37840




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-synuclein









4 Experimental information (i)

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=179.304°, rise=2.37 Å, ax-	Depositor
	ial sym= $C1$	
Number of segments used	46892	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	1.05	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.057	Depositor
Minimum map value	-0.033	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	232.4, 232.4, 232.4	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.83, 0.83, 0.83	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Ullaill	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.29	0/454	0.49	0/614
1	В	0.29	0/454	0.48	0/614
1	С	0.29	0/454	0.48	0/614
1	D	0.29	0/454	0.49	0/614
1	Ε	0.28	0/454	0.49	0/614
1	F	0.29	0/454	0.48	0/614
1	G	0.29	0/454	0.48	0/614
1	Н	0.29	0/454	0.48	0/614
1	Ι	0.29	0/454	0.48	0/614
1	J	0.29	0/454	0.49	0/614
All	All	0.29	0/4540	0.48	0/6140

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	452	478	475	7	0
1	В	452	478	475	10	0
1	С	452	478	475	11	0
1	D	452	478	475	11	0
1	Е	452	478	475	6	0
1	F	452	478	475	8	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	452	478	475	12	0
1	Н	452	478	475	12	0
1	Ι	452	478	475	12	0
1	J	452	478	475	8	0
All	All	4520	4780	4750	59	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:E:50:HIS:NE2	1:J:57:GLU:OE2	2.18	0.77
1:G:57:GLU:OE2	1:D:50:HIS:NE2	2.24	0.68
1:D:57:GLU:OE2	1:J:50:HIS:NE2	2.20	0.68
1:A:50:HIS:NE2	1:F:57:GLU:OE2	2.23	0.66
1:H:52:VAL:HG22	1:I:52:VAL:HG12	1.84	0.59
1:G:80:LYS:HD2	1:H:46:GLU:HB3	1.84	0.58
1:C:52:VAL:HG12	1:D:52:VAL:HG22	1.84	0.58
1:A:52:VAL:O	1:A:52:VAL:HG23	2.04	0.58
1:E:52:VAL:O	1:E:52:VAL:HG23	2.04	0.57
1:G:46:GLU:HB3	1:J:80:LYS:HD2	1.85	0.57
1:G:52:VAL:HG23	1:G:52:VAL:O	2.04	0.57
1:J:52:VAL:O	1:J:52:VAL:HG23	2.04	0.57
1:D:52:VAL:HG23	1:D:52:VAL:O	2.04	0.57
1:H:52:VAL:HG23	1:H:52:VAL:O	2.04	0.57
1:I:52:VAL:O	1:I:52:VAL:HG23	2.04	0.57
1:F:52:VAL:HG12	1:I:52:VAL:HG22	1.87	0.57
1:F:52:VAL:O	1:F:52:VAL:HG23	2.04	0.57
1:C:52:VAL:HG23	1:C:52:VAL:O	2.04	0.56
1:B:52:VAL:O	1:B:52:VAL:HG23	2.04	0.56
1:C:50:HIS:NE2	1:H:57:GLU:OE2	2.29	0.56
1:A:52:VAL:HG12	1:B:52:VAL:HG22	1.87	0.56
1:G:52:VAL:HG12	1:J:52:VAL:HG22	1.88	0.55
1:G:50:HIS:NE2	1:C:57:GLU:OE2	2.29	0.55
1:G:52:VAL:HG22	1:H:52:VAL:HG12	1.88	0.55
1:B:46:GLU:HB3	1:C:80:LYS:HD2	1.88	0.55
1:D:52:VAL:HG12	1:E:52:VAL:HG22	1.88	0.55
1:B:52:VAL:HG12	1:C:52:VAL:HG22	1.87	0.55
1:A:57:GLU:OE2	1:I:50:HIS:NE2	2.31	0.54
1:B:57:GLU:OE2	1:H:50:HIS:NE2	2.31	0.54

Continued on next page...



Continued from pre		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:D:46:GLU:HB3	1:E:80:LYS:HD2	1.90	0.53
1:B:50:HIS:NE2	1:I:57:GLU:OE2	2.32	0.53
1:C:46:GLU:HB3	1:D:80:LYS:HD2	1.90	0.53
1:H:80:LYS:HD2	1:I:46:GLU:HB3	1.94	0.49
1:C:51:GLY:O	1:D:51:GLY:HA2	2.14	0.48
1:G:51:GLY:HA2	1:H:51:GLY:O	2.14	0.47
1:B:51:GLY:O	1:C:51:GLY:HA2	2.15	0.47
1:G:51:GLY:O	1:J:51:GLY:HA2	2.15	0.47
1:A:46:GLU:HB3	1:B:80:LYS:HD2	1.97	0.46
1:B:91:ALA:HB1	1:B:94:PHE:HB2	1.98	0.46
1:A:91:ALA:HB1	1:A:94:PHE:HB2	1.98	0.46
1:C:91:ALA:HB1	1:C:94:PHE:HB2	1.98	0.46
1:F:91:ALA:HB1	1:F:94:PHE:HB2	1.98	0.46
1:H:91:ALA:HB1	1:H:94:PHE:HB2	1.98	0.46
1:I:91:ALA:HB1	1:I:94:PHE:HB2	1.98	0.46
1:G:91:ALA:HB1	1:G:94:PHE:HB2	1.98	0.46
1:D:91:ALA:HB1	1:D:94:PHE:HB2	1.98	0.46
1:E:91:ALA:HB1	1:E:94:PHE:HB2	1.98	0.45
1:J:91:ALA:HB1	1:J:94:PHE:HB2	1.98	0.45
1:D:51:GLY:O	1:E:51:GLY:HA2	2.18	0.44
1:H:51:GLY:HA2	1:I:51:GLY:O	2.17	0.43
1:A:51:GLY:O	1:B:51:GLY:HA2	2.18	0.43
1:F:46:GLU:HB3	1:I:80:LYS:HD2	2.00	0.42
1:F:51:GLY:O	1:I:51:GLY:HA2	2.19	0.42
1:G:80:LYS:NZ	1:H:46:GLU:CD	2.74	0.42
1:C:52:VAL:CG1	1:D:52:VAL:HG22	2.49	0.41
1:F:80:LYS:O	1:I:80:LYS:HB2	2.21	0.41
1:G:46:GLU:CD	1:J:80:LYS:NZ	2.74	0.41
1:H:52:VAL:HG22	1:I:52:VAL:CG1	2.49	0.41
1:F:52:VAL:O	1:F:52:VAL:CG2	2.69	0.41

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	В	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	\mathbf{C}	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	D	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	Ε	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	\mathbf{F}	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	G	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	Н	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	Ι	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
1	J	63/65~(97%)	59 (94%)	4~(6%)	0	100 100
All	All	630/650~(97%)	590 (94%)	40 (6%)	0	100 100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	47/47~(100%)	47~(100%)	0	100 100
1	В	47/47~(100%)	47 (100%)	0	100 100
1	С	47/47~(100%)	47 (100%)	0	100 100
1	D	47/47~(100%)	47 (100%)	0	100 100
1	Е	47/47~(100%)	47 (100%)	0	100 100
1	F	47/47~(100%)	47 (100%)	0	100 100
1	G	47/47~(100%)	47 (100%)	0	100 100
1	Н	47/47~(100%)	47 (100%)	0	100 100
1	Ι	47/47~(100%)	47 (100%)	0	100 100

Continued on next page...



Mol	Chain	Analysed	Rotameric	Outliers	Perce	entiles
1	J	47/47~(100%)	47 (100%)	0	100	100
All	All	470/470 (100%)	470 (100%)	0	100	100

Continued from previous page...

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-32636. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

Orthogonal projections (i) 6.1


6.1.1Primary map



The images above show the map projected in three orthogonal directions.

6.2Central slices (i)

6.2.1Primary map




X Index: 140



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 30 nm^3 ; this corresponds to an approximate mass of 27 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




*Reported resolution corresponds to spatial frequency of 0.294 $\mathrm{\AA^{-1}}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.294 $Å^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
Resolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	3.40	-	-	
Author-provided FSC curve	3.37	3.75	3.40	
Unmasked-calculated*	-	-	_	


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-32636 and PDB model 7WNZ. Per-residue inclusion information can be found in section 3 on page 5.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).



9.4 Atom inclusion (i)



At the recommended contour level, 63% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.



Map-model fit summary (i) 9.5

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score	1.0
All	0.6308	0.4680	1.0
А	0.6430	0.4700	
В	0.6341	0.4750	
С	0.6364	0.4710	
D	0.6319	0.4730	
E	0.6297	0.4640	
F	0.6364	0.4620	
G	0.6430	0.4670	
Н	0.6341	0.4690	0.0
I	0.6497	0.4710	<0.0
J	0.6231	0.4630	

