



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2024 – 12:10 AM JST

PDB ID : 8WQ6
Title : Durio zibethinus trypsin inhibitor DzTI-5 (lattice translocation disorder)
Authors : Deentanya, P.; Wangkanont, K.
Deposited on : 2023-10-11
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1935 (1.56-1.56)
Clashscore	180529	2073 (1.56-1.56)
Ramachandran outliers	177936	2037 (1.56-1.56)
Sidechain outliers	177891	2034 (1.56-1.56)
RSRZ outliers	164620	1935 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	 11% 94%
1	B	194	 14% 83% 11% 5%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 3456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 21 kDa seed protein-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	191	Total 1497	C 946	N 255	O 289	S 7	0	0	0
1	B	184	Total 1442	C 911	N 245	O 279	S 7	0	0	0

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	Zn	0	0
			10	10		
3	B	8	Total	Zn	0	0
			8	8		

- Molecule 4 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			16	10	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	240	Total	O	0	0
			240	240		
5	B	227	Total	O	0	0
			227	227		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 21 kDa seed protein-like



- Molecule 1: 21 kDa seed protein-like



ASN

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.93Å 64.10Å 128.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.70 – 1.55 29.70 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.70-1.55) 99.8 (29.70-1.55)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 1.55Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.221 , 0.246 0.221 , 0.245	Depositor DCC
R_{free} test set	3225 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	17.1	Xtrriage
Anisotropy	0.451	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3456	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.59 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0957e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, ZN, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1531	0.61	0/2080
1	B	0.38	0/1476	0.65	0/2005
All	All	0.36	0/3007	0.63	0/4085

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1497	0	1447	6	0
1	B	1442	0	1385	17	0
2	A	8	0	6	0	0
2	B	8	0	6	0	0
3	A	10	0	0	0	0
3	B	8	0	0	0	0
4	B	16	0	22	2	0
5	A	240	0	0	2	0
5	B	227	0	0	5	0
All	All	3456	0	2866	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:THR:HG22	5:A:514:HOH:O	1.69	0.91
1:B:93:THR:HG22	5:B:521:HOH:O	1.72	0.88
1:B:63:ARG:HD2	1:B:68:LYS:HB2	1.62	0.81
1:B:161:THR:HG23	1:B:162:LEU:H	1.50	0.76
1:A:63:ARG:HD3	1:A:115:LEU:HD13	1.80	0.64
1:B:207:LYS:HE2	1:B:210:SER:OG	2.03	0.58
1:B:112:ARG:HH11	1:B:112:ARG:HG3	1.67	0.58
1:B:94:LYS:NZ	5:B:401:HOH:O	2.39	0.55
1:B:27:LYS:HE3	1:B:28:ASN:H	1.73	0.53
1:B:114:LYS:HG3	1:B:116:CYS:H	1.74	0.53
1:B:68:LYS:HZ2	1:B:115:LEU:HB3	1.75	0.52
1:A:114:LYS:HE2	1:A:116:CYS:O	2.09	0.52
1:B:110:PRO:O	5:B:402:HOH:O	2.20	0.47
1:A:27:LYS:NZ	1:A:28:ASN:H	2.13	0.46
1:B:92:ASP:OD2	5:B:401:HOH:O	2.20	0.46
1:B:38:GLU:HG3	1:B:97:ILE:HD12	1.98	0.46
4:B:311:1PE:H241	4:B:311:1PE:H251	1.64	0.45
1:B:68:LYS:HD2	1:B:69:CYS:SG	2.57	0.44
1:B:82:TYR:OH	4:B:311:1PE:H141	2.17	0.44
1:A:96:ASP:OD1	5:A:401:HOH:O	2.21	0.44
1:B:112:ARG:HH11	1:B:112:ARG:CG	2.31	0.43
1:B:27:LYS:HE2	1:B:102:ASN:HD21	1.85	0.42
1:A:133:LYS:NZ	5:B:406:HOH:O	2.43	0.41
1:B:158:LYS:HD3	1:B:158:LYS:HA	1.89	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/194 (97%)	182 (96%)	7 (4%)	0	100	100
1	B	182/194 (94%)	174 (96%)	8 (4%)	0	100	100
All	All	371/388 (96%)	356 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/167 (99%)	164 (99%)	1 (1%)	84	72
1	B	158/167 (95%)	156 (99%)	2 (1%)	65	41
All	All	323/334 (97%)	320 (99%)	3 (1%)	75	58

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LYS
1	B	68	LYS
1	B	209	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 18 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	A	302	-	3,3,3	1.21	0	3,3,3	1.30	0
4	1PE	B	311	-	15,15,15	0.09	0	14,14,14	0.25	0
2	ACT	A	301	-	3,3,3	1.31	0	3,3,3	1.31	0
2	ACT	B	302	-	3,3,3	1.23	0	3,3,3	1.36	0
2	ACT	B	301	-	3,3,3	1.22	0	3,3,3	1.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	1PE	B	311	-	-	7/13/13/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	311	1PE	OH5-C14-C24-OH4
4	B	311	1PE	C24-C14-OH5-C25
4	B	311	1PE	OH6-C15-C25-OH5
4	B	311	1PE	C14-C24-OH4-C13
4	B	311	1PE	OH7-C16-C26-OH6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	311	1PE	C25-C15-OH6-C26
4	B	311	1PE	C16-C26-OH6-C15

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	311	1PE	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/194 (98%)	0.60	21 (10%) 12 14	12, 19, 38, 63	0
1	B	184/194 (94%)	0.89	28 (15%) 6 7	12, 19, 36, 47	0
All	All	375/388 (96%)	0.74	49 (13%) 8 10	12, 19, 36, 63	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	113	ASP	5.9
1	A	111	ILE	5.9
1	B	160	GLY	5.8
1	B	210	SER	5.1
1	B	161	THR	4.7
1	B	114	LYS	4.7
1	A	216	VAL	4.1
1	A	112	ARG	3.9
1	B	68	LYS	3.9
1	A	161	THR	3.9
1	A	215	VAL	3.5
1	A	115	LEU	3.4
1	B	67	GLN	3.4
1	B	173	CYS	3.2
1	B	104	LEU	3.2
1	B	66	ASP	3.1
1	A	162	LEU	3.0
1	B	69	CYS	3.0
1	A	199	ASN	2.9
1	A	173	CYS	2.9
1	A	63	ARG	2.9
1	B	111	ILE	2.8
1	A	67	GLN	2.8
1	B	65	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	208	ALA	2.8
1	A	27	LYS	2.8
1	A	114	LYS	2.7
1	B	112	ARG	2.6
1	B	27	LYS	2.6
1	B	209	SER	2.6
1	A	198	GLU	2.5
1	A	211	THR	2.5
1	B	37	ASP	2.5
1	B	162	LEU	2.5
1	A	66	ASP	2.4
1	B	115	LEU	2.4
1	B	174	GLU	2.4
1	B	63	ARG	2.4
1	B	159	SER	2.3
1	B	113	ASP	2.3
1	A	212	ILE	2.3
1	B	110	PRO	2.3
1	B	158	LYS	2.3
1	A	174	GLU	2.2
1	A	110	PRO	2.1
1	B	199	ASN	2.1
1	B	117	LEU	2.1
1	A	217	ASN	2.0
1	B	29	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	1PE	B	311	16/16	0.78	0.20	24,29,36,37	16
3	ZN	A	310	1/1	0.79	0.17	31,31,31,31	1
2	ACT	B	302	4/4	0.81	0.18	27,30,30,30	4
3	ZN	B	310	1/1	0.85	0.12	24,24,24,24	1
3	ZN	A	308	1/1	0.87	0.13	26,26,26,26	1
2	ACT	A	302	4/4	0.88	0.13	23,23,24,28	4
3	ZN	A	307	1/1	0.89	0.10	36,36,36,36	1
3	ZN	B	306	1/1	0.89	0.09	31,31,31,31	1
3	ZN	B	307	1/1	0.92	0.08	36,36,36,36	1
2	ACT	B	301	4/4	0.92	0.10	11,14,16,17	4
2	ACT	A	301	4/4	0.92	0.10	11,15,16,17	4
3	ZN	A	309	1/1	0.93	0.08	30,30,30,30	1
3	ZN	A	305	1/1	0.95	0.05	17,17,17,17	1
3	ZN	A	311	1/1	0.95	0.07	39,39,39,39	1
3	ZN	A	312	1/1	0.95	0.07	36,36,36,36	1
3	ZN	B	305	1/1	0.95	0.07	15,15,15,15	1
3	ZN	B	309	1/1	0.96	0.05	21,21,21,21	1
3	ZN	A	303	1/1	0.97	0.05	19,19,19,19	1
3	ZN	B	303	1/1	0.97	0.06	16,16,16,16	1
3	ZN	B	308	1/1	0.98	0.04	17,17,17,17	1
3	ZN	B	304	1/1	0.98	0.04	22,22,22,22	1
3	ZN	A	304	1/1	0.99	0.03	15,15,15,15	1
3	ZN	A	306	1/1	1.00	0.03	14,14,14,14	1

6.5 Other polymers [i](#)

There are no such residues in this entry.